



UNIVERSITY OF  
FLORIDA

EENY-244

EXTENSION

Institute of Food and Agricultural Sciences

## Citrus Whitefly Parasitoid, *Encarsia lahorensis* (Howard) (Insecta: Hymenoptera: Aphelinidae)<sup>1</sup>

Ru Nguyen<sup>2</sup>

### Introduction

*Encarsia lahorensis* (Howard) is a specific parasitoid of the citrus whitefly, *Dialeurodes citri* (Ashmead). This parasitoid was discovered on citrus whitefly by R.S. Woglum, in 1911, while searching for natural enemies in India (Woglum 1913). Attempts were made to introduce *E. lahorensis* into Florida in 1911, however, cultures arrived in December when few susceptible host stages were available and the parasite was not able to survive (Woglum 1913). Apparently, no further attempts were made to introduce this parasite into Florida for the control of citrus whitefly until 1977.

In May 1977, the author transported *E. lahorensis* to Florida from California where it was established. Subsequently, it became established in Alachua and Polk counties, Florida (Nguyen and Sailer 1979). By the summer of 1979, *E. lahorensis* had suppressed the citrus whitefly population on viburnum and gardenia in those counties. Because of this success, 800 potted gardenia plants infested by citrus whitefly nymphs were exposed to *E. lahorensis*. After becoming parasitized these

whiteflies and plants were distributed to 66 of 67 Florida counties. Field surveys during 1981-82 indicated the presence of established populations of *E. lahorensis* in 59 counties (Sailer et al. 1984).

### Distribution

*Encarsia lahorensis* has been reported as native to India and Pakistan (Woglum 1913), and introduced to California in 1966 (Rose and DeBach 1981), Florida in 1977 (Nguyen and Sailer 1979), Georgia, Alabama, Texas, Louisiana, North Carolina (Wendel, personal communication), Italy (Viggiani and Mazzone 1978), and Israel (Rossler, personal communication) for controlling the citrus whitefly.

### Description and Biology

The female is small (0.54-0.84 mm long, 1.42 mm wing span), body white, head yellow, antennae with segments, funiculus segment longer than wide (3 times as long as wide), wing without shading in median. The male is a little smaller than female (0.62 mm long), and different in color, head brown, and abdomen dark brown (Howard 1911, Grissell 1979). Under experimental conditions (26±1°C, 70% Rh)

1. This document is EENY-244 (originally published as DPI Entomology Circular No. 290), one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: November 2001. This document is also available on Featured Creatures Website at <http://creatures.ifas.ufl.edu>. Please visit the EDIS Website at <http://edis.ifas.ufl.edu>. Additional information on these organisms, including many color photographs, is available at the Entomology and Nematology Department WWW site at <http://entnemdept.ifas.ufl.edu/>.

2. Ru Nguyen, Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville, FL.

longevity of the female is nine days; male, eleven days; and unfed adults, three days. Mated females deposited fertilized eggs in 3rd and 4th nymphal stages of *D. citri* and produced female offspring. Virgin females laid unfertilized eggs in the body of female fully- developed larvae or pupae of their own species (adelphoparasitic insect) and produced male offspring. Under laboratory conditions the development from egg to adult at 24°C required 12 to 15 days for males and 24 to 25 days for females (Nguyen and Sailer 1979, Viggiani and Mazzone 1978). *E. lahorensis* overwinters at various stages (larva, pupa). However, it has high mortality during winter in North Florida, especially for the first larval stage. The sex ratio changed from generation to generation. During the main period of adult emergence in March through April in Central Florida, the sex ratio favored females, but the number of males gradually increased later.



**Figure 1.** Citrus whitefly parasite, *Encarsia lahorensis* Howard, ovipositing in an immature citrus whitefly. Credits: University of Florida


### Host

*Encarsia lahorensis* is a specific parasitoid and *Dialeurodes citri* is the only host reported.

### Selected References

Grissell, E.E. 1979. The *Prospaltella* of Florida. Fla. Dept. Agri. Consumer Serv., Div. Plant Ind. Cir. 203: 1-4.

Howard, L.O. 1911. A note on the Indian enemies of *Aleyrodes citri* R. & H., with description



**Figure 2.** Top: immature larva of the parasite, *Encarsia lahorensis* Howard, in a larva of a citrus whitefly, *Dialeurodes citri* (Ashmead); middle: *E. lahorensis* pupa in citrus whitefly larva; bottom: emergence hole of an adult *E. lahorensis* in a dead citrus whitefly larva. Credits: University of Florida

of a new species of *Prospaltella*. J. Econ. Entomol. 4: 130-132.

Nguyen, Ru, and R.I. Sailer. 1979. Colonization of a citrus whitefly parasite, *Prospaltella lahorensis*, in Gainesville, Florida. Florida Entomol. 62: 59-65.

Rose, M., and P. DeBach. 1981. Citrus whitefly parasites established in California. California Agri. 35: 21-23.

Sailer, R.I., R.E. Brown, B. Munir, and J.C.E. Nickerson. 1984. Dissemination of the citrus whitefly (Homoptera: Aleyrodidae) parasite *Encarsia lahorensis* (Howard) (Hymenoptera: Aphelinidae) and its effectiveness as a control agent in Florida. Bull. Entomol. Soc. America 30: 36-39.

Viggiani, G., and P. Mazzone. 1978. Morfologia, biologia e utilizzazione di *Prospaltella lahorensis* How. (Hym.: Aphelinidae), a parassita esotico introdotto in Italia per la lotta biologica al *Dialeurodes citri* (Ashm.). Boll. Lab. Entomol. Agro. "F. Silvestri" Portici. 35: 99-161.

Woglum, R.S. 1913. Report of a trip to India and the Orient in search of the natural enemies of the citrus whitefly. USDA Bur. Entomol. Bull. 120: 1-58.