



UNIVERSITY OF
FLORIDA

ENY-472

EXTENSION

Institute of Food and Agricultural Sciences

Insect Management for Sweet Corn¹

G. S. Nuessly and S. E. Webb²

Foliar, ear and root feeding insects can routinely cause economic losses to sweet corn if left untreated. The most important pests of sweet corn in Florida are the fall armyworm, corn earworm, lesser cornstalk borer, cutworms, corn silk fly, cucumber beetles, aphids, and wireworms. Less common pests of sweet corn include grasshoppers, corn blotch leafminer, twospotted spider mites, sap beetles, stink bugs, maize weevils and billbugs, white grubs, and white fringed beetles.

Fall Armyworm, *Spodoptera frugiperda* (J.E. Smith)

Description

Adults (Figure 1) are light brown to ash gray with a 1 1/2 in. wing span. Several dark and light spots and lines are found on the front wings. The hind wings are lighter with a dark band near the margin. Eggs are deposited in masses of up to 200 eggs, often with more than one layer, and covered with scales from the moth's abdomen. Newly emerged larvae have black heads with all white bodies that become darker and patterned as they grow. Mature larvae grow to about 2 in. and are light green to tan or

brown to nearly black. Larvae have six dark bumps (tubercles) on the upper surface of each abdominal segment greater than or equal to the size of their spiracles (small, oval-shaped openings to the trachea on the lateral sides of most segments). The eighth abdominal segment has four distinct dark bumps on the top surface. The top of the first thoracic segment appears as a dark shield, often with three light stripes. Sutures on the front of head form an inverted "Y." Larvae (Figure 2) have three pairs of true legs and five pairs of prolegs. The 3/4 in. long pupae are dark reddish-brown.




Figure 1. Fall armyworm adult.

1. This document is ENY-472 (which replaces ENY-449), one of a series of the Entomology & Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: November 2001. Revised: July 2003. For more publications related to horticulture/agriculture, please visit the EDIS Website at <http://edis.ifas.ufl.edu/>.

2. G. S. Nuessly, Associate Professor, Crop Protection, Everglades REC, Belle Glade and S. E. Webb, Associate Professor, Entomology and Nematology Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 32611-0640. The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition. All chemicals should be used in accordance with directions on the manufacturer's label. Use pesticides safely. Read and follow directions on the manufacturer's label.




Figure 2. Fall armyworm

Biology

Adults feed on nectar and other moisture sources. Egg masses are usually deposited under leaves. Larvae emerge from eggs in 3 to 4 days. Prior to tassel push, young larvae feed between leaf veins often on outer portions of leaves before moving into the whorl to feed on young developing leaves. Later in the season young larvae first feed on tassels before moving to the ears or feed directly and complete development on ears. Last (6th) instar larvae leave the plants to pupate in shallow chambers beneath the soil surface. Larvae complete development within 2 to 3 weeks. Adults emerge from the pupal cases in 7 to 10 days to complete a generation in 24 to 35 days.

Damage

Long thin holes between the leaf veins indicate feeding by first instar larvae. Later instar larvae leave larger holes across veins. Leaves damaged by small to medium sized larvae within the whorl often emerge to display rows of holes. Feeding by mid to late instar larvae frequently results in a shredded appearance to the emerging leaves. Late instar larvae can completely consume the growing point of the plant, and can severely damage the tassel before it emerges from the whorl. Presence of these larvae can be detected by looking into the whorl for a plug of wet excrement (frass plug) blocking the opening. Tassels should be examined for early through mid instar larvae and their damage, including small holes, missing pieces, excrement and tassel pieces on the upper surface of leaves. Fourth through sixth instar larvae pushed from the whorl by the emerging tassel can quickly ruin the ears of at least one of the surrounding plants. All age larvae feed on ears. Young larvae feed on silk and often enter the tip of the ear to feed on silk before damaging kernels, and cob. Examine the silk for damage and presence of small frass. Mid to late instar larvae can enter the ear through the husk, bypassing the silk and causing direct damage to kernels and cob. The husks of ears

should be examined for direct entry, particularly in the protected area between the ear and plant stem.

Table 1. Fall armyworms

Management Option	Recommendation
Scouting/ Thresholds	Fields should be monitored at least weekly until tassel push for young larvae on leaves and in whorls. Thresholds used for treatment are plant age specific, with seedlings and plants pushing tassel often treated at 10% or lower infestations. Plants between these stages are usually treated when infestations reach between 15 and 20%. Fields should be scouted at least twice weekly during the ear stage to detect and control populations before they cause economic damage. Treatments during this period are usually initiated as soon as larvae are detected. Infestations above a few percent should elicit more frequent sampling. Pheromone traps can be used to monitor relative presence of the adults in the area.

Table 1. Fall armyworms

Management Option	Recommendation
Notes	It is important to eliminate the larvae being pushed from the whorl by the emerging tassel so they do not damage the sensitive emerging ear shoots or flag leaves and do not pupate within the field. Larvae completing development at tassel push can emerge as adults to lay eggs on the crop during the period of active silk growth. Chemical controls are most effective against the younger instars with higher rates and more frequent applications needed to try to control later instars. Granular formulations of several pesticides are available that reduce the exposure to non-target organisms by concentrating the pesticides in the whorls and leaf axils, particularly when applied in tight bands over the rows. Granular formulations are one of the most effective treatments available to kill larvae once they have formed a frass plug within whorls. The other pesticides are used as broadcast or banded sprays. Control of newly emerged larvae on ears is important to prevent economic damage.
Natural Enemies	Birds and insect natural enemies (predators and parasitoids) aid in fall armyworm control from germination through tassel push, but generally do not exert enough pressure to prevent yield loss in sweet corn. Birds can cause more damage than fall armyworms by feeding on kernels at ear tips within a week of harvest.
Resistant Varieties	Host plant resistance is emerging as an effective control strategy against FAW in sweet corns, including varieties enhanced with maysin and Bt genes that allow plants to produce their own natural control substances.

Corn Earworm, *Heliocoverpa zea* (Boddie)

Description

The front wing color is sex dependent. Female fore wings are yellow to pinkish-brown, while those of males are light greenish-brown. Both sexes usually have a dark spot in the middle of the fore wing (Figure 3). Both fore and hind wings have dark margins and the hind wings have a short, narrow band near the middle and two pale spots near the edge of the wing in the dark margin. Both sexes have a wingspan of 1 1/2 in. The ball shaped, white to yellow to green eggs have ridges running from top to bottom. Newly emerged larvae are translucent white to yellow with a light brown head. Older larvae vary considerably from dark yellow to green or tan to dark brown. Mature larvae can reach 2 in. Larvae (Figure 4) have three pairs of true legs and five pairs of prolegs. The skin (cuticle) of larvae are covered with microspines (need good hand lens or microscope to see microspines). The dark colored tubercles (larger bumps) on abdominal segments 1, 2 and 8 do not have microspines more than 1/4 the way to their apex. The central area on the inside surface of the mandibles do not have a separate toothed area (observed with microscope after spreading mandibles with pins or forceps). The 3/4 in. long pupae are reddish-brown in color.




Figure 3. Corn earworm adult.




Figure 4. Corn earworm.

Biology

Corn earworm moths feed on nectar and other moisture sources. Females deposit their eggs singly or in small groups of less than five on leaves and ears. The eggs are not protected by scales. Larvae emerge from eggs in 3 to 4 days to feed in corn whorls or ears, completing development in 18 to 26 days. Larvae leave plants to pupate within the soil. Adults emerge in 7 to 10 days. Corn earworms complete their development from egg to adult in 28 to 40 days.

Damage

Young larvae feed between leaf veins often on outer portions of leaves before moving into the whorl to feed on young, developing leaves. Larger larvae can completely consume the growing point of the plant and severely damage the tassel before it emerges from the whorl. More commonly it attacks the ears causing severe damage to silk, kernels and cob. Twenty years ago this was the primary pest of corn throughout Florida. Fall armyworm has largely replaced it as the primary pest, particularly in central and southern Florida, but corn earworm still causes severe damage in corn fields throughout the state. Corn grown in northern Florida counties during the late spring and summer is more likely to have large populations of corn earworm.

Table 2. Corn earworms

Management Options	Recommendation
Scouting/ Thresholds	Fields should be monitored at least weekly until tassel push for young larvae on leaves and in whorls. Thresholds used for treatment are plant age specific, with seedlings and plants pushing tassel often treated at 10% or lower infestations. Plants between these stages are usually treated when infestations reach between 10 and 20%. Fields should be scouted at least twice weekly during the ear stage to detect and control populations before they cause economic damage. Treatments during this period are usually initiated as soon as larvae are detected. Infestations above a few percent should elicit more frequent sampling. <i>Helicoverpa</i> pheromone traps can be used to monitor relative presence of the adults in the area.
Notes	It is important to eliminate the larvae being pushed from the whorl by the emerging tassel so they do not damage the sensitive emerging ear shoots and flag leaves and do not pupate within the field. Earworm larvae completing development at tassel push can emerge as adults to lay eggs on the crop during the period of active silk growth. Insecticides are most effective against the younger instars with higher rates and more frequent applications needed to try to control later instars. Granular formulations of several pesticides are available that reduce the exposure to non-target organisms by concentrating the pesticides in the whorls and leaf axils, particularly when applied in tight bands over the rows. The other pesticides are used as broadcast or banded sprays. Control of newly emerged larvae on ears is important to prevent economic damage.

Table 2. Corn earworms

Management Options	Recommendation
Natural Enemies	Birds and insect natural enemies aid in earworm control, but generally do not exert enough pressure to prevent yield loss, particularly in sweet corn. Birds can cause more damage than corn earworms by feeding on kernels at ear tips within a week of harvest.
Resistant Varieties	Host plant resistance is emerging as an effective control strategy against corn earworms in sweet corns, including varieties enhanced with Bt genes that allow plants to produce their own natural control substances.


Lesser Cornstalk Borer, *Elasmopalpus lignosellus* (Zeller)

Description

The adults are narrow-winged moths 1/2 to 5/8 in. long (Figure 5). The body and wings of female moths are covered with mostly gray to brown and reddish shiny scales. Males are pale yellow to medium brown with wings bordered with a band of darker scales. Adults fly quickly when disturbed in the field, often landing on the soil several yards away. Males are easily spotted against the darker organic soils. The flat, shingle-like eggs are creamy green and become pink to red at the time of larval emergence. Larvae are patterned with alternating narrow transverse bands of maroon to brown and aqua blue over a cream base. Larvae reach 1/2 to 5/8 in. long at maturity. Larvae have three pairs of true legs and five pairs of prolegs. Larvae are very active when disturbed. Pupae change color from greenish to brown with time and are surrounded by a flimsy silken cocoon.

Biology

Lesser cornstalk borer moths feed on nectar and other moisture sources. Eggs are deposited singly or in small groups on the soil surface or directly on the stem near the soil surface. Temperature greatly affects development. Larvae emerge in 18 days at 64° F, but in less than 3 days at 91° F. They hide

**Figure 5.** Lesser cornstalk borer.

and feed at base of corn stalks from narrow soil-covered silken tubes 1/4 to 1/2 in. beneath the soil surface. Larvae complete development in 17 to 42 days. The larvae pupate within the soil and adults emerge in 10 to 24 days. Complete generation time is 30 to 84 days.

Damage

Larvae bore into the corn stem of young plants just below the soil surface and feed up and down the stem. They also bore into larger brace roots of nearly mature plants. Young plants exhibit severe wilting of the youngest two to three leaves. Plants that do not die from this damage often develop several suckers that are bushy and stunted, but they do not produce marketable ears. While wireworms produce similar damage symptoms, corn plants attacked by these beetle larvae are much less likely to survive to produce suckers as when they are attacked by lesser corn borer larvae. Infestations can cause severe stand loss usually in irregular shaped patterns in the field. Damage is usually more severe during dry, warm periods. Larvae may move down a row to kill several seedlings before completing development.

Table 3. Lesser cornstalk borer

Management Options	Recommendation
Scouting/ Thresholds	Control efforts can be maximized by estimating the potential for damage based on adult monitoring, surrounding crops and previous field history. Delta wing pheromone and black light traps can be used to monitor adults. Adults are easily visible in fields as they are quick to fly as one approaches. Carefully examine the soil immediately around the base of the injured plants for soil-covered silken feeding tubes loosely attached to the entry hole at plant base or brace roots.
Notes	They tend to be worse in drier years and on well-drained soils. However, they can cause severe damage to sweet corn on heavy organic soils planted throughout the year. Larvae may already be present in the field when seeds are planted. Wet, cool growing conditions increase their mortality and greatly reduce their developmental rates. In fields with a high potential for infestation, at-plant pesticide application should be considered to reduce damage. Unfortunately, by the time field infestations are observed in corn, damage to the crop has already occurred. Post-emergence applications applied as soon as damage symptoms appear can limit additional damage, but may be too late to prevent economic losses. Post-emergence applications work best when banded over the rows and lightly incorporated into the soil around the plants.
Natural Enemies	Natural enemies are not thought to have a significant effect on lesser cornstalk borers due to their subterranean habits, silken feeding tubes and sporadic nature.

Table 3. Lesser cornstalk borer

Management Options	Recommendation
Resistant Varieties	Host plant resistance is probably the best strategy for control of this borer in the future. Resistance has been noted in lines tested at CIMMYT in Mexico and domestic sweet corn varieties enhanced with a Bt toxin provided overall control several times greater than that of insecticides.
Cultural Controls	Field cultivation several weeks in advance of planting is advised to allow for decomposition of the plant residues and completion of borer development. Planting corn following corn, sorghum, small grains, sugar cane, beans, peanuts, or too quickly behind recently turned weedy fields can predispose a crop to damage. Lesser cornstalk borer does very well on nutsedge and corn planted behind sugarcane infested with these weeds can suffer severe damage.

Cutworms

Description

Cutworm adults are large bodied moths with various dark patterns over light to medium brown or gray base color of fore wings. Shapes and coloration of spots on front wings are diagnostic in the identification of adult cutworm species. The wing spans of both the black (*Agrotis ipsilon* (Hufnagel)) and variegated (*Peridroma saucia* (Hubner)) cutworms ranges from 1 5/8 to 2 1/8 in, while the wings of the granulate cutworm (*Feltia subterranea* (F.)) are slightly narrower at 1 1/4 to 1 3/4 in. Male and female black cutworm moths (Figure 6) colored differently. However, in both sexes the kidney-shaped (reniform) spot is a dark crescent-shaped ring with a dark line emanating outward and hind wings are whitish to gray with dark scales on veins. The claviform spot (proximal to and beneath reniform spot) is an elongate oval, and the orbicular spot (proximal to reniform spot) is small and round to tear-shaped. Wing color in granulate

cutworms (Figure 7) varies greatly. Both sexes are recognized by the black bar between the usually pale, basal, round (orbicular) and kidney-shaped (reniform) spots on the fore wings. Hind wings are off-white with veins and margins sometimes darker. There is less distinction between the sexes in wing coloration for variegated cutworms (Figure 8). The kidney-shaped reniform and nearly round orbicular spots (proximal to reniform spot) in the central fore wing are both large and ringed with black. The rear half of the forewings is clay-colored with blackish spots. Hind wings are iridescent to pearly white with brown veins and margins.




Figure 6. Black cutworm adult.




Figure 7. Granulate cutworm adult.




Figure 8. Variegated cutworm moth.

Eggs of all three species are whitish and become variegated brown with age and have ribs radiating downward from the top. Black cutworms eggs are spherical shaped, those of granulate cutworms are

rounded with flat bottoms, and variegated cutworms eggs are rounded with rounded bottoms.

All cutworm larvae have sutures on the front of the head that form an inverted "V" and all have three pairs of true legs and five pairs of prolegs. Black cutworm larval color ranges from light gray to black on top and lighter below (Figure 9). Two lighter stripes running down the center of the back are visible on lighter specimens, but all larvae lack an obvious band down their back. Two rows of small, black, raised bumps (tubercles) run down the back; two per segment. Their cuticle is covered with large and small convex, isolated granules (need hand lens or microscope to see granules). Their head is brownish with many dark spots. Granulate cutworm larvae (Figure 10) are gray to reddish brown with dull yellowish subdorsal markings on the abdominal segments. The heads are yellowish to brownish. The undersurface of the body is flecked with white. Granules on their cuticle are isolated, bluntly conical, and project slightly backward. Body color in variegated cutworms (Figure 11) is brownish gray to nearly black on top of back and lighter below. A row of whitish to yellow circular spots run down the center of the back, particularly on the first four abdominal segments. The cuticle is smooth without granulation. An orange to brown line may connect the spiracles, below which there is usually some yellowish to orangish coloration. The head is orange-brown and marked with darker spots. The pupae of all species are reddish brown to brown, 5/8 to 7/8 in. long and are found in the soil.




Figure 9. Black cutworm larvae.




Figure 10. Granulate cutworm larva.




Figure 11. Variegated cutworm larva.

Biology

Cutworm moths feed on nectar and other moisture sources. Moths begin depositing eggs on field debris, stubble or leaves near the soil surface 7 to 10 days after emergence. Black cutworm eggs are deposited singly or in groups of up to 30, granulate cutworms eggs are deposited singly or in small groups and variegated cutworms deposit masses of up to several hundred eggs. Larvae emerge from eggs in 3 to 6 days. Larvae tend to curl up into a ring when disturbed or handled. They may also bite and release a greenish-brown fluid. They feed on leaves and stems of mostly young plants. Older larvae (4th instar and later) can reach 2 in. in length and can cut plants off at their bases and drag them to their burrow in soil. Larvae complete development in 20 to 40 days. Larvae pupate within a chamber in the soil. Adults emerge in 10 to 20 days. Generation time for cutworms is 43 to 76 days, depending on temperature.

Damage

These cutworms are pests of corn and most other vegetables throughout Florida. Damage includes leaf feeding and stand loss due to cutting off entire plants. Black cutworms do most of their feeding at ground level. Larvae feed on young plants, cutting off leaves, or in later instars, entire plants. Populations of this pest tend to be higher in weedy and in wet fields. Granulate cutworm larvae can cut off entire seedling plants, as well as climb and feed on leaves of older plants. This cutworm is not associated with weedy fields as is the black cutworm. First instar larvae stay on plants, while older larvae climb and feed on plants only during night. Variegated cutworm larvae cut off seedlings at ground level or defoliate older plants.

Table 4. Cutworms

Management Options	Recommendation
Scouting/ Thresholds	Seedling corn should be scouted as frequently as twice per week to detect cutworms or their damage, particularly in areas known for this pest. Young larvae may be found grouped together on foliage, but older larvae will usually be found singly in soil or beneath leaf trash during the day. Adults can be monitored with black light and pheromone traps.
Notes	Larvae may already be present in the field when seeds are planted. Pesticides are available for at-plant, pre- and post-emergence broadcast and banded applications. Post emergence applications are the most efficient.
Natural Enemies	Natural enemies such as parasitic wasps, flies and predacious ground beetles can exert tremendous control pressure that may approach 80%. Larvae are also targets for attack by pathogenic fungi and viruses. However, seedlings emerging in fields without resident natural enemy populations can experience significant stand loss from first generation cutworms.
Cultural Controls	Weedy fields quickly rotated to corn have higher potential for stand loss due to older larvae cutting off the emerging plants.

Corn Silk Fly, *Euxesta stigmatias* Loew

Description

These 3/8 to 5/16 in. long flies (Figure 12) have a dark green body with a slight metallic sheen, normal length black legs (not long like the predacious long-legged flies), and red to brown eyes. Their wings are patterned with four dark horizontal bands. The end of the male's abdomen is rounded while that of the female is terminated by a trapezoidal segment from which a long ovipositor is extended during egg deposition. Thin, 1/16 in. long white eggs are

deposited in silk channels, between ends of husks and around armyworm and earworm entrance and exit holes in husk of corn ears. The larvae (Figure 13) are white to pale yellow, narrow, legless maggots, reaching 3/8 to 7/16 in. long, with black mouth hooks. Maggots do not molt out of their last cuticle, but complete metamorphosis within it. The resulting delicate puparium is light to medium brown.




Figure 12. Corn silk fly adult. Credits: J. F. Butler




Figure 13. Corn silk fly larva.

Biology

This insect is saprophytic and feeds and reproduces on a wide variety of over-ripe and rotting fruits, vegetables, and sugarcane. Adults lap up fluids and pollen with their extendable mouth parts. Females will deposit eggs into sweet corn ears for at least 3 weeks after silk initiation, but seem to prefer ears with fresh silk. Adults are active runners on plants often performing elaborate behaviors involving wing flapping and running at and around other corn silk flies. Mated females are most often observed on ears below the overhanging silks. Larvae emerge from eggs in 2 to 4 days and start feeding on silks just inside the silk channel. Many larvae complete development on silks without causing any direct injury to kernels. But those that continue into the ear feed extensively on kernels at the tips of the ear, or may disperse randomly throughout the ear to feed on kernels. Larvae complete development in 15 to 21

days. Exposed larvae quickly seek shelter by crawling or flicking themselves from the ears by grabbing and quickly releasing the end of their abdomen with their mouthhooks. The great majority of larvae leave the ears to pupate within the soil, but a few pupate in the silk near the opening to the silk channel. Adults emerge from their puparia in 7 to 8 days to complete a generation in 25 to 32 days.

Damage


The geographic range of this insect has spread up the Florida peninsula during the last decade from the Homestead region. This fly is a year round pest of corn in southernmost Florida. Corn grown just south of Lake Okeechobee is attacked primarily in the late winter and spring. Corn in central Florida is attacked in late spring and early summer. North Florida corn is prone to attack during the summer. This pest damages ears in several ways. By damaging silks, the larvae disrupt pollination and reduce kernel density. Larval feeding on silks and at ear tips that doesn't result in kernel damage can still result in throwing the ears out of grade. Damaged silks in this moist environment turn golden to medium brown and soon begin to ferment. This can lead to secondary infestations of sap (picnic) beetles that are drawn to the released volatiles from the rotting silks. However, near-mature larvae are just as likely to damage individual kernels distributed throughout ears rendering them completely unmarketable. In south Florida, and other areas with appropriate local food reservoirs, these flies are quick to reenter treated fields. Therefore, damage along field margins and across fields with large field edge to acreage ratios can be substantial.

Table 5. Corn silkfly

Management Options	Recommendation
Scouting/ Thresholds	Adults are usually found on the tassels and upper leaves early in the morning and late in the afternoon. This is the best time to scout and to control them with chemical insecticides. They move down the plants, or at least into shaded parts of the plants, during the day.

Table 5. Corn silkworm

Management Options	Recommendation
Notes	Organophosphate materials have the quickest activity and best residue against this pest. Pyrethroids work well on contact, but their residues may only cause temporary knock down, followed by recovery of adults.
Natural Enemies	Web and non-web spinning spiders feed on adults. Fire ants, predacious ground beetles, earwigs and spiders feed on larvae on the soil surface.
Resistant Varieties	While cultivars with high maysin content impart partial resistance to this insect, no commercial varieties are currently available with this compound.
Cultural Controls	Avoid planting to fields surrounded by other host crops, such as tomato or peppers, particularly if it is possible that these crops will be abandoned during the silking to harvest period of sweet corn.

**Figure 14.** Banded cucumber beetle.

Cucumber Beetles

Description

Adult banded cucumber beetles (*Diabrotica balteata* LeConte) (Figure 14) are small (1/4 to 3/8 in. long) and oval-shaped with a dirty yellow to dark green thorax and abdomen. The yellow transverse bands across the wing covers (elytra) and red to reddish-brown heads serve to separate them from the southern or spotted root worm adults (*Diabrotica undecimpunctata howardi* Barber) which have 12 black spots arranged in 3 rows across their elytra. The third tarsal segment on all legs in both species is laterally swollen into two lobes with the last (fourth) tarsal segment originating from between the lobes. Both species produce small, oval-shaped, pale yellow eggs. The white, elongate and soft bodied larvae have three pairs of minute legs. Larvae are indistinguishable between the two species. The white to pale green pupae are found within loosely woven cocoons in the soil.

Biology

Adults feed on young corn leaves, anthers and silk. Adults enter the soil to deposit eggs in clusters of up to 12 near roots. Larvae emerge in 6 to 8 days and feed on roots and other soft tissue at the base of the stalks. Larvae complete development in 15 to 22 days feeding on corn roots. Adults emerge in 16 to 20 days. Banded cucumber beetles complete development in 35 to 50 days.

Damage

Cucumber beetles are found throughout Florida, with the banded species more common in central and south Florida and the spotted one more common in northern Florida. Larvae damage roots and can enter the corn stalk just above the roots where they can eat the crown and kill the buds of young plants. Bacterial wilt and other pathogens can enter the plant through these wounds. Severe root damage can result in lodging. However, estimates of yield effects for larval or adult feeding have not been determined for Florida corn. Adults are common throughout the winter months in south Florida. While preferring to feed on

weeds in the genus *Amaranthus* (pig weeds), adults can cause serious foliar damage to corn and other vegetables. Adult feeding produces irregular shaped, often elongate holes in young leaf tissue within the whorls. Delayed growth, plant stunting, and stand loss can result from heavy feeding damage to whorls of seedlings. Feeding damage to older leaves is evidenced by notched leaves, window paning, and irregular patches with only the veins remaining. Silk pruning by adults can result in blanks within the ears. Adults are good fliers and can invade and damage fields quickly.

Table 6. Cucumber beetles

Management Options	Recommendation
Scouting/ Thresholds	Seedling and silking stage plants are most sensitive to adult feeding. Look for adults, feeding damage and thread-like or oily spots of fecal material at feeding sites while scouting for army- and ear-worms.
Notes	Chemical control of adults is through contact or bait insecticides. This latter type selectively treats the beetles as they eat the baits. Numerous pesticides are labeled for mostly at-plant treatment of cucumber beetle larvae.
Natural Enemies	Several predacious insects and spiders prey on these insects, but migration of large numbers can quickly overwhelm the natural enemy complex.
Cultural Controls	Control of cucumber beetle infested weeds on ditch banks and surrounding fields should be timed so that beetles are not forced to look for food during sensitive periods in sweet corn development.

Wireworms or Click Beetles

Description

The adult stage (Figure 15) of this insect is a slender, somewhat flattened, medium to dark brown or gray beetle between 1/2 and 7/8 in. long. The exoskeleton is smooth or with very short hairs. They have a large tooth-like projection between the rear legs that fits into a groove on the undersurface of the

abdomen. These beetles feign death when disturbed and can then right themselves from their backs by quick flexion at the juncture of the thorax and abdomen. This behavior produces a clicking sound in some species of click beetles. The larvae are called wireworms (Figure 16) and have narrow, hardened, creamy yellow to orangish-brown, tubular bodies. Characteristic hardened projections on the next to last abdominal segment can be used to identify them to species. They have three pairs of short true legs and no prolegs and can reach 1 1/4 in. long. Pupae are not formed within a cocoon and their legs, antennae and wing buds are completely visible.




Figure 15. Adult stage - Click beetle.




Figure 16. Corn wireworm larva.

Biology

Adults feed on living and decaying plant material and are frequently observed in corn leaf axils and whorls during the late spring. Eggs are deposited in soil near plant roots, particularly around grasses, during summer months. Wireworms take two to several years to complete larval development. The pupal stage is passed in the soil from which adults emerge within 2 weeks.

Damage

Larvae attack seeds, roots, and crowns of plants below the soil surface. They chew into the base of plants and then hollow out the stem, eliminating the growing points of young plants. Young plants first exhibit severe wilting and desiccation of the youngest leaves. Such damage to young plants results in stand loss. Wireworm damage symptoms are nearly identical to those of lesser cornstalk borer, although young plants damaged by the latter are much more likely to send up suckers than those damaged by wireworms. Wireworms also do not form silken feeding tubes as do lesser cornstalk borer larvae. Lodging results following strong winds and rains when roots have been pruned or damaged by larval feeding.

Table 7. Wireworms

Management Options	Recommendation
Scouting/ Thresholds	Soil within 8 in. diameter around affected plants should be excavated in search of larvae in or near these plants. If infestation density is in question, then fields should be sampled for these pests to determine the best treatment strategy before corn is planted. Baits of oat, corn or potato buried in fields and recovered in 2 to 3 weeks work well to monitor wireworms. Average counts greater than 2 per bait are enough to recommend treatment.
Notes	Pesticides are available for pre-plant broadcast and at-plant banded applications. Soil moisture plays an important role in releasing insecticides from their carriers, consequently wireworm control in dry soils is more difficult. Pre- or post-plant irrigation of dry soils and adequate mixing of insecticides with the soil around and in the seed furrow will help to improve control.
Natural Enemies	Birds such as cattle egrets that follow farm equipment through the field eat many wireworms exposed during field disking.

Table 7. Wireworms

Management Options	Recommendation
Cultural Controls	Corn planted to weedy fields, or in rotation from pasture or sugarcane, are often exposed to very large wireworm populations. Summer flooding of fields is an effective cultural control, providing it is maintained for at least several weeks. Water temperature should be above 82° F for maximum control. Rotation through a cycle of rice has been shown to eliminate the need for wireworm treatment in the subsequent crop.

Aphids

Description

The two most commonly encountered aphids in sweet corn are the birdcherry oat aphid, *Rhopalosiphum padi* (L.) (Figure 17), and the corn leaf aphid, *R. maidis* (Fitch) (Figure 18). Both species are small to medium sized (3/32 in.), elliptical- to pear-shaped aphids. Aphids have a pair of short, tube-like structures (cornicles) that extend backward and upward from near the end of the top surface of the abdomen. Wingless adult birdcherry oat aphids are medium green with reddish areas toward the end of the abdomen and area around bases of cornicles. Wingless adult corn leaf aphids are light to medium green with short, dark green to black legs and cornicles, but lack the reddish areas around the cornicles as in birdcherry oat aphids. Nymphs of both species are similar in color and shape to adults, only smaller. Winged adult forms of both aphids have black heads and thoraxes, and green abdomens.

Biology

Adult and immature aphids feed on phloem sap sucked from leaves, stems, tassels, and husks through their long piercing-sucking mouth parts. Excess water and sugars are voided from the body. These honeydew deposits on leaves serve as a source of food for sooty mold fungi, ants, sap beetles, and flies. Winged or wingless adult females deposit live nymphs, skipping the external egg stage in Florida. Adults produce from 3 to 5 nymphs per day for up to




Figure 17. Birdcherry oat aphid.




Figure 18. Corn leaf aphid.

several weeks. This allows for rapid population development. The nymphs pass through several instars before molting into adults in 7 to 10 days. There is no pupal stage in aphids.

Damage

Aphids have become a greater problem for spring sweet corn during the last several years. This may be the result of reduced pesticide inputs due to lower fall armyworm and corn earworm populations. Honeydew from aphids makes husks sticky and encourages sooty mold growth that can result in husk discoloration and grade reduction. Large populations of corn leaf aphid feeding on the emerging tassels of many plants within an area can result in poor pollen shed and fertilization. Both of these aphids can also infest sweet corn ears where their control is difficult due to feeding in concealed locations beneath dried silks, under husk wrappers and between ears and stalks.

Table 8. Aphids

Management Options	Recommendation
Scouting/ Thresholds	Look for aphids (particularly corn leaf aphids) within the corn whorls, on husks beneath silk or in region between ear and stalk near ear base. Insecticide applications for fall armyworm and corn earworm control often kill aphids before they can develop to damaging densities.
Notes	Aphid populations are easier to control while still infesting the whorl. Aphids feeding on husks are somewhat protected from chemical control strategies by the corn canopy and silks. Pesticides are available for post-emergence foliar treatments. Birdcherry oat aphids are less susceptible than corn leaf aphids to pesticides frequently used for armyworm control. Pyrethroids have shown good control of aphids within the whorl in pre-tassel stage sweet corn.

Table 8. Aphids

Management Options	Recommendation
Natural Enemies	Many types of natural enemies may control these aphids under low pesticide input situations, such as in field or silage corn. However, these aphids appear in sweet corn at a time when pesticide use increases for armyworm, thereby greatly reducing the potential impact of their predators and parasitoids. Pathogens kill many winged adults before they begin producing nymphs.

Table 9. Selected insecticides approved for use on insects attacking sweet corn.

Chemical Name	REI (hours)	Days to Harvest	Insects	Notes
*Ambush 2EC (permethrin)	12	1	aster leafhopper, corn earworm, corn rootworm (adults), cutworms, fall armyworm, southern armyworm	Do not apply more than 2.0 lb active ingredient per acre per season.
*Asana XL 0.66EC (esfenvalerate)	12	1	aphids, armyworms, banded cucumber beetle, beet armyworm (aids in control), chinch bugs, corn borer, corn earworm, corn rootworm, corn silk fly, cutworms, flea beetles, grasshoppers, sap beetles (adults), stalkborers, tarnished plant bug	
Avaunt (indoxacarb)	12, (14 days for hand harvesting)	3, (35 for fodder & storage)	European corn borer, fall armyworm	Whorl application only. No more than 4 applications per season.
*Baythroid 2 (cyfluthrin)	12	0	chinch bugs, common stalk borers, corn earworm, corn rootworm adult, cutworms, fall armyworm (1st and 2nd instars only), grasshoppers, silk fly, true armyworm	Maximum number of applications: 10.
Bt (<i>Bacillus thuringiensis</i>)	4 (most)	0	armyworms, beet armyworm, corn borers, fall armyworm, loopers, southern armyworm	

Table 9. Selected insecticides approved for use on insects attacking sweet corn.

Chemical Name	REI (hours)	Days to Harvest	Insects	Notes
*Capture 2 EC (bifenthrin)	12	1 - foliar 30 - soil application	aphids, armyworms, aster leafhopper, chinch bugs, common cornstalk borer, corn earworm, corn rootworm adult, corn silk fly, cutworms, cucumber beetle adult, grubs(1), mites, sap beetles, seed corn beetle(1), seed corn maggot(1), tarnished plant bug, webworms, wireworms(1)	Prohibited in FL coastal counties. (1) Soil application. Do not apply more than 12.8 ounces of product per acre per season.
*Counter 20 CR; Lock 'n Load 15 G (terbufos)	48	60	billbugs, chinch bugs(1), corn rootworm, cutworms (suppression), flea beetles, lesser corn stalk borer (suppression), maize billbug, seedcorn beetle, seedcorn maggot, symphylans, thrips, white grubs, wireworms	(1) Early season control of light to moderate infestations.
*Diazinon 4E, *50W (diazinon)	24	7 or preplant	corn earworm, corn leaf aphid, corn rootworm adult, cutworms, flea beetles, grasshoppers, sap beetles, seed corn maggot, spider mites, wireworms	Apply just before planting for seed corn maggot, cutworms, and wireworms. Do not make more than 5 applications per season (4E).
*14G, *5G	24	preplant only	cutworms, seed corn maggot, wireworms	
Extinguish ((S)-methoprene)	4	0	fire ants	Slow-acting IGR (insect growth reculator). Best applied early spring and fall where crop will be grown. Colonies will be reduced after three weeks and eliminated after 8 to 10 weeks. This is the only fire ant bait that is labeled for use on cropland. May be applied by ground equipment or aerially.
*Force 3G (tefluthrin)	0	at planting or cultivation within 30 days of seeding emergence	billbugs, chinch bugs, corn rootworm, cornstalk borers, cutworms, lesser cornstalk borer, red imported fire ant, seedcorn beetle, seedcorn maggot, white grubs, wireworms	Only one application per crop. Granules must be incorporated.
*Furadan 4F (carbofuran)	48, soil foliar app., 14 days	at planting, 7 foliar (machine harvesting only)	corn rootworms, flea beetles, seedcorn maggot, wireworms	See restrictions for very sandy soil. No more than 2 foliage applications, one if applied to soil also.

Table 9. Selected insecticides approved for use on insects attacking sweet corn.

Chemical Name	REI (hours)	Days to Harvest	Insects	Notes
*Lannate LV, *SP (methomyl)	48	0 = ears 3 = forage 21 = stover	aphids, armyworms, beet armyworm, corn earworm, corn rootworm, cutworms, fall armyworm, flea beetles, picnic beetles	Certain hybrid varieties are susceptible to methomyl injury. Treat a small area to determine safety first.
*Larvin 3.2 (thiodicarb)	48	0	beet armyworm, cutworms, corn earworm, fall armyworm, southern armyworm	Do not allow livestock to graze treated field. Do not feed treated corn silage or fodder to livestock. See label for special instructions for cutworms.
*Lorsban 4E (chlorpyrifos)	24	21 7, if in accordance with SLN FL940003	aphids, beet armyworm, chinch bugs, corn earworm, corn rootworm adult, cutworms, fall armyworm, grasshoppers	Do not feed treated corn silage, forage, or fodder, or allow livestock to graze (if used according to supplemental label).
15G	24	at planting	corn rootworm larvae, cutworms, lesser corn stalk borer, seed corn maggot, wireworms	See label. Do not use in conjunction with Lorsban 4E.
Malathion 57 EC, *8 EC (malathion)	12	5	aphids, armyworms, budworms, cereal leaf beetle, corn rootworm, grasshoppers, sap beetles, thrips	Injury may occur in whorl and silk stages.
*Metasystox-R 2EC (oxydemeton-methyl)	48	7 = 1 appl or 21 = 2 or 3 appl	aphids, corn rootworms, leafhoppers, mites, thrips	Do not apply more than 3 times per season.
Mocap 10G (ethoprop)	48	at planting	corn rootworms, cutworms, symphylans, wireworms, (suppression of white grubs)	
Neemix 4.5 (azadirachtin)	12	0	aphids, armyworms, corn earworm, thrips	
Oil, insecticidal	4	0	aphids, armyworms, corn earworms, corn rootworms, mites, thrips	
*PennCap-M 2EC (methyl parathion)	4 days See label	4	aphids, armyworms, corn earworm, corn rootworm adult, cutworms, flea beetles, grasshoppers, sap beetles, silk fly	See restrictions on label.
Phaser 3EC, 50WP (endosulfan)	24	1	aphids, corn earworm, whiteflies	Do not apply to sweet corn to be processed or used to feed livestock.

Table 9. Selected insecticides approved for use on insects attacking sweet corn.

Chemical Name	REI (hours)	Days to Harvest	Insects	Notes
*Pounce 1.5 G (permethrin)	12	at planting	armyworms, cutworms	
		pre-emergence	armyworms, cutworms, stalk borers	
		foliar - 1	armyworms, corn borers, cutworms, stalk borers	
Pyrellin EC (pyrethrin + rotenone)	12	12 hours	aphids, flea beetles, leafhoppers, loopers, mites, thrips	
Sevin 80S; 4F; XLR (carbaryl)	12	2 - Ears 14 - Forage 48 - Fodder	armyworms, chinchbugs, corn earworms, corn rootworm adult, cutworms, fall armyworm, flea beetles, leafhoppers, sap beetles	Highly toxic to bees.
Sevin 5 Bait (carbaryl)	12	2 - Ears 14 - Forage 48 - Fodder	ants, crickets, cutworms, grasshoppers, mole crickets, sowbugs	
Soap, insecticidal	12	0	aphids, armyworms, leafhoppers, mites, thrips	
SpinTor 2 SC (spinosad)	4	1	armyworms, corn earworm	
Spod-X LC (insect virus)	4	0	beet armyworm	Treat when larvae are young.
*Telone C-35 (dichloropropene)	5 days - See label	preplant	symphylans, wireworms	See supplemental label for use restrictions in south and central Florida.
*Thimet 20G (phorate)	48	at planting, see label	corn rootworms, flea beetles, mites, seedcorn beetle, seed corn maggot, white grubs, wireworms	
*Warrior (lambda-cyhalothrin)	24	1 21 for feeding livestock	aphids, aster leafhopper, beet armyworm, chinch bugs, corn earworm, corn rootworm, cutworms, fall armyworm, flea beetles, grasshoppers, mites (see label for more details), red imported fire ant(1), southern armyworm, tarnished plant bug, wireworms(1)	(1) Suppression only.
The pesticide information presented in this table was current with federal and state regulations at the time of revision. The user is responsible for determining the intended use is consistent with the label of the product being used. Use pesticides safely. Read and follow label instructions.				
* Restricted Use Only.				