UNIVERSITY OF
FLORIDA

EENY-232

EXTENSION

Institute of Food and Agricultural Sciences

Palmetto Tortoise Beetle, Florida Tortoise Beetle (unofficial common names), *Hemisphaerota cyanea* (Say), (Insecta: Coleoptera: Chrysomelidae: Cassidinae)¹

D. W. Hall and J. F. Butler²

Introduction

The Florida tortoise beetle, *Hemisphaerota cyanea* (Say), is a beautiful small beetle on a variety of native and exotic palms on which it occasionally inflicts damage by its feeding activities. It is our only tortoise beetle that feeds on palms.

Distribution

The Florida tortoise beetle is recorded from Florida, Georgia, Alabama, and Texas, and also probably occurs in Louisiana and Mississippi in areas where palms grow.

Description

The adult beetle is dark blue to purple in color. It has alternating longitudinal rows of pits and ridges on the elytra. Its hemispherical shape is the basis for its genus name. The antennae are yellow except for the basal segment which is black. The tarsi are greatly enlarged and modified for grasping the substrate.

Each of the greatly enlarged tarsi is equipped with approximately 10,000 adhesive bristles. Each bristle has two terminal pads. When walking, only a few of the bristles touch the leaf surface. However, when attacked by a predator, the beetle puts all or nearly all of the bristles in contact with the surface and secretes oil onto the pads. With the adhesive force created by the oil between the leaf surface and tarsi, the beetle is able to clamp its hemispherical shell down tightly against the leaf and has been demonstrated to withstand pulling forces of approximately 60 times its own weight for up to two minutes. This time period is sufficient to thwart the efforts of predatory ants attempting to pry the beetle from the leaf. The oils used are similar to those of the cuticle of insects and are not believed to be unique for the defensive function. The wheel bug, *Arilus cristatus* (L.) (Hemiptera: Reduviidae) is able to overcome the defenses of the beetle -- possibly by injecting toxic saliva to kill the beetle.

-
1. This document is EENY-232, one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: August 2001. This document is also available on Featured Creatures Website at <http://creatures.ifas.ufl.edu>. Please visit the EDIS Website at <http://edis.ifas.ufl.edu>. Additional information on these organisms, including many color photographs, is available at the Entomology and Nematology Department website at <http://entnemdept.ifas.ufl.edu/>.
 2. D. W. Hall, professor and J. F. Butler, professor, Entomology and Nematology Department, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.

The Institute of Food and Agricultural Sciences is an equal opportunity/affirmative action employer authorized to provide research, educational information and other services only to individuals and institutions that function without regard to race, color, sex, age, handicap, or national origin. For information on obtaining other extension publications, contact your county Cooperative Extension Service office. Florida Cooperative Extension Service/Institute of Food and Agricultural Sciences/University of Florida/Christine Taylor Waddill, Dean.

Figure 1. Adult *Hemisphaerota cyanea* (Say) (dorsal view). Credits: J.F. Butler, University of Florida

Figure 2. Adult *Hemisphaerota cyanea* (Say) (ventral view). Credits: J.F. Butler, University of Florida

Life Cycle

Eggs are yellow and elongated. They are cemented to host leaves and then covered with a double row of fecal pellets by the adult female. The fecal covering probably protects the eggs from predators and parasitoids. In Georgia, viable eggs were present from early March to mid-April.

Upon hatching, the young larva begins to excrete strands of dry feces that curl up over its back and accumulate to form a dense thatch-like shield. Larvae have an anal fork to hold the fecal shield in place. The fecal shield has been shown to be an effective defense against a predacious larva of *Cycloneda sanguinea* (L.) (Coleoptera: Coccinellidae) and the

predacious stink bug, *Stiretrus anchorago* (Fabricius) (Hemiptera: Pentatomidae). However, the carabid beetle, *Calleida viridipennis* (Say) (Coleoptera: Carabidae) is able to penetrate the larva's defense by chewing through the shield or by getting underneath it and prying it up.

The larvae cause trough-like feeding damage as they scarify the leaf epidermis, but rarely cause significant economic damage to trees.

Figure 3. Feces-covered eggs and young larva of *Hemisphaerota cyanea* (Say). Credits: J.F. Butler, University of Florida

Figure 4. Dorsal view of feces-covered mature larva of *Hemisphaerota cyanea* (Say). Credits: J.F. Butler, University of Florida

After reaching maturity, the larva clings to the leaf surface and pupates under the fecal thatch. Larvae and pupae are found throughout mid to late summer. Adults are found throughout the entire year.

Figure 5. Lateral view of feces-covered mature larva of *Hemisphaerota cyanea* (Say). Credits: J.F. Butler, University of Florida

Figure 7. Saw palmetto, *Serenoa repens* (Arecaceae), a host of the Florida tortoise beetle, *Hemisphaerota cyanea* (Say). Credits: D.W. Hall, University of Florida

Figure 6. Larva of *Hemisphaerota cyanea* (Say), without fecal shield showing anal fork. Credits: M.W. Sanderson, Illinois Natural History Survey

The Florida tortoise beetle is found most commonly on native saw palmetto, *Serenoa repens*, but also feeds on cabbage palm, *Sabal palmetto* (W. Bartram) Small, dwarf palmetto, *Sabal minor* (Jacq.) Pers., scrub palmetto, *Sable etonia* Swingle ex Nash, and probably other native palms. It also has been reported from a variety of exotic palms in Florida.

Selected References

Attygalle, A.B., D.J. Aneshansley, J. Meinwald, and T. Eisner. 2000. Defense by foot adhesion in a chrysomelid beetle (*Hemisphaerota cyanea*): Characterization of the adhesive oil. *Zoology (Jena)* 103:1-6.

Beshear, R.J. 1969. Observations on the life history of *Hemisphaerota cyanea* in Georgia. *Journal of the Entomological Society of Georgia* 4: 168-170.

Eisner, T. and D.J. Aneshansley. 2000. Defense by foot adhesion in a beetle (*Hemisphaerota cyanea*). *Proceedings of the National Academy of Science* 97:6568-6573.

Eisner, T. and M. Eisner. 2000. Defensive use of a fecal thatch by a beetle larva (*Hemisphaerota cyanea*) *Proceedings of the National Academy of Science* 97:2632-2636.

Jackman, J.A. 1976. A tortoise beetle, *Hemisphaerota cyanea*, on palms in Texas. *The Southwestern Entomologist* 1:181-183.

Woodruff, R.E. 1965. A tortoise beetle (*Hemisphaerota cyanea* (Say)) on palms in Florida (Coleoptera: Chrysomelidae). Florida Department of Agriculture, Division of Plant Industry. Entomology Circular No. 35. Gainesville, FL