

# Citrus Pest Quick Guide:

## Lebbeck Mealybug (*Nipaecoccus viridis* Newstead)

Lebbeck mealybug, *Nipaecoccus viridis*, is an exotic mealybug previously documented on dodder and tallow wood in Palm Beach County and intercepted at ports of entry. In June 2019, it was documented feeding on and damaging citrus in Highlands County, Florida. It has a wide range of hosts in Florida including citrus, mango, and several ornamental plants.

### Life Stages

Each adult female produces approximately 600 eggs during her lifetime. The offspring develop into mobile nymphs called “crawlers.” Lebbeck mealybug nymphs resemble, and are easily confused with, scale nymphs. They are reddish-purple, very small, and have legs and antennae that are visible under magnification.

The adult male is 1.3–2.5 mm long and brownish-purple. It has well-developed front wings.

The female has three nymphal stages and slowly moves between stages to new feeding sites. The adult is oval, reddish-purple, and 2.4–4 mm long and 1.5–3 mm wide. Once the female begins laying eggs, it develops a white to pale yellow wax-covered ovisac (egg sack) that houses tiny red eggs. In their native Middle East, the females can live up to 50 days. They die shortly after laying all of their eggs.

### Damage

Lebbeck mealybug has been mistaken for other wax-producing pests (mealybugs and scales) in several fields, but its damage and feeding sites are distinct from those of these other pests. *N. viridis* prefers to feed and reproduce on fast-growing tissues like new growth, new branches, and fruit. Fruit feeding causes hardened lumps and/or discoloration. Leaves also become twisted/distorted from feeding. Fruit drop is the biggest factor in crop loss.



***N. viridis* crawlers.**  
Credit: T. R. Weeks, UF/IFAS



**Adult female next to ovisac with another female.**  
Credit: T. R. Weeks, UF/IFAS



**Female without ovisac.**  
Credit: L. M. Diepenbrock, UF/IFAS



**Fruit damage from *N. viridis* feeding.**  
Credit: T. R. Weeks, UF/IFAS



***N. viridis* on calyx button.**  
Credit: T. R. Weeks, UF/IFAS



***N. viridis* on blossom end of fruit.**  
Credit: T. R. Weeks, UF/IFAS

1. This document is ENY-1005, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date April 2020. Visit the EDIS website at <http://edis.ifas.ufl.edu> for the currently supported version of this publication.

2. Lauren M. Diepenbrock, assistant professor, Entomology and Nematology Department, UF/IFAS Citrus Research and Education Center; and Jamie D. Burrow, Extension program manager, UF/IFAS Citrus REC; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.