



## Toxicity of Tomato and Bell Pepper Insecticides/Miticides to Beneficial Insects<sup>1</sup>

---

Mark A. Mossler<sup>2</sup>

The following table presents categories that were either obtained from the Environmental Protection Agency, Kopperts Side-Effects Database, UF/IFAS Pest Management Strategic Plans, or label interpretation. When different categories are listed for different life stages (e.g. highly toxic in larval stage but nontoxic as an adult), the more toxic category has been placed in the table.

- 
1. This document is PI-90, one of a series of publications of the Pesticide Information Office, Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. For additional information, contact the Pesticide Information Office, University of Florida, P. O. Box 110710, Gainesville, FL 32611-0710, (352) 392-4721. Published December 2005. Please visit the EDIS Web site at <http://edis.ifas.ufl.edu>.
  2. Mark A. Mossler, Doctor of Plant Medicine, Pesticide Information Office, Agronomy Department. Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 32611-0710.

**The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean**

**Toxicity of chemical pest management tools to beneficial invertebrates in Florida tomato and bell pepper.**

<b>Beneficial Insects/ Mites</b>	<b>Beneficial mites</b>	<b>Big-eyed bugs</b>	<b>Damsel bugs</b>	<b>Ground beetles</b>	<b>Honeybees</b>	<b>Lacewings</b>	<b>Ladybird beetles</b>	<b>Minute pirate</b>	<b>Parasitic wasps</b>	<b>Predatory midges</b>	<b>Predatory thrips</b>	<b>Spiders</b>	<b>Syrphid fly larvae</b>
<b>Pest Management Tools (IRAC MoA Class)</b>													
<b>Registered materials</b>													
Abamectin (6)	H	O	H	O	H	H	S	O	H	H	H	O	S
Acephate (1)*	H	H	H	H	H	H	H	H	H	H	H	H	H
Acetamiprid (4)	S	O	O	O	M	M	M	O	M	H	H	O	O
Azadirachtin (18)	S	O	O	O	S	S	S	O	M	O	O	O	O
Azinphos-methyl (1)**	M	H	H	H	H	H	H	H	H	H	H	H	H
Bifenazate (25)	S	O	O	O	M	S	S	O	S	O	O	O	O
Bifenthrin (3)	H	H	H	H	H	H	H	H	H	H	H	H	H
Boric acid (no class)	M	O	O	O	S	S	S	S	O	M	M	O	M
Buprofezin (16)**	O	O	O	O	O	S	O	O	S	M	O	O	O
Carbaryl (1)	S	H	H	H	H	H	H	H	H	H	H	H	H
Chlorpyrifos (1)*	H	H	H	H	H	H	H	H	H	H	H	H	H
Cryolite (9)	O	O	O	O	O	O	O	O	O	O	O	O	O
Cyfluthrin (3)	H	H	H	H	H	H	H	H	H	H	H	H	H

**Toxicity scale:**

O = nontoxic

S = slightly toxic

M = moderately toxic

H = highly toxic

\*Bell Pepper only

\*\*Tomato only

\*\*\*When applied on foliage

**Toxicity of chemical pest management tools to beneficial invertebrates in Florida tomato and bell pepper.**

<b>Beneficial Insects/ Mites</b>	<b>Beneficial mites</b>	<b>Big-eyed bugs</b>	<b>Damsel bugs</b>	<b>Ground beetles</b>	<b>Honeybees</b>	<b>Lacewings</b>	<b>Ladybird beetles</b>	<b>Minute pirate</b>	<b>Parasitic wasps</b>	<b>Predatory midges</b>	<b>Predatory thrips</b>	<b>Spiders</b>	<b>Syrphid fly larvae</b>
<b>Pest Management Tools (IRAC MoA Class)</b>													
Cyhalothrin (gamma/lambda) (3)	H	H	H	H	H	H	H	H	H	H	H	H	H
Cyromazine (17)	O	M	M	O	O	O	O	M	O	O	O	O	O
Diazinon (1)**	H	H	H	H	H	H	H	H	H	H	H	H	H
Dicofol (20)	H	S	S	S	S	S	S	S	M	H	S	S	H
Diflubenzuron (15)*	O	O	M	O	M	H	M	H	O	O	O	O	O
Dimethoate (1)	H	H	H	H	H	H	H	H	H	M	H	H	H
Dinotefuran (4)	M	M	M	M	H	M	H	H	H	H	H	M	H
Disulfoton (1)*	O	O	O	O	O	O	O	O	O	O	O	O	O
Emamectin (6)	H	O	H	O	H	H	S	O	H	H	H	O	S
Endosulfan (2)	H	H	H	H	H	H	H	H	M	H	H	H	H
Esfenvalerate (3)	H	H	H	H	H	H	H	H	H	H	H	H	H
Fenpropathrin (3)**	H	H	H	H	H	H	H	H	H	H	H	H	H
Imidacloprid (4)	M	M	M	M	H	M	H	H	H	H	H	M	H
Indoxacarb (22)	S	S	S	S	H	S	S	S	H	S	S	O	H

**Toxicity scale:**

O = nontoxic

S = slightly toxic

M = moderately toxic

H = highly toxic

\*Bell Pepper only

\*\*Tomato only

\*\*\*When applied on foliage

**Toxicity of chemical pest management tools to beneficial invertebrates in Florida tomato and bell pepper.**

<b>Beneficial Insects/ Mites</b>	<b>Beneficial mites</b>	<b>Big-eyed bugs</b>	<b>Damsel bugs</b>	<b>Ground beetles</b>	<b>Honeybees</b>	<b>Lacewings</b>	<b>Ladybird beetles</b>	<b>Minute pirate</b>	<b>Parasitic wasps</b>	<b>Predatory midges</b>	<b>Predatory thrips</b>	<b>Spiders</b>	<b>Syrphid fly larvae</b>
<b>Pest Management Tools (IRAC MoA Class)</b>													
Malathion (1)	M	H	H	H	H	H	H	H	H	H	H	H	H
Methamidophos (1)**	H	H	H	H	H	H	H	H	H	H	H	H	H
Methomyl (1)	H	H	H	H	H	H	H	H	H	H	H	H	H
Methoxyfenozide (18)	O	O	S	O	O	O	O	S	O	O	O	O	O
Naled (1)*	H	H	H	H	H	H	H	H	H	H	H	H	H
Oils (no class)	M	O	O	O	O	S	S	O	S	S	S	O	S
Oxamyl (1)***	H	H	H	H	H	H	H	H	H	H	H	H	H
Oxydemeton-methyl (1)*	H	H	H	H	H	H	H	H	H	H	H	H	H
Permethrin (3)	H	H	H	H	H	H	H	H	H	H	H	H	H
Pymetrozine (9)	O	O	O	O	O	O	O	O	O	O	O	O	O
Pyrethrins + Rotenone (3)	M	M	M	M	H	M	M	M	M	M	M	M	M
Pyrethrins + PBO (3)	M	M	M	M	H	M	M	M	M	M	M	M	M
Pyriproxyfen (7)	S	S	S	S	O	S	H	S	S	S	S	S	M
Soaps (no class)	H	M	M	S	O	H	M	S	M	M	M	O	M

**Toxicity scale:**

O = nontoxic

S = slightly toxic

M = moderately toxic

H = highly toxic

\*Bell Pepper only

\*\*Tomato only

\*\*\*When applied on foliage

**Toxicity of chemical pest management tools to beneficial invertebrates in Florida tomato and bell pepper.**

Beneficial Insects/ Mites  Pest Management Tools (IRAC MoA Class)	Beneficial mites	Big-eyed bugs	Damsel bugs	Ground beetles	Honeybees	Lacewings	Ladybird beetles	Minute pirate	Parasitic wasps	Predatory midges	Predatory thrips	Spiders	Syrphid fly larvae
Spinosad (5)	M	S	S	S	H	M	M	S	M	S	S	O	S
Spiromesifen (23)	M	O	O	O	O	S	S	S	S	S	S	O	S
Sulfur (8)	M	S	S	S	S	S	S	S	S	S	S	O	S
Tebufenozide (18)	O	O	S	O	O	O	O	S	O	O	O	O	O
Thiamethoxam (4)	S	M	H	M	H	S	H	H	H	H	H	M	H
zeta-Cypermethrin (3)	H	H	H	H	H	H	H	H	H	H	H	H	H

**Toxicity scale:**

- O = nontoxic
- S = slightly toxic
- M = moderately toxic
- H = highly toxic

- \*Bell Pepper only
- \*\*Tomato only
- \*\*\*When applied on foliage