

Fireweed (Heartleaf nettle) Control in Pastures¹

J. A. Ferrell²

Introduction

Fireweed (*Urtica chamaedryoides*) (Figure 1) is native to Florida, but has only recently become problematic. This winter annual species is commonly observed in north and central Florida pastures, particularly in bareground areas (near feeding pens and under fences) as well as along tree lines where forage grasses are less dense.



Figure 1. Fireweed (*Urtica chamaedryoides*).

Fireweed is particularly troublesome because it possesses stinging hairs that easily embed in skin. Once exposed to the toxin, severe irritation can occur for several hours. Though generally avoided by cattle, horses are more likely to browse fireweed and develop stress symptoms. These symptoms commonly manifest themselves as weight loss, or difficulty in swallowing and breathing for many days after consumption. In extreme cases, young horses have died after rolling in fireweed and becoming over-exposed to the toxins in the leaf hairs.

Biology

Fireweed leaves resemble that of a strawberry plant, (Figure 2) but the plant as a whole has little resemblance to strawberry. The plant has square stems and small pale green flower clusters. Small stinging hairs are found on the stems, petioles and leaves. These hairs contain irritants which have been shown to cause respiratory stress and local allergic reactions when ingested or inhaled.

1. This document is SS-AGR-41, one of a series of the Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date May 2006. Reviewed: November 2006. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.
2. J. A. Ferrell, assistant professor, Agronomy Department; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition. Use herbicides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean



Figure 2. Fireweed leaves.

Control

Little work has been reported for control of this seasonal species. Therefore, experiments were conducted in March 2006 to determine which pasture herbicides are most effective on fireweed.

It was observed that 2,4-D and Telar were ineffective on fireweed (Table 1). Glyphosate, which can be applied when pasture grasses are dormant, provided as much as 80% control. However, in other experiments we have found glyphosate to be inconsistent on fireweed, sometimes providing as little as 30% control. Since glyphosate can severely injure pasture grasses that are not fully dormant, and because of its inconsistency on fireweed, we do not recommend the use of this product.

Weedmaster at 1.5 qt/A did not provide acceptable levels of control, but Milestone, Remedy, and Pasturegard were found to be highly effective. Within 2 weeks of application, over 90% of the fireweed plants were dead and the remaining individuals were yellow and dying. By 6 weeks after treatment, no fireweed could be found.

It is our recommendation that Milestone, Remedy, or Pasturegard herbicides be used for effective control of fireweed. These herbicides can be applied any time of year to warm-season forage grasses. There are no grazing restrictions for beef cattle with these herbicides, but lactating dairy animals must be removed for 0 and 14 days with Milestone and Remedy, respectively, and one season for Pasturegard.

Mowing provides no benefit to control of this species. In fact, mowing has been found to result in smaller plants, but with many more stinging hairs. Additionally, the seed is surrounded by a sticky substance that can be transported by mower blades to areas not infested with this weed.

If not controlled, fireweed generally disappears in May with the onset of summer temperatures.

Table 1. Control of fireweed with various herbicides.

Herbicide	Rate Product/A	Herbicide cost ¹ \$ per acre	% Control	
			2 wat ²	6 wat
2,4-D amine	2 qt	6	10	0
Weedmaster	1.5 qt	11	33	65
Remedy	1 qt	25	93	100
Pasturegard	1.5 qt	20	92	100
Milestone	5 fl. oz	14	95	100
Glyphosate	1 qt	5	70	80
Telar	0.5 oz	10	30	10

¹These are approximate values taken from "Approximate herbicide pricing - 2006" (<http://edis.ifas.ufl/WG056>) and do not include application costs.

²wat=weeks after treatment.