



## Weed Control in Okra<sup>1</sup>

---

William M. Stall<sup>2</sup>

Okra is grown throughout the state during the warm season of the year. During this time, warm-season annual weeds can be a serious problem in the field.

Herbicides labeled for use in okra are severely limited (see Table 1). Several have not been relabeled during the reregistration process.

Roundup may be used to reduce the weed population in the field, either as a broadcast spray or as a spot treatment when applied to emerged and growing weeds before okra planting.

Trifluralin (Treflan, etc) is the only preplant incorporated (preemergent to weeds) treatment now labeled in okra.

Cultivation is commonly used in okra, either with sweeps or rolling cultivators. Care should be taken not to throw soil too far up on the okra stems. This practice can increase stem rot and stem decay.

In some areas of the state, okra has been planted as a second crop, behind mulched vegetables, after the mulch has been removed.

This production practice has been quite successful, but okra must not be planted as a second crop into soil containing residual herbicides from the first crop that are phytotoxic to okra. Read the label of each herbicide to find out the length of time that must elapse before non labeled crops may be planted in the field.

---

1. This document is HS192, one of a series of the Horticultural Sciences Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Last revision date: October, 2006. Please visit the EDIS Website at <http://edis.ifas.ufl.edu>.

2. William M. Stall, professor, Horticultural Sciences Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 32611.

**The use of trade names in this publication is solely for the purpose of providing specific information. It is not a guarantee or warranty of the products named, and does not signify that they are approved to the exclusion of others of suitable composition.**

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean

**Table 1.** Chemical weed controls: okra.

Herbicide	Labeled crops	Time of application to crop	Rates (lbs. AI./Acre)	
			Mineral	Muck
Carfentrazone (Aim)	Okra	Preplant Directed-hooded Row-middles	0.031	0.031
<b>Remarks:</b> Aim may be applied as a preplant burndown treatment and/or as a post-directed hooded burn-down application to row middles for the burndown of emerged broadleaf weeds. May be tank mixed with other registered herbicides. May be applied at up to 2 oz (0.031 lb ai). Use a quality spray adjuvant such as crop oil concentrate (coc) or non-ionic surfactant at recommended rates.				
Glyphosate (Roundup, Durango, Touchdown, Glyphomax)	Okra	Chemical fallow Preplant, pre emergence, Pretransplant	0.3 - 1.0	----
<b>Remarks:</b> Roundup, Glyphomax and touchdown have several formulations. Check the label of each for specific labeling directions.				
Pelargonic Acid (Scythe)	Okra	Preplant Preemergence Directed-Shielded	3-10% v/v	3-10% v/v
<b>Remarks:</b> Product is a contact, non-selective, foliar applied herbicide. There is no residual activity. May be tank mixed with glyphosate. May be tank mixed with trifluralin for soil residual activity. Consult label for rates and other information.				
Trifluralin (Several Brands)	Okra	Preplant incorporated	0.5 - 0.75	----
<b>Remarks:</b> Controls germinating annuals, especially grasses. Incorporate 4/ or less within 8 hours. Results in Florida are erratic on soils with low organic matter and clay contents. Note precautions against planting non-registered crops within 5 months.				