



## Limpograss: A Quick Reference<sup>1</sup>

Yoana C. Newman<sup>2</sup>

In Florida, limpograss (*Hemarthria altissima*) is one of the warm-season perennial grasses that has been increasingly adopted by ranchers during the past 15 years because of its superior digestibility, adaptation to seasonally flooded soils, and tolerance of light frost. It is grown throughout the state; however, it is extensively used in South and Central Florida. Limpograss may yield 8 to 10 tons of hay per acre under good fertility and soil moisture.

Recommended cultivars are Floralta (latest release), and Bigalta which adapts better to sands higher in organic matter. Older cultivars, Redalta, and Greenalta, are not recommended because of lower forage quality.

### Origin

Native to South Africa.

### Use

Grazing, haying, and stockpiling (standing hay crop). It has superior digestibility at longer regrowth periods compared to other warm-season grasses (may be as high as 70%), but, in mature stockpiles, this can drop to as low as 40%. Protein is adequate in spring and fall, but in summer it may drop below 7%. Stockpiled material (12 to 16 weeks old may only have 3 to 4%

protein and cattle needs to be supplemented with protein or non-protein nitrogen source).

### Description

Tall prostrate or decumbent growth (sod type) that can reach over 3 feet tall.

### Adaptation

pH: Adapts to acid soils. Nevertheless, if soil is lower in pH it should be limed to the target pH, 5.5.

Soil: Fertile or fertilized sand to clay. Better adapted to flatwoods than well drained sands. Adapts to seasonally flooded sites.

Rainfall: High moisture and rain.

Climate: Subtropical to tropical wet. Tolerates light frost and begins growth soon after occurrence of freezing temperatures. No winter kill with minimum temperatures of 14 degrees F in Gainesville.

However, shows some winter kill when temperatures are as low as 8 degrees F.

1. This document is SS AGR 302, one of a series of the Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date March 2008. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.

2. Yoana C. Newman, assistant professor, forage extension specialist, Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

All chemicals should be used in accordance with directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean

Management Practices

Planting date: April-May (if irrigated); best during rainy season (June-August).

Planting rate: 1000 to 1500 lb/acre of stem tops or stolons (requires vegetative establishment as it produces very few viable seed).

Planting depth: 2-3 inches.

Stubble height: No less than 12 inches when rotationally grazing (stocking). When continuously stocking, no less than 16 inches of stubble height should be left to ensure adequate stand persistence regardless of where in the state it is planted. Plant material that reaches 20 to 24 inches tends to lay flat where cattle will likely trample on it, if not used, it will build up an undesired thatch.

Fertilization

Planting: Whenever root growth has initiated (one to two weeks after planting), apply 30 lb N/acre, all phosphorus ( $P_2O_5$ ) and 50% of potassium ( $K_2O$ ) recommended in soil test. Thirty to 50 days later apply rest of the potassium, plus 70 lb N/acre.

Grazing: For grazed, established stands, apply 60 lb N/A and all of the  $P_2O_5$  and  $K_2O$  in late winter or early spring. Apply an additional 60 lb N in late summer or early fall. For a minimum fertilization alternative, ignore the P and K recommendation and apply only 60 lb N per year.

Hay: 80 lb N/acre/cut + all phosphorus ( $P_2O_5$ ), and all potassium ( $K_2O$ ) in late winter or early spring. After each cutting, apply 80 lb N plus 40 lb of  $K_2O$ /acre. Apply fertilizer up to 6 weeks prior to end of season. Include 20 lb of  $P_2O_5$ /acre in the supplemental fertilizer if the soil tested low or medium in P.

Stockpiling: Remove cattle in mid August from area to be stockpiled; follow by fertilization (80 lb N plus 40 lb of  $K_2O$ /acre).

Broad-leaf Weed Control

(Check with County Agent or Extension Weed Specialist for updates and restrictions, and always follow label directions).

Do not use 2,4-D (several brands) or 2,4-D containing herbicides such as Weedmaster (2,4-D + dicamba) because limpogross may be injured or killed.

Use dicamba herbicide (brand examples are Banvel, Clarity or Vanquish). Limpogross has shown more tolerance to dicamba than 2,4-D.

Pests and control

(Check with County agent or extension weed specialist for updates on rates and restrictions, and always follow label directions).

- Spittlebugs are likely to be present if a thatch has built up in the pastures. Graze or cut accumulated summer growth to avoid this situation.
- Chinchbugs are likely to be present in very dry years.
- Isolated cases of Armyworm damage have been reported.

**Additional Information**

- Newman, Y.C. and L.E. Sollenberger. 2005. Grazing management and nitrogen fertilization effects on vaseygrass persistence in limpogross pastures. *Crop Science* 45:2038-2043. <http://crop.scijournals.org/cgi/reprint/45/5/2038?ck=nck>
- Newman, Yoana C., Lynn E. Sollenberger, Carrol C. Chambliss, William E. Kunkle. 2003. Yearling Beef Heifer Performance on Limpogross Pastures Grazed. 2003 Florida Beef Report. *Animal Sci. Dep. University of Florida*. [http://www.animal.ufl.edu/extension/beef/beef\\_cattle\\_report/2003/Forages-Chap5.pdf](http://www.animal.ufl.edu/extension/beef/beef_cattle_report/2003/Forages-Chap5.pdf)
- Newman, Y.C., L.E. Sollenberger, and C.G. Chambliss. 2003. Canopy characteristics of continuously stocked limpogross swards grazed to different heights. *Agron. J.* 95:1246-1252. <http://agron.scijournals.org/cgi/reprint/95/5/1246>

- Newman, Y.C., L.E. Sollenberger, A.M. Fox, and C.G Chambliss. 2003. Canopy height effects on vaseygrass and bermudagrass spread in limpograss pastures. *Agron. J.* 95:390-394.  
<http://agron.scijournals.org/cgi/reprint/95/2/390>
- Newman, Y.C., L.E. Sollenberger, W.E. Kunkle, and C.G. Chambliss. 2002. Crude protein fractionation and degradation parameters of limpograss herbage. *Agron. J.* 94 (6):1381-1386.  
<http://agron.scijournals.org/cgi/reprint/94/6/1381>
- Newman, Y.C., L.E. Sollenberger, W.E. Kunkle, and D.B. Bates. 2002. Canopy height and nitrogen supplementation effects on performance of heifers grazing limpograss. *Agron. J.* 94 (6):1375-1380.  
<http://agron.scijournals.org/cgi/reprint/94/6/1375>