**ENH1100** 



# New Florida Foliage Plant Cultivar: 'Orange Hot' *Anthurium*<sup>1</sup>

R.J. Henny, J. Chen, and T.A. Mellich <sup>2</sup>



**Figure 1.** A mature *Anthurium* 'Orange Hot' that was grown in a 1.6 L (6-inch) container.

## Description

Mature leaves of a one year old 'Orange Hot' plant are lance-shaped, 18 to 20 cm (7.2 to 8 in.) in length and 9 to 11 cm (3.6 to 4.4in.) wide at the widest part. The upper surface leaf color is dark green (RHS 146B-C; Royal Horticultural Society, 1995). The underside of the leaf is a lighter green (RHS 139A). Petioles are uniform yellow-green (RHS 146B-C). *Anthurium* 'Orange Hot' produces spathes

that are 6 to 7 cm (2.4 to 2.8 in.) long and 4 to 5 cm (1.6 to 2.0 in.) wide. They are held on peduncles 25 to 35 cm (10 to 14 in.) tall that locate the inflorescences higher than the leaf canopy where they are easily seen and enjoyed. When they unfurl, new spathes are a light salmon orange color (RHS 39B-C), and they gradually fade over a period of two to three weeks to a lighter orange (RHS 36B-C). The spadix at anthesis is a light orange color (RHS 38A) at the base and increases in intensity to orange (RHS 39A-B) at the apex. In older flowers, the spadix becomes a uniform lighter orange color (RHS 37D). Peduncles are yellow green at the base (RHS 152A-B) and darken to grayed-orange (RHS 172D) near the spathe.

#### **Performance**

Performance characteristics of *Anthurium X* 'Orange Hot' were determined in a growth study. Thirty liners were selected from well-rooted cuttings in 25-mm (72 insert) cell trays. Liners were potted into 1.6L (6-inch) plastic pots containing a growing medium of 3:2 Canadian peat: perlite by volume. The soil was amended with 0.9 kg m<sup>-3</sup> of Micromax, (Sierra Chemical Company, Milpitas, CA) and 4.1

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean

<sup>1.</sup> This document is ENH1100, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date May 2008. Visit the EDIS Web Site at http://edis.ifas.ufl.edu.

R.J.Henny, Professor, Department of Environmental Horticulture; J. Chen, Associate Professor, Department of Environmental Horticulture; and Terri Mellich, Biologist, University of Florida, Institute of Food and Agricultural Sciences, Mid-Florida Research and Education Center, 2725 Binion Road, Apopka, FL 32703

kg m<sup>-3</sup> dolomite. Plants were grown in a shaded greenhouse with a maximum irradiance of 125µmol s-1m<sup>-2</sup> under a natural photoperiod and a temperature range of 15 to 34°C (63 to 97°F). Ten plants were grown at each of three fertilizer levels. The fertilizer was applied as a liquid at 100ml/pot per week. Fertilizer levels were derived from stock solutions made in a 3N-1P-2K ratio. The test was set in a completely randomized design. At the end of the test, data recorded canopy height and width, length and width of largest leaf and number of open flowers. Plants were also rated for visual quality on a scale of 1 to 5, where 1 = dead, 3 = acceptable, saleable and 5 = excellent quality.

Anthurium X 'Orange Hot' reached marketable size in 7 months. Fertilizer level did not significantly affect any of the variables measured except canopy width, which increased linearly as fertilizer levels increased (Table 1). All plants were rated between good and excellent at each fertilizer level. Overall plant quality averaged 4.2 out of a possible 5.0. Mature plants were wider than they were tall. Both primary and secondary shoots bloomed resulting in average flower counts of four to five per plant. Plants were well branched. Canopy height and width averaged 20.4 cm by 42.9 cm (8.2 by 17 inches) for the 1.6L (6-inch) pot size. For additional details the reader is referred to the 2003 'Orange Hot' Anthurium article in HortScience (Henny et al. 2003).



**Figure 2.** Anthurium X 'Orange Hot' growing in a Florida foliage plant nursery.

## **Availability**

Anthurium X 'Orange Hot' is intended for commercial producers growing finished plants in 1.6 or 3.9L (6-inch or 8-inch) containers. Stock plants have been released to University of Florida licensed cooperating Florida tissue culture labs for

propagation and distribution. Inquiries regarding participating labs may be obtained by writing the Florida Foundation Seed Producers, Inc., P.O. Box 309, Greenwood, FL 32443. Plants for research purposes may be obtained directly from the author.

### References

R.J Henny, D.J. Norman, and J. Chen. 2003. 'Orange Hot' *Anthurium*. HortScience, Vol. 38(1):133-134.

R.J. Henny, and D.J. Norman. 1999. 'Red Hot' *Anthurium*. HortScience, Vol. 34(1):153-154.

Kamemoto, H. and A.R.Kuehnle. 1996. Breeding Anthuriums in Hawaii. Univ. of Hawaii Press. 132p.

Royal Horticultural Society. 1995. The Royal Horticultural Society Colour Chart. 3<sup>rd</sup> ed. Royal Horticulture Society, London.

**Table 1.** Final canopy height and width, length and width of largest leaf, visual quality and total number of open flowers produced by *Anthurium* X 'Orange Hot' grown in 1.6L pots for 7 months.

Fertilizer(N)	Canopy		Leaf		Open	Visual
g/m <sup>-2</sup> /yr	height (cm)	width (cm)	length (cm)	width (cm)	Flowers	Quality <sup>z</sup>
34	19.8	39.2	18.4	10.0	4.4	3.8
68	21.3	44.5	18.6	10.1	5.6	4.5
101	20.2	45.0	19.5	10.2	4.5	4.4
Significance <sup>y</sup>	NS	L	NS	NS	NS	NS

ZVisual rating where 1= dead, 3= saleable, and 5= excellent

<sup>&</sup>lt;sup>y</sup>NS and L\* = Nonsignificant or Linear significance at 5% level respectively.