

Tropical Foliage Plant Development: Breeding Techniques for Anthurium and Spathiphyllum¹

R.J. Henny, J. Chen and T.A. Mellich²

Anthurium and *Spathiphyllum* are popular tropical ornamental plants. They have lush leaves and exotic flowers and are prized for interior decoration in both residential and public settings.

Anthurium and *Spathiphyllum* are among the many plants in the Araceae family that are commercially cultivated as tropical ornamentals. Members of this plant family, including *Anthurium* and *Spathiphyllum*, are commonly referred to as aroids. The Araceae family includes at least 10 important ornamental tropical foliage plant genera.

Anthurium, also called flamingo lily, exhibit large, showy-red, orange, pink or white flowers arising from plants with glossy, arrow-shaped evergreen leaves. *Anthuriums* are often cropped for cut flowers, and large-sized *Anthurium* plants that yield long stems are favored for the floral market. New interspecific hybrids -- such as *Anthurium* X 'Lady Jane' and *Anthurium* X 'Southern Blush' -- are popular for the tropical flowering-foliage market because of their overall smaller size. *Anthurium* X 'Red Hot' (Henny, 1999; Henny et al., 2008) is one such interspecific hybrid that exhibits numerous showy-red spathes and a compact, freely branching growth habit (Figure 1).




Figure 1. *Anthurium* 'Red Hot' plants growing in a commercial Florida foliage plant nursery.

Spathiphyllum has become one of the most popular foliage-plant genus in the foliage industry. The wholesale value of finished *Spathiphyllum* totaled \$32 million in 1998 (USDA, 1999). These easy-care, attractive plants, commonly called peace lily, originate from the tropical rainforests of Central America. The plants have dark, evergreen foliage.

Spathiphyllum cultivars are popular interiorscape plants in part because sizes available on the market range from very small varieties, such as Petite, to the larger forms of 'Mauna Loa' and 'Sensation'.

From an ornamental viewpoint, the spathes and spadices (technically both are part of an inflorescence) are called flowers and consist of a

1. This document is ENH1102, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date June 2008. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.
2. R.J.Henny, professor, J. Chen, associate professor, and T.A. Mellich, biological scientist, Environmental Horticulture Department, University of Florida Institute of Food and Agricultural Sciences Mid-Florida Research and Education Center-Apopka, FL

white, sail-shaped spathe surrounding a spadix (Chen et al., 2003). Market demand for new varieties of both *Anthurium* and *Spathiphyllum* has stimulated interest in breeding new cultivars.

However, hybridization within the aroid family can be difficult to achieve because of sporadic flower production and specific requirements for successful pollination.

Research by the Foliage Breeding Program at the University of Florida Mid-Florida Research and Education Center (MREC)-Apopka has given plant breeders new techniques to manipulate flowering and pollination in ornamental aroids for hybridization purposes. These techniques improve breeding potential for *Anthurium*, *Spathiphyllum*, and other aroid genera that were previously difficult to hybridize.

Factors Affecting Flowering

For breeding purposes, flower production is vital. Pollinations can take place only when parents with desired traits are blooming simultaneously. Therefore, careful planning is required to ensure a sufficient supply of flowers. Differences in the natural flowering cycles, even within the same genus, can be frustrating for breeders of tropical plants. Flowering times may be affected by environmental conditions. For example, in a greenhouse heated to a minimum night temperature of 60°F, plants of the same species will flower approximately one month earlier than those in a comparable greenhouse heated to 50°F. Cultural conditions -- such as light, irrigation and nutrition -- can affect flower initiation. Plant size and maturity can also be factors.

In the case of *Spathiphyllum*, natural flowering occurs once annually -- in late winter or, in Central Florida, in very early spring. However, different *Spathiphyllum* species and cultivars may not flower simultaneously or at all. Flower production in *Anthurium* plants is not a problem since, once mature, plants produce flowers continuously throughout the year.

Flower Induction

In *Spathiphyllum* the potential barrier to breeding caused by flower unavailability can be overcome by the use of gibberellic-acid (GA_3) sprays to stimulate flowering. At the MREC in Apopka, research in the aroid-breeding program showed that treatment of *Spathiphyllum* with gibberellic-acid (GA_3) sprays induces plants to flower throughout the year (Henny, 1981).

Treatment generally consists of a single foliar spray of 250 ppm GA_3 from a commercially available source (GibGro, 4% gibberellic acid, Agtrol Chemical Products, Houston, Texas 77074). Following treatment, plants flower within three-to-five months depending on the time of year. Flowers will appear more quickly if greenhouse temperatures are warm and the crop is rapidly growing. Within a genus, different species and cultivars generally flower close together

following GA_3 -treatment, so potential parental selection can be controlled. An additional benefit of using GA_3 to induce flowering is that treated plants produce more flowers than plants flowering under natural conditions. Some inflorescences produced following GA_3 -treatment may be deformed, but no detrimental effects on fertility -- as measured by pollen production or seed set -- have been observed.

Flower Structure

Anthurium and *Spathiphyllum* produce bisexual flowers (figures 2 and 3). Their inflorescences consist of a spadix enclosed by a spathe. The spadix is a fleshy spike covered with many small, complete flowers. (The term "complete" indicates that male and female structures are contained within each single floret.)

Pollen Collection and Transfer

In *Anthurium* and *Spathiphyllum* (aroid genera with bisexual flowers), the spathe unfurls and exposes the spadix several days before flowers are receptive for breeding.

The unfurling of the spathe reveals numerous, uniformly sized bisexual flowers located along the



Figure 2. An *Anthurium* spadix with receptive stigmas that are ready to be pollinated.

entire spadix. In *Anthurium*, new flowers become receptive each day, beginning at the spadix base and advancing gradually towards the tip over a two-week period (Figure 2). Pistillate (female) receptivity is indicated by a glistening shine of stigmatic surfaces and stickiness to the touch. Small drops of exudate are sometimes present. Flowers may remain receptive for more than one day. In contrast, all flowers on a *Spathiphyllum* spadix mature simultaneously (Figure 3).




Figure 3. *Spathiphyllum* spadix that is ready to be pollinated.

The stigmatic surfaces of both *Anthurium* and *Spathiphyllum* become dry and brown before pollen is dehiscid, so floral emasculation is not required. *Anthurium* pollen begins to appear along the spadix, usually at the bottom first and then proceeds towards the tip (Figure 4). *Anthurium* pollen is produced for up to two weeks because of the differences in the maturation of individual pistillate flowers. *Anthurium* pollen is not dispersed by wind. To transfer the pollen to the receptive female flowers, collect it on your fingertips then gently rub it onto receptive stigmas (Figure 2.)



Figure 4. Anthurium flowers that are producing pollen. They can no longer be used as a female parent.

A *Spathiphyllum* inflorescence will produce pollen over a period of three-to-four days (Figure 5). The pollen of *Spathiphyllum* is lighter and may be dispersed by air currents, so a brush should be used. Before attempting pollination, gently tap the *Spathiphyllum* inflorescence over a container, and collect the pollen as it falls. Subsequent transfers are made using a soft brush.




Figure 5. *Spathiphyllum* flowers that are producing pollen. Stigmas are no longer receptive and cannot be successfully pollinated.

Pollen Storage

It is always best to use fresh pollen. However, if pollen is in short supply, it can be stored in a container in a high-humidity environment in a refrigerator. Collect the pollen in a container, such as a petri dish. Place a lid over the petri dish, and set it on top of a wet paper towel or other moistened material. Enclose the wrapped petri dish within a sealed storage container. At no time should the pollen be directly in contact with the wet paper towel. Avoid splashing water droplets onto the pollen.

Spathiphyllum pollen may be stored for several days or weeks if necessary (Henny 2000).

Seed Production

To achieve maximum seed production, an *Anthurium* spadix needs to be pollinated more than once. However, it is possible to fertilize an entire *Spathiphyllum* spadix with a single pollination. To ensure seed set for *Anthurium* or *Spathiphyllum*, no special environmental manipulation -- such as controlling relative humidity -- is needed after pollen transfer. Pollinated flowers develop mature fruits within four-to-six months. *Anthurium* fruit will require up to five-to-eight months to mature. As seeds mature, *Anthurium* and *Spathiphyllum* spadices begin to turn to a yellowish color and soften.

Breeding for ornamental tropical foliage plants can be a lengthy process due to the limited numbers of seed obtainable and prolonged germination time. Several years may be needed to grow the hybrids to reach sufficient size for evaluation. Once a hybrid is selected and tested, tissue-culture methods may be applied so that a new aroid hybrid cultivar can be increased rapidly enough to reach commercial production levels within two-to-three years. The production of new hybrids leads to new plant introductions, and these foundations keep the foliage market fresh and profitable.

Literature Cited

Chen, J., D.B. McConnell, R.J. Henny, and K.C. Everitt. 2003. Cultural guidelines for commercial production of interiorscape *Spathiphyllum*. <http://edis.ifas.ufl.edu/EP161>.

Henny, R. J. 'Red Hot' Anthurium. 1999. HortScience. 34(1): 153-154.

Henny, R.J., J. Chen and T.A. Mellich. 2008. New Florida Foliage Plant Cultivar: Red Hot Anthurium. <http://edis.ifas.ufl.edu/EP363>

Henny, R.J. 1981. Promotion of flowering in *Spathiphyllum* Mauna Loa with Gibberellic Acid. HortScience 16 (4): 554-555.

Henny, R.J. 2000. Breeding ornamental aroids. p. 121-132. In: D.J. Callaway and M.B. Callaway (eds.), Breeding ornamental plants. Timber Press, Portland, OR.

USDA (United States Department of Agriculture). 1999. Floriculture crops 1998 Summary. Washington, DC.