***Hypogeococcus pungens* Granara de Willink (Insecta: Hemiptera: Pseudococcidae), a Mealybug¹**

Amanda Hodges and Greg Hodges²

Introduction

Hypogeococcus pungens is native to South America and was described by Granara de Willink (1981). A previous description by Williams (1973) classified *H. pungens* incorrectly as *Hypogeococcus festerianus* (Lizer y Trelles), and this confusion resulted in numerous publications listing *H. pungens* as *H. festerianus*. Little biological information is available for either species, but much of the available literature for *H. festerianus* actually is *H. pungens* information.

Field specimens of *H. pungens* are sometimes confused with the pink hibiscus mealybug, *Maconellicoccus hirsutus* (Green), but the potential hosts of *H. pungens* are more restrictive. The only other species of *Hypogeococcus* that occurs in Florida is the native *Hypogeococcus margaretae* Miller. The only known host for *Hypogeococcus margaretae* is the hatpin plant, *Eriocaulon decangulare*, and it has only been reported from St. Lucie and Indian River Counties in Florida.

Figure 1. Infestation of the mealybug, *Hypogeococcus pungens* Granara de Willink. Credits: Lyle J. Buss, University of Florida

Synonymy

A related species, *H. festerianus* is a valid species, but *H. pungens* has been misidentified as *H. festerianus* in the following publications: Hamon (1984), McFadyen and Tomley (1981), Suss and Trematerra (1986), and Williams (1973).

1. This document is EENY-459, one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: June 2009. This document is also available on Featured Creatures Web site at <http://creatures.ifas.ufl.edu>. Please visit the EDIS Web site at <http://edis.ifas.ufl.edu>. Additional information on these organisms, including many color photographs, is available at the Entomology and Nematology Department website at <http://entnemdept.ifas.ufl.edu/>
2. Amanda Hodges, SPDN assistant director in entomology and training/education, Department of Entomology and Nematology, Gainesville, FL, Greg Hodges, Florida Department of Agriculture and Consumer Services, Division of Plant Industry.

Details of these misidentifications are available on ScaleNet with a *Hypogeococcus pungens* valid name query.

Distribution

This mealybug has been reported from Australia, South America, the Caribbean, Europe, and North America. The Australian populations are the result of introduction for the biological control of cacti. Reports of *H. pungens* in the U.S. include Florida (1984) and Hawaii (2005).

In Florida, as of 2009, *H. pungens* occurs in the following counties: Brevard, Broward, Charlotte, Clay, Collier, Dade, Escambia, Hendry, Highlands, Hillsborough, Lake, Lee, Leon, Manatee, Miami-Dade, Monroe, Okaloosa, Orange, Palm Beach, Pasco, Pinellas, Polk, Sarasota, Seminole, St. Lucie and Volusia.

Identification Characteristics

Adult females are approximately 3 mm long (0.12 inches), produce a reddish body fluid when pierced, and lack both lateral filaments and an ovisac. *Hypogeococcus pungens* produce large amounts of wax and often form feeding clusters at nodal regions of host plants. The body shape is oval to round, and more round than other pinkish-colored mealybugs (Hodges et. al. 2008).

In Florida, heavy infestations of pink-colored mealybugs on *Portulaca* and *Alternanthera* species are frequently identified as *H. pungens*. This species is rarely found on cacti in Florida.

Immature and adult specimens may not be visible until the cottony wax is removed. Clusters of *H. pungens* tend to occur at nodal regions of leaves and stems.

Prior to confirming a diagnosis, scale and mealybug insects (Hemiptera: Coccoidea) generally must be slide-mounted. One of the unique morphological characteristics about *H. pungens* is the presence of three circuli (Miller et. al. 2007, Williams and Willink 1992). The circulus, a structure on the ventral surface of mealybugs, is thought to serve an adhesive function for the insect. *Hypogeococcus*

Figure 2. Shown infesting cacti is *Hypogeococcus spinosus* Ferris, which is not established in Florida. This species is related to *Hypogeococcus pungens* Granara de Willink. Credits: Lyle J. Buss, University of Florida

Figure 3. Late and early instar nymphs of the mealybug *Hypogeococcus pungens* Granara de Willink. Credits: Lyle J. Buss, University of Florida

Figure 4. Adult male (dorsal view) and nymphs of the mealybug *Hypogeococcus pungens* Granara de Willink. Credits: Lyle J. Buss, University of Florida

festerianus, the species originally confused with *H. pungens*, has only one circulus. Also, the pink

Figure 5. Lateral view of an adult male of the mealybug *Hypogeococcus pungens* Granara de Willink. Credits: Lyle J. Buss, University of Florida

hibiscus mealybug and *H. margaretae* have only one circulus.

Figure 6. Slide mounted specimen of *Hypogeococcus pungens* Granara de Willink. Arrows pointing to three circuli. Credits: Greg Hodges, Florida Department of Agriculture and Consumer Services, Division of Plant Industry

More detailed taxonomic diagnosis information is available the va Scale Insects: Identification tools, images, and diagnostic information for species of quarantine significance provided by the USDA-ARS Systematic Entomology Laboratory.

Life Cycle

Very little biological or life history information is available specifically for this species. The primary available biological work is Suss and Trematerra (1986). Populations of *H. pungens* were found both on roots and host leaves or stems. Miller et. al. (2007) also report that *H. pungens* has been intercepted on the roots of Cactaceae.

Hosts

In comparison to other mealybug species, the host range for *H. pungens* is fairly limited.

- **Amaranthaceae** - *Achyranthes aspera* (Devil's horsewhip), *Gomphrena globosa* (globe Amaranth)
- **Cactaceae** - several species with reported genera including *Cereus*, *Eriocerus*, *Harrisia*, *Hickenia*, *Parodia*
- **Polygonaceae** - *Alternanthera pungens* (khaki weed), *A. bettzickiana* (joyweed)
- **Portulacaceae** - *Portulaca oleracea* (purslane, verdolaga, pigweed, little hogweed, or pusley), *P. quadrifida* (chickenweed)

Figure 7. Infestation of the mealybug, *Hypogeococcus pungens* Granara de Willink. Credits: Lyle J. Buss, University of Florida

Figure 8. Infestation of the mealybug, *Hypogeococcus pungens* Granara de Willink. Credits: Florida Department of Agriculture and Consumer Services, Division of Plant Industry

Economic Importance

Hypogeococcus pungens has a very limited host range and appears to be of little economic significance as a pest. Significant pest populations of *H. pungens* rarely occur in Florida. Although *H. pungens* was reported occurring on Cactaceae in Italy,

subsequent surveys have reported reduced populations (Longo 2009). Exceptions to the general non-pest status of *H. pungens* could occur in areas with rare or endemic Cactaceae hosts, particularly those that reproduce primarily by seeds instead of budding.

Due to its fairly limited host range, *H. pungens* was successfully introduced as a biological control agent for the cacti species *Eriocereus bonplandii*, *E. martini*, *E. tortuosus*, and *Acanthocereus pentago* in Australia (McFadyen and Tomley 1981). This species has also been introduced into South Africa for the control of *E. martini* (Moran and Zimmermann 1991). *Hypogeococcus pungens* may be a more effective biological control agent for cacti species that are primarily dependent upon seeds for reproduction. Moran and Zimmerman (1991) noted that the inhibition of flowering and fruiting by *E. martini* due to *H. pungens* limited spread of the pest weed. Although *H. pungens* was a successfully biological control agent for pest weeds in Australia and South Africa, limited natural dispersal abilities of *H. pungens* require constant reintroduction of the mealybug as new infestations of the cacti weed occur.

Management

For high populations of *H. pungens*, it is often important to confirm the identification of your species prior to implementing a management strategy. This species may be confused with other similar mealybugs, and it is particularly important to report and submit unusual damage in terms of pest frequency or host range to your local diagnostic lab.

The following identification resources are available for Florida residents:

- your local Cooperative Extension Service office
- University of Florida, IFAS Extension Insect Identification Laboratory
- University of Florida, IFAS Mealybug Web site
- University of Florida Distance Diagnostic and Identification System

For many mealybug species, natural enemies will often effectively manage populations. The

mealybug destroyer, *Cryptolaemus montrouzieri*, is commonly found in association with many mealybug species as a generalist predator. Moran and Zimmermann (1991) report the presence of a ladybird beetle, *Exochomus* sp. (Coleoptera: Coccinellidae) feeding on *H. pungens*, but detailed observations on the impact of this species on populations is not provided. For additional information on natural enemies see Beneficial Insects and Mites.

Natural enemy populations will be negatively impacted by any pesticide applications applied to the host for control of *H. pungens* or other species. Mealybug populations may also be controlled by horticultural oil applications, but natural enemy populations may be adversely impacted by this as well.

Additional mealybug management information is available at:

- Florida Insect Management Guide for houseplant arthropod pests
- Florida Insect Management guide for scale insects and mealybugs on ornamental plants

Acknowledgements

The authors acknowledge Renato Inserra, FDACS-DPI, for translation of the Suss and Trematerra (1986) manuscript and for communications with Santi Longo, University of Catania in Sicily, regarding status of *H. pungens* in Italy.

Selected References

- Granara de Willink MC. 1981. Nueva especie de *Hypogeococcus* Rau de Tucumán, República Argentina (Homoptera Pseudococcidae). (In Spanish; Summary In English). Neotrópica 27: 61-65.
- Hamon AB. 1984. A cactus mealybug, *Hypogeococcus festerianus* (Lizer y Trelles), in Florida. Florida Department of Agriculture and Consumer Services, Division of Plant Industry. Entomology Circular 263.

- Hodges A, Hodges G, Buss L, Osborne L. (2008). Mealybugs and mealybug look-alikes of the Southeastern United States. *North Central IPM Center*.
<http://www.ncipmc.org/alerts/phmb/mealybugs.pdf> (19 June 2009).
- Longo S. (2009). Personal Communication. University of Cantania in Sicily. Professor of Entomology Department.
- McFadyen RE, Tomley AJ. 1981. The successful biological control of *Harrisia cactus* (*Eriocereus martinii*) in Queensland. pp 139-143. In Wilson BJ and Swarbrick JT. (Editors), Proceedings of the 6th Australian Weed Conference. Queensland Weed Society for the Council of Australian Weed Science Societies, Australia. 235 pp. (2 vols.)
- Miller DR, Rung A, Venable GL, Gill RJ. (August 2007). Scale Insects: Identification tools, images, and diagnostic information for species of quarantine significance. *Systematic Entomology Laboratory* USDA-ARS.
<http://www.sel.barc.usda.gov/scalekeys/ScaleInsectsHome/ScaleInsectsHome.html> (19 June 2009).
- Moran VC, Zimmermann HG. 1991. Biological control of cactus weeds of minor importance in South Africa. *Agriculture, Ecosystems, and Environment* 37: 37-55.
- Süss L, Trematerra P. 1986. *Hypogeococcus festerianus* (Lizer y Trelles), nocivo alle Cactacee ornamentali in Leguria. (In Italian). *Informatore Fitopatologia* 10: 43-46.
- Unknown. (2006). Annual Report FY 2006. *Hawaii Department of Agriculture*.
http://hawaii.gov/hdoa/meetings_reports/legislative-reports/2007LegReports/Appendix%204%20-%20PI%20Annual%20Rpt..pdf (19 June 2009).
- Williams DJ. 1973. Two cactus-feeding mealybugs from Argentina (Homoptera, Coccoidea, Pseudococcidae). *Bulletin of Entomological Research* 62: 565-570.
- Williams DJ, Granara de Willink MC. 1992. *Mealybugs of Central and South America*. CAB International, London, England. 635 pp.