Boisduval Scale, *Diaspis boisduvalii* Signoret (Insecta: Hemiptera: Diaspididae)¹

Adriana Espinosa, Heidi Bowman, Amanda Hodges and Greg Hodges²

Introduction

The boisduval scale, *Diaspis boisduvalii* Signoret, is an economically important pest of orchids, and was reported as the most important insect pest of orchids in Florida (Dekle 1965). Miller and Davidson (2005) list boisduval scale as one of the 43 most serious worldwide armored scale pests.

Figure 1. Adult female boisduval scales, *Diaspis boisduvalii* Signoret. Photograph by: Lyle J. Buss, University of Florida

Distribution

The boisduval scale occurs throughout the tropics, and as a greenhouse pest in more temperate climates (Dekle 1965, Miller and Davidson 2005).

Description

Females are white to light yellow, approximately 0.05-0.09 inches (1.2-2.25 mm) in diameter, circular or oval in shape, and covered with a centrally located, white-transparent, flat circular or oval shed skin. When the scale cover is removed, a single, horn-like projection on either side of the body, near the head and thorax may be visible (Dekle 1965, Howard et al. 2001, Miller and Davidson 2005).

Males are oval to elongate in shape, with a white cover and marginal shed skin. Males measure approximately 0.04 inches (1mm) in length.

Males are oval to elongate in shape, with a white cover and marginal shed skin. Males measure approximately 0.04 inches (1mm) in length.

-
1. This document is EENY-467 (IN838), one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: November 2009. This document is also available on Featured Creatures Web site at <http://entomology.ifas.ufl.edu/creatures>. Please visit the EDIS Web site at <http://edis.ifas.ufl.edu> and the Entomology and Nematology Department Web site at <http://entnemdept.ifas.ufl.edu/>.
 2. Adriana Espinosa, former extension assistant; Heidi Bowman, graduate student; and Amanda Hodges, SPDN assistant director in entomology and training/education, University of Florida; Greg Hodges, Florida Department of Agriculture & Consumer Services, Division of Plant Industry.

Figure 2. Adult female boisduval scales, *Diaspis boisduvalii* Signoret. Photograph by: Lyle J. Buss, University of Florida

Figure 3. Adult female boisduval scales, *Diaspis boisduvalii* Signoret, with exuviae present. A centrally-located circle shape is part of the exuviae. Photograph by: University of Florida

Life Cycle

Each adult female may produce up to 200 eggs and live as long as seven months. Eggs hatch within five to seven days and become crawlers. Egg color depends upon maturity, and ranges from clear to yellow, and finally orange. The development from egg to the adult stage averages 33 days for males and 50 days for females (Miller and Davidson 2005).

Hosts

This species has been recorded in over 15 families and 65 genera in Florida alone, but is most commonly found on orchids and palms.

Figure 4. Yellow colored body of the adult female boisduval scale, *Diaspis boisduvalii* Signoret. The oval-shaped cover has been removed. Photograph by: Florida Department of Agriculture and Consumer Services, Division of Plant Industry

Figure 5. Cluster of male boisduval scales, *Diaspis boisduvalii* Signoret, on Cattleya leaf. Notice the "fluffy" or "cottony" appearance. Photograph by: Heidi M. Bowman, University of Florida

Figure 6. Male, female, and crawler life stages of boisduval scale, *Diaspis boisduvalii* Signoret, on Cattleya leaf. Photograph by: Heidi M. Bowman, University of Florida

Within the palm hosts (family Arecaceae) the following genera are susceptible (Howard et al. 2001, Miller and Davidson 2005):

- *Acoelorrhaphe*
- *Acrocomia*
- *Archontophoenix*
- *Areca*
- *Bactris*
- *Butia* - pindo palm
- *Caryota* - fishtail palms
- *Chamaedorea*
- *Chamaerops* - European fan palm or Mediterranean fan palm
- *Cocos* - coconut palm
- *Corypha* - gebang palm, buri palm or tailpot palm
- *Dictyosperma*
- *Dypsis*
- *Elaeis* - oil palms
- *Euterpe*
- *Howea*
- *Hyophorbe*
- *Latania* - latan palm
- *Livistona*
- *Nannorrhops* - mazari palm
- *Phoenix*
- *Ptychosperma*
- *Rhapidophyllum* - needle palm
- *Raphis*

- *Roystonea* - royal palms
- *Sabal* - palmetto or fan palms
- *Syagrus*
- *Thrinax*
- *Trachycarpus* - fan palms
- *Washingtonia*

Recorded orchid hosts (Orchidaceae) include the following genera:

- *Acineta*
- *Angraecum*
- *Anguloa*
- *Bletia*
- *Brassavola*
- *Brassia*
- *Brassocattleya*
- *Broughtonia*
- *Bulbophyllum*
- *Cattleya*
- *Caularthron*
- *Coelogyne*
- *Cycnoches*
- *Cymbidium*
- *Dendrobium*
- *Encyclia*
- *Epidendrum*
- *Laelia*
- *Maxillaria*
- *Miltonia*

- *Neofinetia*
- *Odontoglossum*
- *Oncidium*
- *Ornithidium*
- *Peristeria*
- *Pleurothallis*
- *Renanthera*
- *Rhynchosyilis*
- *Schomburgkia*
- *Sophronitis*
- *Stanhopea*
- *Trichopilia*
- *Vanda*
- *Xylobium*

Other recorded host plant families include:

- Agavaceae
- Amaryllidaceae - *Agave* sp.
- Anacardiaceae - *Mangifera indica*, mango; *Schinus* sp., pepper trees
- Araliaceae - *Hedera helix*, common ivy
- Asteraceae - *Baccharis* sp.
- Bromeliaceae - *Aechmea*, *Ananas*, *Aregelia*, *Billbergia*, *Bromelia*, *Catopsis*, *Guzmania*, *Neoglaziovia*, *Pitcairnia*, *Puya*, *Ronnbergia*, *Tillandsia*, and *Vriesia* species
- Cactaceae
- Cyperaceae
- Fabaceae - *Acacia*, *Baikiaea*, *Cassia*, and *Leucaena* species
- Heliconiaceae

- Lauraceae - *Persea* sp.
- Liliaceae
- Moraceae - *Ficus* sp.
- Musaceae - *Musa* sp.
- Rosaceae - *Rosa* sp.
- Rubiaceae - *Coffea*, *Dracaena*, and *Citrus* species
- Vitaceae - *Vitis* sp.

A full listing of recorded hosts for boisduval scale is available at: ScaleNet: A Database of Scale Insects of the World.

Damage

Boisduval scales are normally found on the leaves and stems of palms. For orchids, this scale tends to prefer the leaf midrib and the portion of the petiole that is covered by the sheath. Boisduval scale is also capable of infesting orchid pseudobulbs and all aerial portions of the plant, including bark and fruit (Miller and Davidson 2005).

Figure 7. Chlorosis and "pitting" caused by boisduval scale, *Diaspis boisduvalii* Signoret, feeding on *Cattleya* leaf. Photograph by: Heidi M. Bowman, University of Florida

Management

Various cultural, biological, and chemical control options may be effective for management. Nutritional problems may enhance a host's susceptibility.

Cultural control. Exclusion is the first measure to be taken to avoid boisduval scale infestation. Carefully examine all plants and propagative materials before purchasing them. However, buyer beware, it is easy to miss scales located underneath leaf sheaths and not all life stages are visible to the unaided eye. New plants should be isolated from collections and nursery stock for at least two weeks to ensure they are pest free. Spacing plants so that their leaves do not touch can help avoid boisduval crawlers moving from infested plants onto clean neighboring plants (Beardsley and Gonzalez 1975). The crawlers can also move from infested to clean plants through strong wind currents in greenhouses and rooms. If it is possible, isolate infested plants to avoid boisduval scale spread and begin a treatment regiment.

Biological control. *Coccidencyrtus* sp. (Hymenoptera: Encyrtidae) have been reported as parasites of boisduval scale (Miller and Davidson 2005, Tenbrink and Hara 1992)

Figure 8. *Coccidencyrtus* sp., a parasitoid of the boisduval scale, *Diaspis boisduvalii* Signoret. Photograph by: Florida Department of Agriculture and Consumer Services, Division of Plant Industry

Chemical control. Horticultural oils are often very effective for controlling scale insects, but product labels should be carefully followed. For example, dormant oils applied to the actively growing stage of a plant may result in a burning effect on the plant material. Because crawlers tend to establish themselves on the upper and lower leaf surfaces, near the base of the plant, and in leaf sheaths, thorough spray coverage is important.

For small collections, using 70% isopropyl alcohol and a gentle cloth or cotton swab to wipe away the scales can provide effective control. However, some soft-leaved orchids may be damaged by isopropyl.

The American Orchid Society has a helpful, online video demonstrating how to treat scales with alcohol (AOS Video Library 2008).

If a chemical application is used, remember that timing the application to target the immature or crawler life stage may be critical for appropriate control. As boisduval scale is an armored scale, the waxy covering may still be present even after it has been effectively killed.

Florida Insect Management Guide for ornamentals

American Orchid Society management recommendations

Selected References

AOS Video Library. (2008). Boisduval Scale. *American Orchid Society*.
http://www.aos.org/AM/Template.cfm?Section=Video_Library&CONTENTID=6350&TEMPLATE=/CM/ContentDisplay.cfm (28 October 2009).

Beardsley Jr JW, Gonzalez RH. 1975. The biology and ecology of armored scales. *Annual Review of Entomology* 20: 47-73.

Buss EA, Turner JC. (June 2006). Scale insects and mealybugs on ornamental plants. *EDIS*.
<http://edis.ifas.ufl.edu/MG005> (28 October 2009).

Dekle GW. 1965. *Arthropods of Florida and Neighboring Land Areas: Florida Armored Scale Insects*. Vol. 3. Florida Department of Agriculture & Consumer Services, Division of Plant Industry. Gainesville, FL.

Howard FW, Moore D, Giblin-Davis RM, Abad R. 2001. *Insects on Palms*. CABI Publishing, Wallingford, UK. 400 pp.

Johnson PJ. (November 2001). Scale. *American Orchid Society*.

<http://www.aos.org/AM/>

Template.cfm?Section=Search&template=/CM/

HTMLDisplay.cfm&ContentID=5671 (28 October 2009).

Miller DR, Gimpel ME. (2009). Diaspididae: Diaspidinae and Leucaspinae. *ScaleNet*.

<http://www.sel.barc.usda.gov/scalenet/scalenet.htm>
(3 September 2009).

Miller DR, Davidson JA. 2005. Armored Scale Insect Pests of Trees and Shrubs (Hemiptera: Diaspididae). Cornell University Press. Ithaca, NY. 456 pp.

Tenbrink VL, Hara AH. (March 1992). *Diaspis boisduvalii* (Signoret). *Crop Knowledge Master*.

http://www.extento.hawaii.edu/kbase/crop/Type/d_boisdu.htm (4 November 2009).