

Crapemyrtle Pruning¹

Gary W. Knox, Edward F. Gilman, Teagan Young, Ryan W. Klein, and Leynar Leyton²

What is a “crape murder”?

Pruning is one of the most controversial aspects of maintaining crapemyrtle. Traditionally, many crapemyrtles were routinely topped, leaving large branch and stem stubs. This practice has been called “crape murder” because most people dislike the winter appearance and many professionals believe the practice impacts crapemyrtle health and structural integrity. UF/IFAS research has clarified the effects of various crapemyrtle pruning practices that resulted in the recommendations in this publication.

When properly placed, crapemyrtle is a low-maintenance plant needing little or no pruning. Problems with overgrown, misshapen, or misplaced crapemyrtle can be reduced greatly with the proper selection of crapemyrtle cultivars, choice of high-quality plants at the nursery, and placement in the landscape. Unfortunately, landscape professionals and homeowners often must maintain crapemyrtles planted by others and, therefore, must deal with the consequences of poor cultivar selection and/or placement.

Why is pruning necessary?

Crapemyrtle may need little or no pruning if planted in full sun, away from walkways and roads, and in areas with room for plant growth. In these locations, trees form a round canopy that often extends to the ground (Figure 1).



Figure 1. Crapemyrtles form a round canopy when planted in full sun with room for plant growth.

Credits: Edward F. Gilman, UF/IFAS

Some reasons for pruning crapemyrtle (or any plant) are to improve plant structure and alter plant size, shape, and appearance according to human perceptions and landscape function. Lower limbs of crapemyrtle are removed to increase clearance for pedestrians or vehicles. Stems are cut to increase branching. Other pruning may be conducted to direct growth away from structures; stimulate flowering; and remove spent flowers, seed capsules, and dead or damaged branches and twigs.

1. This document is ENH1138, one of a series of the Department of Environmental Horticulture, UF/IFAS Extension. Original publication date September 2009. Revised April 2010 and December 2024. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Gary W. Knox, emeritus professor (former author), Department of Environmental Horticulture, UF/IFAS North Florida Research and Education Center; Edward F. Gilman, emeritus professor (former author), Department of Environmental Horticulture; Teagan Young, graduate assistant; Ryan W. Klein, assistant professor, Department of Environmental Horticulture; and Leynar Leyton, assistant professor, Department of Environmental Horticulture, UF/IFAS North Florida Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

Before the introduction of disease-resistant cultivars, thinning the interior twigs and branches was a recommended pruning technique. This allowed better air movement to keep foliage dry and prevent or reduce powdery mildew disease. This practice is unnecessary for modern, disease-resistant crapemyrtle cultivars.

Types of Pruning

If pruning crapemyrtle is necessary, recommended types of pruning include tipping and pollarding. The pruning method of topping is not recommended. Topping is the drastic removal of large-diameter wood (typically several years old) that shortens all stems and branches. Topping is also known as heading, stubbing, rounding, and dehorning (Figure 2) or “crape murder.”



Figure 2. A crapemyrtle that has been topped. “Crape murder” consists of topping at position 1 in the first year, position 2 in the second year, and position 3 in the third year, whereupon topping in the fourth year often occurs below position 1 and repeats the sequence in subsequent years.

Credits: Edward F. Gilman, UF/IFAS

Tipping (or tip pruning) is different than topping in that cuts are made through smaller-diameter branches, which are typically one year old, on the outer edge of the plant canopy (Figure 3). Tipping is sometimes called “rounding over” or “pencil pruning” because cuts are made through stems about the diameter of a pencil. Tipping is very time-consuming, but it creates an attractive tree.

Pollarding involves initially making cuts through branches up to about three years old and then annually removing all or most sprouts back to the original cut. Over time,

woundwood and dead branch stubs form a “pollard head,” a swollen area at the end of the branch where sprouts are removed each year (Figure 4). Pollarding was used extensively in Europe to maintain trees at a small size. Pollarded trees are cut back to exactly the same size each year because cuts are made back to the pollard heads. The pollard head contains abundant stored energy with little decay and should not be removed.



Figure 3. Crapemyrtles that have been tip pruned.

Credits: Edward F. Gilman, UF/IFAS



Figure 4. A crapemyrtle that has been pruned using the pollarding method.

Credits: Edward F. Gilman, UF/IFAS

Pollarding, when done correctly, can allow a crapemyrtle to be maintained at a given height (the height of the pollard head) indefinitely. However, once started, pollarding requires annual pruning to remove regrown sprouts, increasing the maintenance required on an otherwise

low-maintenance plant. Topping and pollarding are considered “hard pruning” because of the larger-diameter pruning cuts that result.

Crapemyrtle Response to Pruning During the Growing Season

Crapemyrtle may be tip pruned during the growing season to promote flowering or to remove spent flowers and seed capsules. Tip pruning promotes flowering because dormant buds below the cuts are stimulated to grow, and flowers form at the tips of new growth. This practice is often unnecessary with newer cultivars because most have been selected for long seasons of bloom without repeated pruning. Also, tipping is impractical with large plants. Usually, hard pruning during the growing season is not recommended because of the impact on plant growth.

Crapemyrtle Response to Winter/Dormant Pruning Flowering

Topping and pollarding typically delay flowering up to one month compared to unpruned crapemyrtle. On some cultivars, topping and pollarding also shorten the season of bloom. In contrast, tip pruning does not affect flower timing or duration. With some cultivars, the greatest number of flowers occurs on unpruned crapemyrtle. The number of flowers decreases as pruning severity increases (i.e., as pruning cut diameter increases). However, pruned crapemyrtles may produce slightly larger flower panicles than unpruned plants.

Plant Shape and Appearance

Crapemyrtle shape becomes more upright as pruning severity increases. Although pruned plants are smaller than unpruned plants, new growth is extremely vigorous following hard pruning (Figure 5).

Hard pruning (topping and pollarding) also stimulates crapemyrtle sprouting from roots, upper stems, or the base of main stems. Most people dislike the appearance of sprouts on tree-form crapemyrtle, and typically, these sprouts are removed at least once during the growing season. If basal and root sprouts are not removed, one or more may form woody stems that eventually compete with existing main stems. These additional or competing stems, such as stems that rub against each other, may also result in poor form and structure. Hard pruning can cause excessive sprouting, so this method may indirectly

increase the maintenance requirements of an otherwise low-maintenance plant. Unpruned crapemyrtles form the fewest sprouts.



Figure 5. Crapemyrtle shape becomes more upright as pruning severity increases.

Credits: Edward F. Gilman, UF/IFAS

Long-stem sprouts emerge just below large-diameter cuts as a result of topping and pollarding. These sprouts usually develop into upright, unbranched stems that eventually flower, often bending under their own weight. Rain or wind storms can cause extreme bending of these long, unbranched stems; some will break because they are weakly attached to the main stem or because they are too slender.

Tip pruning dormant crapemyrtles to remove seed capsules is a personal aesthetic choice. Tip pruning does not induce sprouting or reduce flowering. However, this practice becomes increasingly labor intensive and impractical as a crapemyrtle grows larger. The presence of seed capsules in winter will not affect flowering the following growing season. Wind and rain eventually knock seed capsules off plants, usually prior to the growing season.

Spanish moss can be removed from the tree at any time to improve appearance. Although Spanish moss does not directly cause decline in the plant, heavy infestations can block sunlight from foliage, which could impact health.

Plant Size

Pruning will reduce plant size for at least part of the year. However, crapemyrtle regrowth often is so vigorous that

the plant grows to its original size within a couple of years. Many years of hard pruning are necessary to produce a plant smaller than an unpruned crapemyrtle.

Plant Health

Topping removes large amounts of starches and other food reserves stored within branches because large-diameter stems are removed. Topping dramatically reduces the size of the plant canopy, ultimately decreasing the plant's ability to produce these starches and other food reserves through photosynthesis. The large branch stubs caused by topping expose large areas of wood to insects and wood-rotting organisms, which weaken the plant's structure. Finally, topping results in many dead stubs throughout the tree (Figure 6).



Figure 6. Large branch stubs caused by topping.
Credits: Edward F. Gilman, UF/IFAS

Pollarding also results in large-diameter pruning cuts, but usually, these cuts are covered by woundwood faster than those on topped trees because the pruning cuts are smaller. Pollarding produces fewer dead branch stubs and less internal wood decay than topping. Vigorous, healthy crapemyrtles have a low risk of stem decay with either of these pruning methods because crapemyrtle rapidly and effectively walls off damaged wood. However, pruning large, old plants can cause more stress and decay and can even kill the plant. Tip pruning does not appear to significantly affect plant health.

Maintenance Time

With proper selection and placement, a crapemyrtle may need little or no pruning. Unpruned trees produce few sprouts to be removed during the growing season. Topping takes the least time to complete of the three pruning methods, especially during the dormant season. The structure of the plant is easy to see at this time, and stems

can be removed easily from the tree. Topped trees produce more summer sprouts, but total maintenance time is low even when adding time for de-sprouting. The reduced pruning time may have contributed to the apparent popularity of topping crapemyrtle despite the delayed flowering, unattractive form, weakened stem structure, and excessive sprouting that all result.

Time needed for tip pruning increases each year because of annual growth in size and the increased number of shoots that result from the prior year's tip pruning. Tip pruning requires the most time during the dormant season, and the most time overall, but very little summer de-sprouting is needed.

Pollarding crapemyrtle requires slightly more time than topping for dormant pruning because more cuts are required. Both topping and pollarding require about the same amount of summer de-sprouting time. One advantage of pollarding is that it requires about the same amount of winter pruning each year, regardless of age, because the number of sprouts remains about the same. The time it takes to top or tip a tree increases with the age and size of the plant. This likely explains why many crapemyrtles are topped below the original topping cut after several years of annual topping. In other words, five or six cuts several feet from the ground efficiently remove the entire top of the tree, whereas topping several dozen stems takes more time.

All pruning methods generate clippings and debris. Cleanup and disposal of the numerous clippings are another maintenance consideration. Factors other than total maintenance time may influence pruning practices. Topping requires less skill than other pruning methods, although it also results in delayed flowering, weak growth, and unattractive sprouting. Seasonally, landscape professionals and homeowners may have more time available during the dormant season than during the growing season, when mowing and other landscape maintenance practices may compete with de-sprouting time.

Recommended Crapemyrtle Pruning

Crapemyrtles selected to match the landscape site require little or no pruning. A crapemyrtle that requires routine pruning to fit into its surroundings should be considered for replacement with a smaller-maturing cultivar.

If pruning is necessary, use the following recommendations:

- Pruning for safety may be done anytime. This may involve removing damaged or weak branches or pruning lower limbs for pedestrian and vehicle clearance and visibility.
- Pruning to improve plant structure, redirect growth, or alter plant shape and appearance should occur when plants are leafless and dormant—typically December through February. Although this can be accomplished at any time, the branching structure without leaves is visibly easier to determine appropriate branches for pruning.
- Prune to remove branches that are crossing or rubbing each other.
- Prune dead, damaged, or diseased branches at the branch collar.
- Remove vigorous branches growing toward the center of the canopy.
- Severe pruning should be performed late in the dormant period. Pruning too early might stimulate new growth that could be damaged by low temperatures.
- Pruning to reduce plant size is best accomplished with pollarding rather than topping. Pollarding results in fewer dead stubs and less stem decay. Pollarding annually maintains crapemyrtle at a set height and results in a predictable flower display. It is also convenient to cut branches back to the same position each year instead of pruning at the ever-increasing height of topped trees.

Proper selection of crapemyrtle cultivars, selecting high-quality plants at the nursery, and proper placement in the landscape may prevent the need for significant pruning.

Crapemyrtle Selection to Avoid Pruning

Selecting the right cultivar for your space is essential to avoid extensive pruning. The correct crapemyrtle cultivar should be chosen to fit the intended planting area (see Table 1 for recommendations). This ensures that crapemyrtles are well-suited to their location, reducing the need for pruning and maintaining their health and structural integrity. Properly selected crapemyrtles become low-maintenance plants, needing little to no pruning.

Resources

Gilman, E. F. 2015. “*Lagerstroemia* x ‘Tonto’ Tonto Crape Myrtle. Southern Trees Fact Sheets.” FPS-329. University of Florida Institute of Food and Agricultural Sciences. <https://edis.ifas.ufl.edu/publication/FP329>

Gilman, E. F., R. W. Klein, and G. Hansen. 2024. “*Lagerstroemia* x ‘Sioux’ Sioux Crape Myrtle.” Southern Trees Fact Sheets. FPS-328. University of Florida Institute of Food and Agricultural Sciences. <https://edis.ifas.ufl.edu/publication/FP328>

Gilman, E. F., and G. W. Knox. 2005. “Pruning type affects crape myrtle.” *Journal of Arboriculture* 31 (1): 49–54. <https://doi.org/10.48044/jauf.2005.006>

Gilman, E. F., G. W. Knox, and P. Gomez-Zlatar. 2008. “Pruning method affects flowering and sprouting on crapemyrtle.” *Journal of Environmental Horticulture* 26 (3): 164–170. <https://doi.org/10.24266/0738-2898-26.3.164>

Gilman, E. F., and D. G. Watson. 2014. “*Lagerstroemia* x ‘Acoma’ ‘Acoma’ Crapemyrtle.” Southern Trees Fact Sheets. ENH-503. University of Florida Institute of Food and Agricultural Sciences. <https://edis.ifas.ufl.edu/publication/ST343>

Gilman, E. F., and D. G. Watson. 2014. “*Lagerstroemia* x ‘Muskogee’ ‘Muskogee’ Crapemyrtle.” Southern Trees Fact Sheets. ENH-506. University of Florida Institute of Food and Agricultural Sciences. <https://edis.ifas.ufl.edu/publication/ST346>

Gilman, E. F., and D. G. Watson. 2014. “*Lagerstroemia* x ‘Natchez’ ‘Natchez’ Crapemyrtle.” Southern Trees Fact Sheets. ENH-507. University of Florida Institute of Food and Agricultural Sciences. <https://edis.ifas.ufl.edu/publication/ST347>

Gilman, E. F., and D. G. Watson. 2014. “*Lagerstroemia* x ‘Tuscarora’ ‘Tuscarora’ Crapemyrtle.” Southern Trees Fact Sheets. ENH-508. University of Florida Institute of Food and Agricultural Sciences. <https://edis.ifas.ufl.edu/publication/ST348>

Gilman, E. F., D. G. Watson, R. W. Klein, and D. R. Hilbert. 2024. “*Lagerstroemia indica* ‘Cherokee’ ‘Cherokee’ Crape Myrtle.” Southern Trees Fact Sheets. ENH-505. University of Florida Institute of Food and Agricultural Sciences. <https://edis.ifas.ufl.edu/publication/ST345>

Table 1. Crapemyrtle cultivar growth characteristics and ornamental features.

Name	Height (feet)	Spread (feet)	Flower Color	Features
Semi-Dwarf Varieties (6–12 feet tall)				
<i>Lagerstroemia indica</i> 'Cherokee'	10–15	6–10	Dark red	Striking summer flowers, attractive fall foliage, and good drought tolerance.
<i>Lagerstroemia</i> x 'Tonto'	5–10	5–8	Dark red	Resistant to powdery mildew and other diseases, providing a long season of vibrant color.
<i>Lagerstroemia indica</i> 'Zuni'	8–10	8–10	Lavender	Disease-resistant with a vase-shaped form, ideal for adding height without overwhelming small spaces.
Medium Varieties (10–20 feet tall)				
<i>Lagerstroemia</i> x 'Acoma'	10–15	6–10	White	Weeping form with arching branches, suitable for small spaces with its graceful appearance.
<i>Lagerstroemia</i> x 'Sioux'	15–20	10–15	Dark pink	Upright growth habit with vibrant flowers and high resistance to mildew.
<i>Lagerstroemia indica</i> 'Catawba'	12–15	10–15	Purple	Upright, vase-shaped growth with striking flowers and good mildew resistance.
<i>Lagerstroemia indica</i> 'Tuscarora'	15–20	10–15	Coral pink	Strong, upright growth with flowers and excellent resistance to leaf spot and mildew.
Large Varieties (20+ feet tall)				
<i>Lagerstroemia</i> x 'Muskogee'	20–25	15–25	Lavender	Light-colored blooms, an extended flowering period, and good disease resistance.
<i>Lagerstroemia</i> x 'Natchez'	25–30	15–25	White	Rapid growth rate, attractive exfoliating bark, large flower clusters, and extended bloom period.
<i>Lagerstroemia indica</i> 'Miami'	20–25	15–20	Pink	Vibrant flowers, vigorous growth, and good disease resistance.
<i>Lagerstroemia</i> x 'Tuscarora'	18–25	15–18	Coral pink	Strong, upright growth with flowers and excellent resistance to leaf spot and mildew.