

***Phoenix sylvestris*, Wild Date Palm¹**

Robert J. Northrop, Michael G. Andreu, Melissa H. Friedman, Mary McKenzie, and Heather V. Quintana²

Family

Arecaceae, palm family.

Genus

Phoenix is the Latin term for the Greek word that means "date palm."

Species

The species name *sylvestris* translates from the Latin term for "of the forest."

Common Names

**wild date palm, toddy palm, sugar date palm,
silver date palm**

Explanations of these common names are not known.

Description

This palm is native to India and southern portions of Pakistan. In both countries, it occurs in areas where there is sparse vegetation mainly

composed of scrub species and along flat lands where monsoons occur. Though slow growing, it can reach heights of up to 50 feet and grows well in areas of the United States where temperatures do not fall below 15°F. Leaves are pinnately compound and blue-green, and they can grow to 10 feet in length. Leaflets can reach approximately 18 inches long and grow opposite to one another on the rachis in such a way that the entire leaf looks flat. The petiole, or stem that attaches the leaf to the trunk, is 3 feet long and armed with spines. Young trunks bear triangular shaped leaf scars (the place where leaves once attached to the trunk) that become more diamond-shaped with age. On older trees, aerial roots tend to be present at the base of the trunk. Yellow inflorescences can reach lengths of 3 feet, are heavily branched, bear small white blossoms, and grow among the leaves. The oblong fruits are 1 inch long and occur in orange clusters, turning dark red to purple when mature.

Allergen

Each individual tree is either a male or a female (as is true for all species within this genus). Male

-
1. This document is FOR 246, one of a series of the School of Forest Resources and Conservation Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date May 2010. Visit the EDIS Website at <http://edis.ifas.ufl.edu>.
 2. Robert J. Northrop, Extension forester, Hillsborough County Extension; Michael G. Andreu, assistant professor of forest systems, School of Forest Resources and Conservation; Melissa H. Friedman, biological scientist, School of Forest Resources and Conservation, IFAS, University of Florida, Plant City Center; Mary McKenzie, research assistant, School of Forest Resources and Conservation; and Heather V. Quintana, research assistant, School of Forest Resources and Conservation

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer-Chancy, Interim Dean

trees are extremely allergenic because their pollen is air-borne, whereas female palms cause minimal to no allergies.

Riffle, R. L. and P. Craft. 2003. *An encyclopedia of cultivated palms*. Portland, OR: Timber Press, Inc.

Applications

Commercial/Practical

In India, sugar and alcohol are made from wild date palm flowers and jelly is made from the fruit.

Horticultural

The wild date palm is an attractive landscape specimen with its blue-green leaves, textured trunk, and yellow inflorescences. The canopy of this palm is dense, exhibits a round shape, and can provide light shade. This palm will thrive and show the best growth when it is planted in direct sunlight. Wild date palm is drought tolerant and prefers well-drained sandy soils, but it grows better when regularly watered. This species is susceptible to lethal yellowing disease (<http://edis.ifas.ufl.edu/pp146>), so it is best to avoid planting wild date palm where the disease is known to be present.

References

Borrer, D. J. 1988. *Dictionary of root words and combining forms* (2nd ed.). Mountain View, CA: Mayfield Publishing Company.

Coombes, A. 1994. *Dictionary of plant names: Botanical names and their common name equivalents*. Portland, OR: Timber Press.

Floridata.com. 2004. *Phoenix dactylifera*, Retrieved from http://www.floridata.com/ref/p/phoe_dac.cfm

Harrison, N. A. and M. L. Elliot. 2009. *Lethal yellowing (LY) of palm* (PP146). Gainesville, FL: UF-IFAS Florida Cooperative Extension Service. Retrieved from <http://edis.ifas.ufl.edu/pp146>

Meerow, A. W. 2004. *Betrock's guide to landscape palms* (9th ed.). Hollywood, FL: Betrock Information Systems.

Ogren, T. L. 2000. *Allergy-free gardening: The revolutionary guide to healthy landscaping*. Berkeley, CA: Ten Speed Press.