

Parasitoids of Dipteran leafminers, *Diglyphus* spp. (Insecta: Hymenoptera: Eulophidae)¹

Jian Li and Dakshina R. Seal²

Introduction

Diglyphus spp. wasps are promising biological control agents for agromyzid leafminers (Diptera: Agromyzidae). The species occurring in North America are *Diglyphus isaea* (Walker), *D. begini* (Ashmead), *D. websteri* (Crawford), *D. intermedius* (Girault), *D. pulchripes* (Crawford) and *D. carlylei* (Girault) (Lasalle and Parrella 1991, Stegmaier 1972).

Distribution

This genus of leafminer parasitoids occurs widely in Asia, Europe, North America, New Zealand and Northern Africa (Minkenberg 1989).

Description

Adult

The adult parasitoid is a tiny wasp 1.5–2 mm long, depending on the species. The head, thorax and dorsal abdomen are generally metallic green in color, while the eyes are red. The scutellum has two pairs of setae, a submarginal vein with more than two dorsal setae, and the funicle is 2-segmented (Lasalle and Parrell 1991).



Figure 1. Adult *Diglyphus* sp. on a bean leaf. Larvae in this genus are external parasitoids of dipteran leafminers. Photograph by: Jian Li, University of Florida

Egg

The adult female *Diglyphus* parasitoid stings the dipteran host larva to paralyze it. Then the female may lay one or more eggs on the late instar leafminer larva (Minkenberg 1986).

-
1. This document is EENY-484 (IN877), one of a series of the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date December 2010. This document is also available as a Featured Creature at <http://entomology.ifas.ufl.edu/creatures>. Please visit the EDIS website at <http://edis.ifas.ufl.edu>.
 2. Jian Li, and Dakshina R. Seal, Department of Entomology and Nematology, University of Florida-IFAS, Tropical Research and Education Center, Homestead, FL 33031

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer-Chancy, Interim Dean



Figure 2. Adult *Diglyphus* sp. on a bean leaf. Larvae in this genus are external parasitoids of dipteran leafminers. Photograph by: Jian Li, University of Florida

Larva

The parasitoid larva has three instar stages. The first instar larva is transparent, whereas second and third instars are yellowish. The parasitoid larva feeds externally on the leafminer larva, eventually killing the host. The parasitoid larva then pupates in the leaf mine before emerging as an adult. The development time is temperature dependent. *Diglyphus isaea* takes about 10 days at 25°C for complete development on both the American serpentine leafminer, *Liriomyza trifolii*; and the pea leafminer, *L. huidobrensis* (Bazzocchi et al. 2003).



Figure 3. *Diglyphus* sp. larvae, 1st, 2nd and 3rd instar respectively left to right. The larvae were removed from the mine of a bean leaf. Larvae in this genus are external parasitoids of dipteran leafminers. Photograph by: Jian Li, University of Florida

Pupa

The third instar larva pupates in the leaf mine. The pupa is initially both transparent and light green, but eventually turns black.

Hosts

Diglyphus spp. are primary ectoparasitoids of dipteran leafminers in the family Agromyzidae. However, a *Diglyphus* sp. was also recorded



Figure 4. Parasitoid *Diglyphus* sp. larva (top) feeding on a leafminer larva. The larva was removed from the mine of a bean leaf. Larvae in this genus are external parasitoids of dipteran leafminers. Photograph by: Jian Li, University of Florida



Figure 5. Two parasitoid *Diglyphus* sp. larvae (one is feeding horizontally on top, the second is positioned vertically to the right side of the larger host larva) feeding on a leafminer larva. The larvae were removed from the mine of a bean leaf. Larvae in this genus are external parasitoids of dipteran leafminers. Photograph by: Jian Li, University of Florida



Figure 6. Early stage of pupa of a *Diglyphus* sp. showing transparent and light green color, and red eyes (on right). The pupa was removed from the mine of a bean leaf. Larvae in this genus are external parasitoids of dipteran leafminers. Photograph by: Jian Li, University of Florida

parasitizing Lepidoptera (Lyonetiidae) larvae (Boucek and Askew 1968).

Economic Importance

Augmentative release of commercial *Diglyphus* spp. is used for controlling leafminers in greenhouses in North America and Europe. Bazzocchi et al.



Figure 7. Late stage of pupae of *Diglyphus* spp. turn black in color. In this image the head is to the right. Larvae in this genus are external parasitoids of dipteran leafminers. Photograph by: Jian Li, University of Florida

(2003) indicated that *D. isaea* parasitized at least 18 different agromyzid species. *Diglyphis isaea* is the most effective commercial biological control product for controlling the American serpentine leafminer, *Liriomyza trifolii*; the pea leafminer, *L. huidobrensis*; *L. bryoniae*; and the chrysanthemum leafminer, *Phytomyza syngenesiae* (Syngenta-bioline). Lasalle and Parrella (1991) indicated that *D. begini* also parasitizes *L. trifolii*, *L. huidobrensis* and *L. bryoniae* in North America. Kaspi and Parrella (2005) reported that the insecticide Abamectin has little impact on *D. isaea* adults or the larvae within the leaf mines of chrysanthemums.

Selected References

- Boucek Z, Askew RR. 1968. Palearctic Eulophidae (excl. Tetrastichinae). (Hym. Chalcidoidea). Index of Entomophagous Insects. La François, Paris. 260 pp.
- Bazzocchi GG, Lanzoni A, Burgio G, Fiacconi MR. 2003. Effects of temperature and host on the pre-imaginal development of the parasitoid *Diglyphus isaea* (Hymenoptera: Eulophidae). *Biological Control* 26: 74-82.
- Kaspi R, Parrella MP. 2005. Abamectin compatibility with the leafminer parasitoid *Diglyphus isaea*. *Biological Control* 35: 172-179.
- Lasalle J, Parrella MP. 1991. The chalcidoid parasites (Hymenoptera, Chalcidoidea) of economically important *Liriomyza* species (Diptera, Agromyzidae) in North America. *Proceedings of the Entomological Society of Washington* 93: 571-591.
- Minkenberg OPJM. 1989. Temperature effects on the life history of the Eulophid wasp *Diglyphus isaea*, an ectoparasitoid of leafminers (*Liriomyza*

spp.), on tomatoes. *Annals of Applied Biology* 115: 381-397.

Minkenberg OPJM, Van Lenteren JC. 1986. The leafminers *Liriomyza bryoniae* and *L. trifolii* (Diptera: Agromyzidae), their parasites and host plants: a review of Agriculture. University of Wageningen Papers 86: 1-50.

Stegmaier CE. 1972. Parasitic Hymenoptera bred from the family Agromyzidae (Diptera) with special reference to south Florida. *Florida Entomologist* 55: 273-282.

Syngenta-bioline. Dig-line i: Leaf miner control. *Syngenta Bioline Limited*. <http://www.syngenta-bioline.co.uk/controldocs/html/DiglyphusIsaea.htm> (20 December 2010).