

Big-Eyed Bugs, *Geocoris* spp. (Insecta: Hemiptera: Lygaeidae)¹

F. W. Mead²

Introduction

Big-eyed bugs are small insects (approximately 1/6 inch long) that occur in many parts of the world. They are generally regarded as beneficial because they prey upon numerous kinds of insect and mite pests of turf, ornamental and agricultural crops. Big-eyed bugs are among those insects receiving research attention in Florida (and elsewhere) for their value as predators. To aid in identification of big-eyed bugs in Florida, a key to adults and late instar nymphs is provided in this publication.

Distribution

Geocoris bullatus (Say), the large big-eyed bug, is widely distributed in the United States and Canada, from coast to coast. There are numerous Florida records from the northern border south to Key West. *G. punctipes* (Say) is primarily an austroriparian species common throughout Florida and ranging from New Jersey west to southern Indiana and Colorado south and southwest to Texas, Arizona, California, and Mexico. Other localities include Guatemala, Panama, and Hawaii. *G. uliginosus* (Say)



Figure 1. Adult bigeyed bug, *Geocoris* sp. Credits: University of Florida

ranges over most of the United States and southern Canada. In Florida, *G. uliginosus* is known at least as far south as Ft Myers.

Identification

Big-eyed bugs are small, oblong-oval lygaeids having the head broader than long and prominent eyes which curve backward and overlap the front of

1. This document is EENY-252 (originally published as DPI Entomology Circular 121), one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: November 2001. Revised: October 2004. Reviewed: March 2011. This document is also available on Featured Creatures website at <http://entomology.ifas.ufl.edu/creatures>. Please visit the EDIS website at <http://edis.ifas.ufl.edu> and the Entomology and Nematology Department website at <http://entnemdept.ifas.ufl.edu/>.

2. F. W. Mead, Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville, FL.



Figure 2. Adult bigeyed bug, *Geocoris* sp., feeding on a whitefly nymph. Credits: Jack Dykinga, USDA




Figure 4. A bigeyed bug, *Geocoris punctipes* (Say). Credits: Division of Plant Industry



Figure 3. The large bigeyed bug, *Geocoris bullatus* (Say). Credits: Division of Plant Industry

the pronotum; the stylus has a longitudinal groove. These features can be seen on nymphs as well as adults and serve to separate bigeyed bugs from similar bugs. A distinguishing feature of adult




Figure 5. A bigeyed bug, *Geocoris uliginosus* (Say). Credits: Division of Plant Industry

big-eyed bugs is the very short or absent claval commissure. Lygaeids such as chinch bugs (*Blissus* spp.), false chinch bugs (*Nysius* spp.), and pamerars (*Pachybrachius* spp.) are sometimes confused with big-eyed bugs, but these genera have a claval commissure approximately half as long as the scutellum. Also the head has more of a triangular shape in these lygaeids. Caplan (1968) emphasized the need for turf specialists to distinguish between big-eyed bugs and chinch bugs. Misidentification could result in a chinch bug spray directed against geocorines, resulting in needless loss of money and beneficial insects.




Figure 6. A comparison of the claval commissure on a bigeyed bug, *Geocoris* sp., and a pamera bug, *Pachybrachius* sp. Credits: Division of Plant Industry

The following key to geocorines in Florida does not include two species of *Hypogeocoris*, which have been reported in Florida, but apparently are scarce or rare. Some minor variations in *Geocoris bullatus* (Say) and *G. uliginosus* (Say) have been formalized as subspecies, but will not be considered in this circular.

Key to Species of Adult Geocorinae in Florida

1. Beak with segment I (basal) longer than II; head (except in *G. punctipes*) punctulate (with small pits) or rugulose (minutely wrinkled) (*Geocoris*) 2

1'. Beak with segment I subequal to or shorter than II; head smooth, impunctate, shining (*Hypogeocoris*)

2. Nearly all black above except for pale border along each side; scutellum entirely black (Fig. 5) *G. uliginosus* (Say) 1832

2'. Mostly pale above; scutellum with a pair of pale areas or spots (Fig. 3, 4) 3

3. Scutellum with a pair of prominent, smooth (impunctate), calloused basolateral, pale spots, the spots sometimes extending posteriorly (extension punctulate); pronotum with a pair of somewhat lunate impunctate callosities, usually shiny black but sometimes invaded by various amounts of yellow; head smooth, polished, not at all granulose; inner posterior margin of corium not marked with fuscous or at most weakly so; groove of tylus extending back onto vertex and crossed near middle by an arcuate, transverse sulcus (Fig. 7), length 3.5 to 4.2 mm (Fig. 4) *G. punctipes* (Say) 1832




Figure 7. Dorsal aspect of the head of the bigeyed bug *Geocoris punctipes* (Say), showing the transverse sulcus. Credits: Division of Plant Industry

3'. Scutellum with a pair of punctate, non-calloused, submedial, pale-yellow areas; the shape and extent of pale areas variable but usually oblong and partially angulate; pronotum with the pair of impunctate callosities nearly round, pale-yellow; head granulose; inner posterior margin of corium marked with two fuscous "spots", the posterior one larger; groove of tylus not extending back onto vertex; vertex lacking transverse sulcus; length 3.0 to 3.5 mm (Fig. 3) *G. bullatus* (Say) 1832, the large bigeyed bug.

Key to Late Instar Nymphs of Florida Species of *Geocoris*

To make sure a nymph is a lygaeid, consult the key by Herring and Ashlock (1971) and/or the one by DeCoursey (1971). To key a lygaeid nymph to genus, consult Sweet and Slater (1961).

1. Dorsal ground color of head and thorax dark brown (Fig. 8) *G. uliginosus* (Say)

1'. Dorsal ground color of head and thorax pale (irregular dark spots often present) 2




Figure 8. Nymph of *Geocoris uliginosus* (Say), a bigeyed bug. Credits: Division of Plant Industry

2. Mesothoracic wing pads (developing forewings) either unmarked or with only one apical brown spot; scutellum with two pairs of linear brown marks, sometimes coalesced into one large pair, these marks basolaterad; pronotum usually with three pairs of brown spots, variable in shape and degree of pigmentation, often inconspicuous; antennal segments I-III each with prominent dorso-apical pale spot (Fig. 9) *G. bullatus* (Say), the large bigeyed bug

2'. Mesothoracic wing pads each with four or five brown marks (usually three basal streaks, one middle dot, and one prominent apical spot); scutellum

with three to five pairs of dark brown marks (usually four pairs), the most prominent pair near middle; pronotum with five to six pairs of conspicuous dark brown irregular spots; antennal segments I-III each without dorsoapical pale spot (Fig. 10) *G. punctipes* (Say)



Figure 9. Nymph of the large bigeyed bug, *Geocoris bullatus* (Say). Credits: Division of Plant Industry




Figure 10. Nymph of *Geocoris punctipes* (Say), a bigeyed bug. Credits: Division of Plant Industry

Bionomics

The literature on the food habits and life histories of *Geocoris* spp. is too extensive for more than a token review here. The most abundant big-eyed bug in Florida and the southeastern United States is *G. punctipes* (Say). McGregor and McDonough (1917) reported the life history of *G. punctipes* at Batesburg, South Carolina, finding the average development time from egg to adult was 30 days. Nymphs consumed an average of 47 mites, and adults an average of 83 "red spider" mites on cotton per day. York (1944) reported that adult *Geocoris* required either free moisture or plant moisture as well as insect prey. Sweet (1960) found that *Geocoris* adults can survive on sunflower seeds and water, without insect food.

Dumas et al. (1962) found more *G. punctipes* in the morning than at midday or evening, either by sweep net sampling or complete plant examination in Arkansas soybean fields. Bell and Whitcomb (1964) reported that in Arkansas *G. punctipes* and *G. uliginosus* were among the most abundant and important predators of bollworm eggs, *Helicoverpa* (= *Heliothis*) *zea* (Boddie) on cotton from mid-June until September. Whitcomb and Bell (1964) reported that big-eyed bugs preyed upon aphids, plant bugs, eggs, and young larvae of the bollworm and cotton leafworm in Arkansas cotton fields. On the negative side, however, the prey occasionally were beneficial species (*Orius* spp.). Champlain and Sholdt (1967) reported on the life history of *G. punctipes* in the laboratory. Lingren et al. (1968) reported *G. punctipes* was a more effective predator than *G. uliginosus* against *Heliothis* spp. Stoner (1970) found that *G. punctipes* apparently needed prey for proper development and fecundity. Orhanides et al. (1971) reported that *G. punctipes* was an effective predator of the pink bollworm, *Pectinophora gossypiella* (Saunders) in southern California cotton fields. Tamaki and Weeks (1972) listed 46 references, itemized from the literature the prey list of *Geocoris* spp., and presented extensive research results from a five year project on *Geocoris* in the Yakima Valley of Washington, including data on *G. bullatus*.

Selected References

- Anonymous. (May 2000). Color photographs of adult *Geocoris bullatus* (Say) and *Geocoris uliginosus* (Say). *Cedar Creek Natural History Area*. <http://www.cedarcreek.umn.edu/insects/album/020007018ap.html> (29 November 2001).
- Bell KO, Whitcomb WH. 1964. Field studies on egg predators of the bollworm, *Heliothis zea* (Boddie). *Florida Entomologist* 47: 171-180.
- Blatchley WS. 1926. Heteroptera or true bugs of Eastern North America, with especial reference to the faunas of Indiana and Florida. Nature Publishing Co., Indianapolis. 1116 p.
- Caplan I. 1968. Some chinch bugs aren't. *Weeds Trees and Turf* 7: 31-32.
- Champlain RA, Sholdt LL. 1967. Life history of *Geocoris punctipes* (Hemiptera: Lygaeidae) in the laboratory. *Annals of the Entomological Society of America* 60: 883-885.
- DeCoursey RM. 1971. Keys to the families and subfamilies of the nymphs of North American Hemiptera-Heteroptera. *Proceedings of the Entomological Society of Washington* 73: 413-428.
- Dumas BA, Boyer WP, Whitcomb WH. 1962. Effect of time of day on surveys of predaceous insects in field crops. *Florida Entomologist* 45: 121-128.
- Herring JL, Ashlock PD. 1971. A key to the nymphs of the families of Hemiptera (Heteroptera) of America north of Mexico. *Florida Entomologist* 54: 207-212.
- Lingren PD, Ridgway RL, Jones SL. 1968. Consumption by several common arthropod predators of eggs and larvae of two *Heliothis* species that attack cotton. *Annals of the Entomological Society of America* 61: 613-618.
- McGregor EA, McDonough FL. 1917. The red spider on cotton. *USDA Bulletin* 416: 41- 43.
- Orhanides GM, Gonzalez D, Bartlett BR. 1971. Identification and evaluation of pink bollworm

predators in southern California. *Journal of Economic Entomology* 64: 421-424.

Stoner A. 1970. Plant feeding by a predaceous insect, *Geocoris punctipes*. *Journal of Economic Entomology* 63: 1911-1915.

Sweet MH. 1960. The seed bugs: A contribution to the feeding habits of the Lygaeidae (Hemiptera: Heteroptera). *Annals of the Entomological Society of America* 53: 317-321.

Sweet MH, Slater JA. 1961. A generic key to the nymphs of North American Lygaeidae (Hemiptera: Heteroptera). *Annals of the Entomological Society of America* 54: 333-340.

Tamaki G, Weeks RE. 1972. Biology and ecology of two predators, *Geocoris pallens* Stål and *G. bullatus* (Say). U.S. Department of Agriculture Technical Bulletin 1446. 46 p.

Whitcomb WH, Bell K. 1964. Predaceous insects, spiders, and mites of Arkansas cotton fields. *Arkansas Agricultural Experiment Station Bulletin* 690. 84 p.

York GT. 1944. Food studies of *Geocoris* spp., predators of the beet leafhopper. *Journal of Economic Entomology* 37: 25-29.