

# Luffa—an Asian Vegetable Emerging in Florida<sup>1</sup>

Yucong Xie, Guodong Liu, Yuncong Li, and Kati Migliaccio<sup>2</sup>

*Luffa* is the genus name of several tropical and subtropical plants in the cucumber family (*Cucurbitaceae*) (Stephens 2015). Alternatively spelled “Loofa” or “Loofah,” the name is derived from the plant’s use as a material for sponges and dish cloths for bathing and cleaning dishware (Oboh and Aluyor, 2009). The name “Luffa” was introduced to Western botany nomenclature by the German anatomist and botanist, Johann Veslingius (1598–1649). A French botanist, Joseph Pitton de Tournefort (1656–1708), introduced the formal botany genus name “Luffa”. The species is native to tropical Asia, but has also been grown in Egypt since the late medieval era. While *Luffa* is widely cultivated in Asia, particularly in China and Vietnam, the gourd has recently become an attractive commodity in US vegetable markets, including Asian groceries in New York, California, and Florida.

## Description

There are two common species in the *Luffa* genus: smooth luffa (*Luffa aegyptiaca* Mill.), also called sponge gourd, and angled luffa (*Luffa acutangula* (L.) Roxb.). In the 16<sup>th</sup> century, Vestingius named luffa “Egyptian cucumber.” Common English terms for angled luffa include Chinese okra, dish cloth gourd, ribbed loofah, ridged gourd, sponge gourd, silky gourd, sinkwa towelsponge (USDA 2016), strainer vine, and vegetable gourd. Fresh produce of both smooth and angled luffa varieties are emerging in Florida vegetable markets. Smooth luffa, as its name suggests, produces green, squash-like fruit (Figure 1). Common cultivars

for smooth luffa include ‘Smooth Boy’, ‘Smooth Beauty’, and ‘South Winner’. For angled luffa, the fruit slightly resembles a cucumber or zucchini with 8 to 10 longitudinal ridges and ribs and a dark green color (Figure 2). Prominent cultivars include ‘Lucky Boy’, ‘Hybrid Green Glory’, ‘Summer Long’, and ‘Hybrid Asian Pride’.




Figure 1. (A) Fresh immature fruit of smooth luffa; (B) The fruit are sometimes wrapped with Cling Wrap to maintain moisture after harvest.

Credits: Guodong Liu, UF/IFAS (Left); Cuiping Huang, Hubei, China (Right)



Figure 2. (A) Fresh immature fruit of angled luffa; (B) The fruit of angled luffa have a longer shelf life and are more tolerant to shipping than those of smooth luffa. The angled luffa is more popular in Florida’s commercial farms for Asian vegetable crops.

Credits: Guodong Liu, UF/IFAS

Luffa is a fast-growing, long-season, and warm-climate crop that prefers neutral to slightly alkaline soil conditions with high moisture, growing annually in tropical and subtropical

1. This document is HS1285, one of a series of the Horticultural Sciences Department, UF/IFAS Extension. Original publication date September 2016. Visit the EDIS website at <http://edis.ifas.ufl.edu>.

2. Yucong Xie, graduate student, Horticultural Science Department; Guodong Liu, assistant professor, Horticultural Science Department; Yuncong Li, professor, UF/IFAS Tropical Research and Education Center, Department of Soil and Water Sciences; and Kati Migliaccio, professor, Department of Agricultural and Biological Engineering; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county’s UF/IFAS Extension office.

conditions. The stem is glabrous and ribbed and can grow as long as 23 feet. Tendrils have 3 to 5 branches with stems 2 to 3 inches in length and branches 3 to 4 inches in length (Figure 3). A trellis is needed for optimum fruit production and to cultivate straighter fruits (Zong et al. 1993).



Figure 3. Tendrils of luffa for climbing.  
Credits: Guodong Liu, UF/IFAS

## Cultivation Method

Mature luffa seeds are typically dark and hard, resembling watermelon seeds in size and shape. Immature seeds typically have a lighter color and are softer than mature ones (Figure 4).




Figure 4. (A) Immature seeds; and (B) mature seeds of luffa.  
Credits: Cuiping Huang, Hubei, China

The leaves of luffa are alternating (Figures 5 and 7) and palmate like, a pattern which is similar to squash and gourd plant foliage, or other species belonging to the cucumber family. A fully developed leaf is typically up to 12 inches wide and 5 inches long, and has an acute-end lobe, serrated edges, and is hairless (Figure 5).




Figure 5. (A) Cotyledon or seed leaf; and (B) true leaf of luffa.  
Credits: Cuiping Huang, Hubei, China

Both direct seeding (Figure 6) and transplanting are used for growing luffa in Florida. Seedlings can be transplanted once they have two or three true leaves (Figure 7). Direct

seeding is fast and can save labor, but it cannot be used before the frost-free season ends. Transplanting requires that the seedlings be ready and is more labor-consuming, but this method can expand the growing season and provide an earlier harvest with higher market value. The growth habit of luffa is similar to that of cucumber (*Cucumis stivaus* L.), except that *Luffa* must have well drained soil and requires high moisture conditions. Excessive water is detrimental, potentially slowing growth and increasing the risk of root disease. In addition, damping off may threaten young seedlings grown in cold and wet areas. For planting or transplanting, the distance between plants is usually 3 feet, and the distance between rows is 6 feet. Vine length is up to 30 feet. Vertical trellises (Figure 11) of 6 feet in height are recommended because they substantially increase fruit yield (Davis 1994). A mature luffa plant can bear about 6–7 fruit in ideal growing conditions. Luffa can also be grown in appropriate-size containers (Figure 7).



Figure 6. A grower seeding luffa on a commercial farm for Asian vegetable crops in northeast Florida in late July 2015.  
Credits: Guodong Liu, UF/IFAS



Figure 7. Seedlings ready for transplanting. For commercial luffa production, 72 cell pack insert-round trays can be used for establishing seedlings.  
Credits: Guodong Liu, UF/IFAS

The mature luffa leaves have petioles 1 to 3 inches in length. The leaves have 3 to 7 lobes depending on soil fertility. Lamina shape is ovate or broadly ovate. Fully developed laminae are 3 to 6 inches in length and 2 to 5 inches in width, with 3 to 7 broad, rounded, or obtuse lobes (Figure 8). Some varieties are hispid-covered on either the back or both sides of the leaf.



Figure 8. Fully developed leaves of luffa.  
Credits: Guodong Liu, UF/IFAS

Luffa inflorescences are monoecious: males (Figure 9) have elongate racemes with 10 to 16 flowers and females (Figure 10) have a solitary flower. Both male and female flowers are yellow or orange and 2 to 3 inches in diameter. The pollinated flowers develop cylindrical fruit (Figures 1 and 2), and only young, immature fruits, which require the hot growing season of about two to three months, can be harvested for consumption as a vegetable. Luffa seeds are elliptic: a quarter to three eighths of an inch in length, one eighth to a quarter inch in width, and one sixteenth of an inch in thickness. Seeds are smooth or slightly rugose. Immature seeds are white in color whereas mature seeds are dark brown or black (Figure 4). One mature luffa fruit contains 30 or more seeds (Partap et al. 2012).




Figure 9. Male flower buds of Luffa. (A) A single male flower; (B) Male racemes with more than 10 flowers.  
Credits: Guodong Liu, UF/IFAS

UF/IFAS is developing recommendations for fertilizer application rates. Before rates are established, growers can follow UF/IFAS recommendations for cucumber

production (Liu et al. 2016): target soil pH of 6.0 to 6.5; nitrogen supplementation of 150 lbs/acre; both  $P_2O_5$  and  $K_2O$  supplementation of 100–120 lbs/acre for low and 80 lbs/acre for medium nutrient concentrations based on soil testing. Plastic mulch is commonly used in Florida, but must be punctured to create holes large enough to avoid touching the seedlings. In northeast Florida, soil is rich in phosphorus. Growers should use liquid fertilizers of either 8-0-8 or 10-0-10 for fertigation through drip irrigation.



Figure 10. Female flower of Luffa.  
Credits: Guodong Liu, UF/IFAS



Figure 11. Luffa production with trellis in northeast Florida.  
Credits: Guodong Liu, UF/IFAS

Leafminer is a common insect pest targeting luffa (Figure 12). If an infestation is severe, contact your local UF/IFAS Extension agents or see *Vegetable Leafminer*, *Liriomyza sativae* Blanchard (*Insecta: Diptera: Agromyzidae*) <http://edis.ifas.ufl.edu/in507> (Capinera 2014).

Fruit rot, a common disease, can lead to losses if fruits are grown near the ground. If luffa fruit is not harvested before maturation, the fruit will become fibrous, resulting in a tough texture and bitter taste. Seeds from mature fruit can be collected and laid onto paper to dry. Once dry, the seeds can be kept in mason jars with tight-fitting lids, or glass canisters with gasketed lids and stored in cool, dark, and dry conditions for the next cropping.




Figure 12. Luffa leaves afflicted with leafminers.  
Credits: Guodong Liu, UF/IFAS

## Luffa Uses

The use of the luffa fruit is dependent on its growth stage. The immature fruit is a nutritious vegetable and is popular in Asian cuisine. Additionally, another species of luffa, called ball luffa (*Luffa operculata* (L.) Cogn.), is used as a fibrous scrubbing sponge (Figure 13) for household cleaning. Its common names include sponge cucumber and wild loofa (Filipowicz and Schaefer 2014). Ball luffa is rarely, if ever, eaten. Moreover, luffa sponges are recyclable, biodegradable and can be applied as an eco-friendly material in packaging, water absorption, and water waste treatment (Shen et al. 2012).



Figure 13. Luffa sponge.  
Credits: Guodong Liu, UF/IFAS

When harvested, immature luffa fruit can be cooked and eaten like cucumber or squash and is highly nutritious. Immature fruit can also be eaten raw. Abitogun and Ashogbon (2010) studied raw and defatted *Luffa* cylinders to determine the proximate mineral and energy composition, including crude proteins, moisture, fat content, ash content, crude fiber, and carbohydrates (Table 1). A high crude protein content was found in defatted luffa, confirming that luffa could be used to supply nutrients to other farm products (Abitogun and Ashogbon, 2010). However, while

less bitter leaves and roots can be consumed as food, the tough fibrous vines are not edible. Mature *Luffa* fruit look brown and can be harvested for seeds for next growing season. The sponges can be used for cleaning.

*Luffa* fruit are similar in taste to zucchinis and cucumbers, with some varieties exhibiting a sweeter taste than others. In Asian cuisine, edible luffa is usually cooked with olive oil and seasoned with soy sauce and cayenne pepper. A simple luffa recipe involves boiling the sliced luffa fruit as shown below. First, peel luffa fruit with a peeler (Figure 14a). Then, slice the fruit (Figure 14b). Next, boil the sliced fruit (Figure 14c) with approximately 2 fluid ounces of water per pound of sliced fruit. After boiling for 2 to 3 minutes, stir the salt into the boiling mixture. Now the luffa is ready to




Figure 14. (A) A peeler is useful for peeling luffa; (B) Peeled luffa; (C) Sliced luffa ready to be boiled; (D) Boiled luffa ready for consumption.  
Credits: Guodong Liu, UF/IFAS

be drained and served (Figure 14d). It can also be served as a luffa soup without draining.

*Luffa* fruit is rich in phosphorus, potassium, vitamin A, and vitamin C (Table 1), with benefit to the immune system and general health. *Luffa* has diuretic and purgative effects. It helps alleviate asthma and skin disease (Dhiman 2012). *Luffa* also contains triterpenoid saponins such as lucyosides and flavonoids such as apigenin. The seed extracts contain polypeptides, e.g., luffins. These physiologically active compounds have pharmacological effects such as anti-inflammation, anti-fungal, anti-allergy, anti-asthma, and anti-tussive (Partap et al. 2012).

Currently, luffa is primarily consumed by Asian populations, and consumption in Asian restaurants and homes is predominant in New York and other large cities with sizable Asian populations. A key economic venture for Florida growers would be to partner with distributors in these geographical locations. As cleansing sponges, processed mature luffa has more widespread value for bathroom and kitchen cleaning purposes and can be sold in gift shops. In 2008, it was reported that luffa sponges can sell at a range

of 1–5 dollars per unit and can bring up to 40 cents at whole sale. A grower can expect outputs in excess of 20,000 sponges per acre, reaching sizable profits (Jeanine 2008). In Florida, two production seasons are usual: from early March to late July and from late July to late November.

## Summary

Luffa is becoming an increasingly profitable vegetable crop in Florida. This fruit creates market and food diversity and also expands competitiveness of Florida's vegetable industry. Florida's unique geography and climate present clear advantages in luffa production. This geographical advantage is particularly apparent in the fall season, during which other states cannot grow this crop profitably due to limitations of climate conditions. Florida growers can grow this crop well into the mediate cold winter season, providing a unique economic opportunity.

## References and Further Reading

- Abitogun A.S and A.O. Ashogbon. 2010. "Nutritional Assessment and Chemical Composition of Raw and Defatted *Luffa cylindrica* Seed Flour." *Ethnobotanical Leaflets*, 14: 225–35.
- Arnott, J. 2007. *Luffa sponge gourd*. Cobb County Extension Service. Horticulture of University of Georgia. Available at : <http://www.caes.uga.edu/extension/cobb/anr/Documents/luffa.pdf>
- Capinera, J.L. 2014. *Vegetable Leafminer, Liriomyza sativae* Blanchard (Insecta: Diptera: Agromyzidae). IN507. Gainesville: University of Florida Institute of Food and Agricultural Sciences. Accessed September 15, 2016. <http://edis.ifas.ufl.edu/in507>
- Davis J.M. 1994. "Luffa Sponge Gourd Production Practices for Temperate Climates." *HortScience* 29(4): 263–266.
- Davis, J. 2008. *Commercial Luffa Sponge Gourd Production*. NC State University & A&T State University Cooperative Extension. Available at <http://content.ces.ncsu.edu/commercial-luffa-sponge-gourd-production>
- Dhiman, A. K., Gupta, D.K. Sharma, N.S. Gill, and A. Goyal. 2012. "A Review on the Medicinally Important Plants of the Family Cucurbitaceae." *Asian Journal of Clinical Nutrition* 4(1): 16–26.
- Egel, D.S., R. Weinzierl, P. O'Malley, R. Cloyd, B. Hutchinson, J. Pinero, and C. Welty. 2016. "Asian Vegetables."
- Midwest Vegetable Production Guide for Commercial Growers*. Purdue University.
- Filipowicz, N.H. and H. Schaefer. 2014. "Revisiting Luffa (Cucurbitaceae) 25 Years After C. Heiser: Species Boundaries and Application of Names Tested with Plastid and Nuclear DNA Sequences." *Systematic Botany* 39(1): 205–215.
- Jianhu S., M.X. Yi, X.D. Huang, S.W. Zhou, and R. Dong. 2012. "Mechanical properties of luffa sponge." *Journal of the Mechanical Behavior of Biomedical Materials*, 15:141 – 152.
- Lee, S. and J.G. Yoo. 2006. "Method for preparing transformed luffa cylindrica Roem." *World Intellectual Property Organization*.
- Liu, G.D., E. H. Simonne, K. T. Morgan, G. J. Hochmuth, M. Ozores-Hampton, and S. Agehara. 2016. *Chapter 2. Fertilizer Management for Vegetable Production in Florida*. CV296. Gainesville: University of Florida Institute of Food and Agricultural Sciences. Accessed September 15, 2016. <http://edis.ifas.ufl.edu/cv296>.
- Oboh, I. O. and E.O. Aluyor. 2009. "Luffa cylindrica - an emerging cash crop." *African Journal of Agricultural Research*, 4 (8): 684–688.
- Partap, S. A. Kumar, N.K. Sharma, and K.K. Jha. 2012. "Luffa Cylindrica: An important medicinal plant." *J. Nat. Prod. Plant Resour.* 2 (1):127–134.
- Shelagh, K., R. Anton, C. Sally, D. Samantha, and M. Nigel. 2013. "Growing Luffa. *Luffa cylindrica* and *L. acutangula* (syn *L. aegyptica*)." *Garden Organic the National Charity for Organic Growing*.
- Stephens, J. M. 2015. Gourd, Luffa—*Luffa cylindrica* (L.) Roem., *Luffa aegyptica* Mill., and *Luffa acutangula* (L.) Roxb. MV071. Gainesville: University of Florida Institute of Food and Agricultural Sciences. Accessed September 15, 2016. <http://edis.ifas.ufl.edu/mv071>
- USDA, Natural Resources Conservation Service. "Luffa acutangula (L.) Roxb." Available online at: <http://plants.usda.gov/core/profile?symbol=LUAC2>. Accessed on August 31, 2016
- Zong, R.J., C. Marita, and L. M. Leonard. 1993. "Angled luffa, bitter melon, fuzzy melon, yard-long bean-postharvest handling of Asian specialty vegetables under study." *California Agriculture*, 47(2):27–29

Table 1. Proximate compositions and mineral composition of raw and defatted *Luffa cylindrial*.

Nutrient	Unit	Value per 100 g		1 cup (1" pieces) = 95.0g		1 gourd = 178.0g	
<b>Proximates</b>							
Water	g	93.85		89.16		167.05	
Energy	kcal	20		19		36	
Protein	g	1.2		1.14		2.14	
Total lipid (fat)	g	0.2		0.19		0.36	
Carbohydrate, by difference	g	4.35		4.13		7.74	
Fiber, total dietary	g	1.1		1		2	
Sugars, total	g	2.02		1.92		3.6	
<b>Minerals</b>							
Calcium, Ca	mg	20	2	19	1.9	36	3.60
Iron, Fe	mg	0.36	3	0.34	2.8	0.64	5.33
Magnesium, Mg	mg	14	4	13	3.5	25	6.82
Phosphorus, P	mg	32	4	30	4.1	57	7.73
Potassium, K	mg	139	3	132	2.8	247	5.15
Sodium, Na	mg	3		3	0.0	5	0.00
Zinc, Zn	mg	0.07	1	0.07	0.7	0.12	1.26
<b>Vitamins</b>							
Vitamin C, total ascorbic acid	mg	12	15	11.4	13.9	21.4	26.18
Thiamin	mg	0.05	4	0.048	4.1	0.089	8.01
Riboflavin	mg	0.06	5	0.057	4.7	0.107	8.03
Niacin	mg	0.4	3	0.38	2.8	0.712	5.09
Vitamin B-6	mg	0.043	4	0.041	3.4	0.077	7.70
Folate, DFE	µg	7	2	7	1.8	12	3.10
Vitamin B-12	µg	0		0		0	
Vitamin A, IU	IU	410		390		730	
Vitamin E (alpha-tocopherol)	mg	0.1		0.1		0.18	
Vitamin D (D2 + D3)	µg	0		0		0	
Vitamin D	IU	0		0		0	
Vitamin K (phylloquinone)	µg	0.7		0.7		1.2	
<b>Lipids</b>							
Fatty acids, total saturated	g	0.016		0.015		0.028	
Fatty acids, total monounsaturated	g	0.037		0.035		0.066	
Fatty acids, total polyunsaturated	g	0.087		0.083		0.155	
Fatty acids, total trans	g	0		0		0	
Cholesterol	mg	0		0		0	
<b>Amino Acids</b>							
Other							
Caffeine	mg	0		0		0	

Source: USDA-ARS, National Nutrient Database for Standard Reference 28 available at <https://ndb.nal.usda.gov/ndb/foods/show/2972?fgcd=&manu=&facet=&format=&count=&max=35&offset=&sort=&qlookup=luffa> (2016).