

Florida's Bats: Florida Bonneted Bat¹

Holly K. Ober, Terry J. Doonan, and Emily H. Evans²

The Florida bonneted bat (Figure 1) is one of only two endangered species of bat in Florida, and the state's only endemic flying mammal ("endemic" means that it is found nowhere in the world but in Florida). With a wingspan of 20 inches (50 cm), it is Florida's largest bat and the third largest of all 48 species of bats in the United States. Fur color varies from brown to gray to black.



Figure 1. Florida bonneted bat (*Eumops floridanus*).

Credits: Merlin Tuttle, Merlin Tuttle's Bat Conservation <http://www.merlintuttle.com/>

Little is known about these rare bats. They occur only in southern Florida, with recent confirmation of presence in Miami-Dade, Broward, Collier, Hendry, Lee, Charlotte, Glades, Highlands, De Soto, and Polk Counties (Figure 2).




Figure 2. Florida bonneted bats can be found in just 10 counties in southern Florida.

Credits: Emily Evans

These bats roost (sleep during the day) in cavities or beneath peeling bark of live and dead pine trees (*Pinus elliotti*, *P. palustris*), royal palms (*Roystonea regia*), rock outcrops, bat houses, chimneys, and under barrel roof tiles. They roost alone or in groups of up to 50 individuals.

1. This document is WEC381, one of a series of the Department of Wildlife Ecology and Conservation, UF/IFAS Extension. Original publication date October 2016. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. Holly K. Ober, associate professor and wildlife Extension specialist, Department of Wildlife Ecology and Conservation; Terry J. Doonan, mammal conservation coordinator, Florida Fish and Wildlife Conservation Commission; and Emily H. Evans, mammal conservation biologist, Florida Fish and Wildlife Conservation Commission; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Florida bonneted bats typically leave their roosts shortly after sunset. They fly long distances to forage on insects at high altitudes. Bonneted bats' broad, funnel-shaped ears point forward, helping them detect insects at long range while in flight.

The Florida bonneted bat was listed as federally endangered in 2013 because of concerns over habitat loss, degradation, and modification caused by humans. Additional concerns include the species' small population size and restricted range, the small number of known colonies, their slow reproduction, and the relative isolation of separate populations of bonneted bats.



Figure 3. Free-tailed bats (those in the family Molossidae) have long tails that extend more than an inch beyond their tail membrane, like the bat on the left. In comparison, other bats have short, stubby tails, like the bat on the right (not a free-tailed bat).

Credits: Merlin Tuttle, Merlin Tuttle's Bat Conservation <http://www.merlintuttle.com/>

How do you know if you have found a bonneted bat?

First, check the rump to determine if a tail is visible. If you see a tail longer than one inch (Figure 3), the bat is one of Florida's three species of free-tailed bats (in the family Molossidae).

Read below to determine how to differentiate the three species of free-tailed bats found in Florida.

Florida Bonneted Bat (*Eumops floridanus*)

- Is found only in central and south Florida (as far north as Polk County)
- Roosts in pine tree cavities, palms, rock crevices, and man-made structures (beneath barrel tile roofs and in bat houses)
- Roosts in small groups (under 50 individuals)
- Has ears that are not fully separated from one another (they are joined at the base; Figure 4)

- Is the largest bat in Florida, about the size of a standard i-Phone when it is at rest and its wings are folded

Brazilian Free-Tailed Bat (*Tadarida brasiliensis*)

- Is found throughout the entire state of Florida, except the Keys
- Often roosts during the day in man-made structures (attics, stadiums, bridges, and bat houses)
- Typically roosts in large congregations (1,000–500,000 individuals)
- Produces a musky odor that is noticeable from some distance away
- Has ears that are fully separated from one another (not joined at the base like bonneted or velvety free-tailed bat ears; Figure 4)
- Is about the size of a standard business card when it is at rest and its wings are folded



Figure 4. The ears of the Florida bonneted bat are joined at the base (left), whereas there is a gap between the ears of the Brazilian free-tailed (right).

Credits: : Kathleen Smith, FFWCC, used with permission; Elizabeth Braun de Torrez, UF/IFAS

Velvety Free-Tailed Bat (*Molossus molossus*)

- Is found only in extreme south Florida (the Keys and perhaps the Everglades region)
- Typically roosts beneath roofs of man-made structures
- Roosts in the Keys in moderately-sized groups (50–300 individuals)
- Has ears that are not fully separated from one another (they are joined at the base)
- Is about the size of a standard business card when it is at rest and its wings are folded

If you find a bonneted bat, do not disturb it. Contact U.S. Fish and Wildlife Service in Vero Beach, FL (772-562-3909 or verobeach@fws.gov) to alert them to the location of the bat(s).

Additional Information

Florida Fish and Wildlife Conservation Commission.
[Living with bats.](#)

Ober, H. K., M. B. Main, and G. M. Allen. 2004. *Bats of Florida*. WEC186. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/UW203>.

Ober, H. K., and F. J. Mazzotti. 2008. *Conservation of Bats in Florida*. WEC247. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/UW291>.

Ober, H. K. 2008. *Effective Bat Houses for Florida*. WEC246. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/uw290>.

Ober, H. K. 2008. *Insect Pest Management Services Provided by Bats*. WEC245. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ufl.edu/UW289>.

Wisely, S. M., and H. K. Ober. 2015. *Facts about Wildlife Diseases: Rabies*. WEC239. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ufl.edu/UW282>.

Ridgley, F. N. 2015. Florida bonneted bat house plans. Miami Zoo. Available at <http://zoomiamiconservation.com/wp-content/uploads/2014/03/Florida-Bonneted-Bat-House-Plans.pdf>.