



UNIVERSITY OF
FLORIDA

EXTENSION

Institute of Food and Agricultural Sciences

SS-AGR-153

Liming of Agronomic Crops¹

E. B. Whitty and C. G. Chambliss²

The primary reason for liming acid soil is to improve the yield or quality of the crop being grown. It is difficult to determine the precise factor that is responsible for the improved growth after liming because a number of soil parameters change simultaneously as soil acidity is altered.

When mineral soil pH is below 5.5, aluminum toxicity seems to be the main reason for reduced plant growth. Organic soils contain little Al, thus plants can tolerate much lower pH levels on those soils without adverse effects. Many Florida soils are low in magnesium (Mg) and application of dolomitic limestone serves two purposes: it raises the soil pH and it provides Mg as a nutrient element.

On the other hand, excessive liming can be detrimental. Many Florida soils are quite low in manganese and deficiencies of Mn can occur in soils that are over-limed. The problems can begin to appear any time soil pH is raised above 6.3 or so, depending on the level of Mn present and the crop being grown.

Some physiological disorders of plants, such as frechning of tobacco, are associated with high levels of lime. Certain plant diseases, such as black shank of

tobacco, are more virulent as the soil pH increases above pH 5.8.

It has been noted in many Florida fields that are routinely irrigated from deep wells, the soil pH may not decline over time and may actually increase in some instances. Since irrigation water drawn from limestone aquifers contains low levels of dissolved calcium carbonate, this added lime accumulates over time and affects soil pH. Use of soil samples as described below can indicate if irrigation water contributes to the soil pH. Also, the need for lime can be affected by the source and amount of fertilizer applied. Again, a soil test can help reveal the practical effects on soil pH and the need for lime, if any.

In order to obtain the maximum benefits from liming and few, if any, of the detriments, it is necessary to plan a liming program. Soil factors and plant factors must be taken into account in determining the type and quantity of lime to apply.

The first step is to take a soil sample that is a representative of the field and have it tested by a laboratory which runs a lime requirement test. Since interpretation of soil test results are dependent on the test used and the field correlations of the test, no interpretation will be made here. Refer to SL-129

1. This document is SS-AGR-153, one of a series of the Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. First printed December 1992. Revised April 2002. Please visit the EDIS Web site at <http://edis.ifas.ufl.edu>.
2. E. B. Whitty, professor and C. G. Chambliss, associate professor, Agronomy Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. It is not a guarantee or warranty of the products named and does not signify that they are approved to the exclusion of others of suitable composition.

The Institute of Food and Agricultural Sciences is an equal opportunity/affirmative action employer authorized to provide research, educational information and other services only to individuals and institutions that function without regard to race, color, sex, age, handicap, or national origin. For information on obtaining other extension publications, contact your county Cooperative Extension Service office. Florida Cooperative Extension Service/Institute of Food and Agricultural Sciences/University of Florida/Christine Taylor Waddill, Dean.

UF/IFAS Standardized Fertilization

Recommendations for Agronomic Crops for the target pH for agronomic crops.

The decision of whether to use dolomitic or calcitic ("hi cal") lime should be based primarily on the cost of the material to the producer. When both lime and Mg are needed, dolomite can serve as the liming material and as a source of the nutrient Mg. However, if the cost of dolomite is significantly higher than calcite, the producer should consider the alternative of applying calcite as the liming material and Mg in the fertilizer. Application of dolomite as a source of Mg without regard to the liming effect can lead to other nutritional problems in soils with pH above 6.3.

Producers frequently may have access to by-product materials that can serve very well for liming agricultural land if the nature of the material is understood and proper precautions are followed. Lime from municipal water treatment plants is an example. Some suggestions follow about the handling and use of lime from water treatment plants:

1. As received from the water treatment plant, the lime usually has the consistency of a thick paste. Pile and allow to dry before attempting to spread.
2. Turn with a front-end loader to promote drying. Spread before completely dry and on a calm day to minimize drift.
3. Use about 80% as much material as you would agricultural limestone. It will react quickly due to its fineness and thus carry more potential for overliming if not properly used.

Materials sold as aglime are covered by the Florida Commercial Fertilizer Law and must meet specifications of fineness of grind, carbonate equivalence, and Mg content (in the case of dolomite). This affords some consumer protection. Lime by-products are not covered by the law and the consumer must realize more personal responsibility when dealing with such products. Liming is probably the most important soil fertility practice on strongly acid mineral soils. However, many field crops in Florida produce just as well on moderately acid soils as they do on only slightly acid soils.

LOWERING SOIL pH

Soil pH is sometimes too high for optimum growth and yield of particular plant species. **Caution** : Most plant species are tolerant to a wide range of soil pH. Do not attempt to lower soil pH unless there is evidence that plant growth is being adversely affected by the pH.

If the source of the high pH is naturally occurring carbonates (ex. the rockland soils of Dade County or soil containing limestone outcroppings), it is impractical to lower the soil pH on a field-wide basis. In those situations, application of elemental sulfur and micronutrients together in a band is recommended. The micronutrients will remain soluble in the acid band and principal adverse effects of high pH may be avoided.

If the soil pH is too high as a result of excessive liming, take note and prevent a re-occurrence of this costly mistake. The soil pH will gradually become more acid with time and unless the crop is being adversely affected, time is the best cure for over-limed soil in Florida.

When a more rapid lowering of soil pH is desired, elemental sulfur broadcast and worked into the soil will hasten acidification. **Caution** : Sulfate forms of sulfur will not lower pH. Elemental sulfur (ex. ag grade sulfur, wettable sulfur, flowers of sulfur) is acted upon by the soil microorganisms and sulfuric acid is produced. It is the acid, not the sulfate, that neutralizes the excess carbonate in the soil. A rate of 500 lb/A sulfur is recommended for the unfortunate cases of overliming. The effect on soil pH will probably be slow because of microlinal action.