



Management and Cultural Practices for Peanuts¹

D.L. Wright, B. Tillman, E. Jowers, J. Marois, J. A. Ferrell, T. Katsvairo, and E. B. Whitty²

Introduction

Peanuts have been one of the more profitable crops grown in the SE Coastal Plains for many years. Farm programs with the quota system made peanut production an attractive enterprise. The elimination of the quota system has led to a shift in acreage resulting in a 50% increase in total acreage in Florida since 2000. With current price, budgets should be studied carefully to determine where peanuts fit into your operation. Enterprise budgets for irrigated and dry land peanuts can be found at <http://nfrec.ifas.ufl.edu/Hewitt/budgets.htm>. Where peanuts are grown in new areas, yields are often much higher than where peanuts have been produced for many years. This is due to lower pest pressure and peanut yield responds more to good rotation than most other crops. Peanuts fit into many rotation schemes used by growers including corn, cotton, and other grass crops, and do best when planted after perennial grasses. Peanuts in the United States are grown for peanut butter, peanut candy, roasted peanuts, peanut oil, and other peanut products.

Peanut Production and TSWV

Peanut production has changed dramatically in the last 5-10 years and many of the management considerations are driven by tomato spotted wilt virus (TSWV). Factors affecting spotted wilt incidence and severity are shown below in approximate order of importance:

- History and location
- Cultivar
- Planting date
- Plant population
- Insecticide use
- Row pattern
- Tillage
- Herbicides

1. This document is SS-AGR-74, one of a series of the Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. First published February 2000. Revised September 2006. Please visit the EDIS Web site at <http://edis.ifas.ufl.edu>

2. D.L. Wright, professor, North Florida Research and Education Center--Quincy, FL; B. Tillman, assistant professor, North Florida Research and Education Center--Marianna, FL; E. Jowers, County Extension Director and Extension agent IV M.S.--Jackson County, FL; J. Marois, professor, North Florida Research and Education Center--Quincy, FL; J. A. Ferrell, assistant professor, Agronomy Department, Gainesville FL; T. Katsvairo, graduate assistant, research, North Florida Research and Education Center--Quincy, FL; and E. B. Whitty; professor emeritus, Agronomy Department, Gainesville, FL; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean

There are dramatic differences between years in the amount and severity of TSWV. Weather, thrips, and other factors shown in Table 1 will largely determine the severity of the disease. Variety selection and planting date are the most consistent factors that influence TSWV severity. However, weather conditions and other farm activities determine when peanuts are planted. Optimum planting dates have shifted over the years. Peanut planting once began in mid to late April and would end by mid May or until peanuts were planted. This date has continued to shift and is now focused on the period between May 11 and May 25. The Risk Index is modified each year as better information is gained to help farmers adapt to a shifting disease. Plant stands are critical for minimizing spotted wilt incidence and severity regardless of cultivar used. Although most insecticides have little, if any, effect on spotted wilt incidence, use of phorate (Thimet or Phorate) in-furrow at planting will reduce thrips damage. Twin rows and strip tillage have provided consistent suppression of spotted wilt in recent years as well as higher grades due to more uniform maturity. Use of Classic herbicide tends to increase severity of spotted wilt. Recent research in Florida has shown not only that strip tillage reduces TSWV as compared to conventional tillage but that perennial grasses killed in the fall reduces TSWV even further. Getting a good plant stand (4-6 plants/ row ft.) is critical to minimizing TSWV.

Table 1. Peanut Disease Risk Index

	Spotted Wilt	Leaf Spot	White Mold	Limb Rot
Rotation	N/A	25	25	20
Variety	50	35	25	25
Field History	N/A	10	15	10
Irrigation	N/A	10	5	10
Planning Date	30	10	5	5
Row Pattern	15	0	5	0
Tillage	15	10	0	5
Insecticide	15	N/A	N/A	N/A
Classic Herbicide	5	N/A	N/A	N/A
Plant Stand	25	N/A	N/A	N/A

Peanut Types

There are four market types of peanuts in the United States: runner, Virginia, Spanish, and Valencia. Runners have an intermediate size seed and are the primary type grown in Florida and the Southeast. Virginia peanuts have a large seed and are grown mostly in North Carolina, Virginia, and Texas, but some are grown in Florida. Spanish peanuts have small kernels and are grown mostly in Texas and Oklahoma. Virginia and Spanish peanuts are grown in Florida, but acreage is very minor in relation to runners. Valencia peanuts are usually about the same size as Spanish peanuts but are considered special-use peanuts, such as for boiling. Florida is a major producer of boiling peanuts which are usually planted early and throughout the year (February-July) to keep a continuous supply.

There are two vegetative types (growth habit) of peanuts: runner and bunch. Runner peanuts have a spreading growth habit because lateral branches are long and grow close to the ground. The bunch type grows mostly erect stems and does not spread out like the runner type. The peanut seed is called a *kernel*, and is found inside pods or hulls. Peanuts flower above ground and then a "peg" or reproductive stem grows down from the flower and enters the soil. The peanut pod forms at the end of this peg. The peanut leaf is made up of four leaflets. Peanuts are harvested by digging them from the ground with a digger-shaker-inverter, letting them partially dry in the field and then combining them. As the peanuts are dug, the vines are shaken to remove excess dirt and then the peanut vines are inverted so that the peanuts are exposed to the sunlight for quick drying. After they dry in the field, the nuts are removed from the vine with a peanut combine or picker. Peanuts are partially dried by natural drying in the field before combining. After combining, they are placed on trailers where heated air is passed through the peanuts to complete the drying process. Peanuts should be stored at a moisture content of 10 percent or less.

Variety Selection

Variety trial information on peanut can be found on the web at http://edis.ifas.ufl.edu/TOPIC_Peanut_Varieties for

Florida, <http://www.swvt.uga.edu/> for Georgia, and <http://www.ag.auburn.edu/aaes/communications/peanuts/ay274.pdf> for Alabama. Deciding on best varieties is a very important decision for yield and markets. Many varieties have resistance to diseases as well as having other good characteristics. There is often a 10-20% difference between some of the best varieties and the lower yielding varieties. Quality or grade may also vary making a difference in the prices received for the commodity. You have only one chance to make the right decision with the variety and yield limits are set with the variety. Runner peanuts are grown for the dry market and the most information will be available for them. Valencia's are grown because they grow rapidly and mature in less time than other types and also produce a peanut with a good flavor. Virginia type varieties have larger seed and the peanuts are especially suitable for roasting purposes. The runner type peanuts are also planted in home gardens because of their flavor and good yield. Seed for planting are relatively easy to find. The Spanish varieties are mostly grown in Texas because of being a short season crop, requiring less water and for the flavor and the small kernel. Whatever your choice of variety, consult your local seed dealer for availability. If you are growing peanuts for market purposes, be sure to check with the buying point where you intend to sell your peanuts to be sure they can process the variety you have chosen. Most of the peanuts grown for the Southeast are runner types and are shown in Table 2.

Plant Population

Peanut seed have always been expensive as compared to most row crops. Seed were more expensive under the quota system and as few as 3 seed/row ft. were planted with little difference in yield. After TSWV became a problem, it was found that seeding rates needed to be higher (5-6 seeds/row ft.) to keep disease levels lower. This meant going from 85-100 lbs/A seed to 100-120 lbs/A depending on seed size and variety. Current recommendations suggest 6 seed per ft. of row in single rows and 3 seed per ft. of row in twin rows to reduce TSWV levels. Where there are skips in the row, adjacent plants tend to get TSWV.

Crop Rotation and Land Selection and Preparation

Crop rotation is an important cultural practice that is recommended for all crops and especially for peanuts. Rotation is recommended to reduce the effects of pests (disease, nematode, insects, and weeds). In many cases, low value crops are the only alternative for rotating with high cash value crops like peanuts. In other cases, good systems are known and are not used because fencing or other costs will be involved. Extensive research has been conducted on rotation for most crops and their impacts on pests. Leafspot control is usually accomplished with a routine fungicide program but data shows that rotation with cotton or corn can reduce the amount and severity of disease. Recent data shows that rotations with bahiagrass can reduce the disease even further and will result in higher yields. In one study peanut yield were 19% higher after 2 years of corn and 41% higher after 2 years of bahiagrass. Other data from Georgia indicates that growers use shorter rotations under pivots due to the large capital investments required for irrigation systems and a one, two, and three year rotation under the pivot resulted in a 7, 36, and 34% yield increase as compared to 11, 25, and 28% for non irrigated as compared to continuous peanuts under both systems. Rotations produce more profit, and reduce the amount of management and capital inputs needed. Peanuts grow best on well-drained soils and in full sun. If planted on poorly-drained soils, diseases will be prevalent and yields are normally low. Peanuts can be grown on sandy or excessively-drained soils, but irrigation may be needed for consistent production.

Peanut Fertilization

Soil tests from soil samples taken immediately after harvest of cotton or peanut can be used to determine lime as well as fertility requirements for crops for the coming year. If soil pH needs adjusting, fall is a good time of the year to apply needed lime since it may take as long as 6 months for full reaction. However, some reaction does occur soon after application. Bacteria, that fix nitrogen in relation to the peanut plant, do better and form more nodules with an adequate calcium (Ca) level and with pH around 6.0 or higher. If the crop that precedes

peanuts is well-fertilized, there may be enough residual nutrients in the soil to make direct fertilization unnecessary for the peanuts. The analysis and report of the soil sample will also tell you which nutrients are needed to grow peanuts on a particular soil. Calcium and magnesium (Mg) may be supplied by the limestone. If no lime is needed, Ca can be supplied by gypsum, and Mg can be included in the fertilizer. If phosphorus (P) is low, be sure to add sufficient P in the fertilizer. The same is also true for potassium (K) and other fertilizer that is needed by the peanuts. Many of the micronutrients are needed by peanuts. Boron (B) is normally the most deficient, especially on sandy soils. Therefore, if peanuts are to be planted on a sandy soil, it would be advisable to use B in the peanut fertilizer. The B should be applied at the rate of 0.5-0.75 pound of elemental B per acre. Higher rates of B can be toxic to the plants. Applications of B should be split since it is a highly leachable, mobile nutrient. High application rates of other nutrients can make B deficiency more pronounced. Deficiencies are most often found on highly weathered, sandy soils. The deficiency that we most often associate with B deficiency is internal fruit damage that we call "hollow heart", which reduces the quality and value of the crop. However, in more severe cases, B deficiency can result in split stems and roots, on the lower part of the stem with shortened internodes, terminal death, and extensive secondary branching. Leaves may be dark green and mottled with few or no peanuts developing on stubbed pegs. Some fields have been observed with these symptoms recently and this is a reminder that there is a reason to apply B even if little visual observations of deficiencies are seen. B may be applied early with herbicides or with fungicides to keep from making additional trips across the field. Split applications are desirable on sandy fields with a total of 1/2 to 3/4 pound of B per acre for the year. The crop may take up less than a tenth of a pound per acre but it is still important for crop production.

Peanuts may have a yellow cast to them during the growing season at times. There could be several causes for this including poor nodulation, micronutrient deficiencies, water logged soils, or herbicide damage. Some fields are yellow due to being water logged which results in poor nodulation and nitrogen fixation. Plants usually grow out of this

as soils dry out. Iron and other micronutrients may be limiting in water logged soils too, which will grow out of it as soils dry out. Manganese deficiencies may occur on soils that have been limed for years and have a pH above 6.3. These symptoms can be seen as a light green to yellow cast to the peanut canopy. Manganese applications will have to be made for the crop to overcome this. It is possible to lower the pH through acid forming fertilizers; however, applications of a few pounds of micronutrient may be more cost effective and the response will be quicker than changing the pH.

Calcium, Liming, pH, and Gypsum

Lime should be added if the soil pH is below 5.8, with the target pH being 6.2-7.0. If lime is needed, both a dolomitic or calcitic lime can be used and additional Ca as gypsum may be needed on larger seed varieties. In addition to reducing soil acidity, calcitic lime supplies the plant nutrient Ca while dolomitic lime supplies both Ca and Mg. Peanut responds very little to direct fertilization of most nutrients. However, Ca is needed in high levels by peanut for developing a viable seed. High levels of Ca are not necessary to grow a healthy plant. The amount of Ca taken up by the plant is dependent on the concentration in soil solution and on the amount of water moving into the plant. Calcium is the element most commonly deficient for peanut. Calcium deficiency results in high incidence of pod rot and unfilled pods called "pops". These peanuts also have much lower germination if saved for seed. Georgia research has shown that Ca applied as lime should not be turned under and that turned under lime had yields similar to no lime. Even though peanut has a lower Ca requirement than for soybean or cowpea, peanut does have an exceptional need for Ca for seed maturation and quality. Lime should be applied to fields well in advance of planting and may be applied to strip tilled fields as a surface application. For those growers who use minimum tillage and strip tillage, surface applications are acceptable. We have long term plots that have not been turned or had lime incorporated for 28 years that are still producing good crop yields. A high Ca and P layer can develop in the top 2-3 inches after many years of surface applications of fertilizer and lime. Calcium is routinely applied as gypsum at pegging time on sandy

soils for rapid replenishment of soil solution Ca. This is not as necessary on heavier soils that have higher diffusion gradients toward the pods. Since peanuts are often grown on sandy soils, which are drought prone, there is a limited ability of these soils to replenish the soil solution Ca. Heavier soils and irrigated soils are better able to supply the needed Ca for proper uptake. The Ca needs are primarily for pod and seed development and not for growing a healthy plant. Test soils and apply the needed amounts of Ca for good yields and quality.

The critical period for Ca absorption begins about 20 days after pegs start entering into the soil and may extend for an additional 60 days. However, some researchers have reported that 69% of total Ca uptake occurred between day 20 and 30 after pegging begins. It is then a necessity that proper amounts of Ca are supplied for the first 30 days after pegging begins. The problem occurs when limited soil moisture coincides with the high Ca need period and there is no moisture for Ca to be in solution for uptake by peanut. Sandy soils in the peanut region have low moisture retention capacity which leads to moisture induced Ca deficiency. Much of the irrigation installed in the SE was due to corn in rotation with peanuts. High levels of K and Mg in the soil can inhibit uptake of Ca and therefore peanuts are often not fertilized and "high cal" lime is used instead of dolomite. Soil test levels of about 450 lbs./A of Ca result in maximum yields of runner type peanuts while levels almost double this are necessary for maximum yield of Virginia type peanut as noted from research by S. Hodges at UGA. The larger peanuts have a smaller surface-to-weight ratio and requires a higher concentration of soil solution Ca in order to provide adequate Ca to the pod. Therefore, if soil samples show only Ca to be low, then calcitic lime would be satisfactory. However, if both Ca and Mg are low, then dolomitic limestone should be selected. Lime should be applied 3 to 6 months before planting peanuts, but can be planted up to time of planting. Lime can be applied in the fall to allow time to react with the soil. If Mg soil test is not low, use calcitic lime so that Mg will not compete with Ca for uptake by peanuts. Gypsum is often applied to peanuts at pegging time so that peanuts will have an adequate Ca supply. Rates of 250 lbs/A of dry gypsum in a band to 1000 lbs/A of wet gypsum are

often applied to peanuts that are to be saved for seed or when the soil test shows a need.

Inoculating Peanuts

Peanuts may not always respond to rhizobium inoculation. The main reason for this is that there is an indigenous population of rhizobium called cowpea miscellany that is common to many native weeds. These organisms are potentially able to nodulate a crop of peanuts grown for the first time in a field. Peanuts are only moderately efficient in fixing and translocating atmospheric nitrogen (N). With soybean, as much as 80% of the plant N comes from the atmosphere while about 55% of the plant N needs of peanut are from N fixation. Calcium is important to nodulation and maximum peanut root growth occurs at a pH of about 7.3 while shoot growth, nodulation, and N fixation is best at a pH range of 5.9 to 6.3. An application of lime can improve the availability of Ca, Mg, and P and decrease aluminum toxicity. Inoculants may be applied to soils that have not had peanuts grown on them for several years or at all. In some cases inoculation can increase nodulation and in others little response may be noted. However, inoculants are cheap insurance in providing needed N for plant growth. Nitrogen fertilizer will not normally increase yields unless the N-fixing bacteria that live in nodules on the peanut roots are not present. If needed, a commercial inoculant can be applied in the seed furrow at planting to provide the needed bacteria.

Planting Date and Depth

Peanuts were traditionally planted in April before TSWV. This disease has pushed the planting window later and the recommended date is now May 11-25. Peanuts can do well planted through the first week of June and will suffer yield loss planted later than this. Early planted peanuts (any time in April) could suffer a serious yield loss from TSWV in many areas of Florida. Highest numbers of thrips (*F. fusca*) that vector the virus occurred on April planted peanuts, whereas peanuts planted in May had smaller populations. The population dynamics of thrips in non-crop plants or volunteer peanuts early in the season is a reason for these effects since these plants may serve as reservoirs for TSWV.

Modeling work with the PNUTGRO model in Gainesville indicated that the highest yield could be expected from an April 15 planting while planting April 1 would result in about a 6-8% yield decrease and May 1 about 1-2 % yield decrease. However, the Gainesville area has not had the spotted wilt pressure that most of the other areas of the Southeast have experienced and it is located further to the south than much of the peanut growing area. The Southeast Climate Consortium has a peanut model for Florida. This model shows potential yields according to their climate predictions. The website can be found at (<http://www.agclimate.org>). If conditions are to be dry, cover can be killed early for those who strip till or land can be bedded early to have enough moisture to plant on time. Peanuts may be planted deeper than any of the other row crops. It will emerge from 2-3 inches deep or deeper. However, like other seed that are high in oil content, seed should be planted into moisture for rapid germination.

When planting peanuts, be sure to handle them gently and carefully. If they are handled roughly or the bags are dropped, seeds will split and germination will be reduced and a poor stand results. Peanut seed should be planted in rows that are 24 to 36 inches apart. The distance between rows will be determined by the variety being planted and the equipment available. However, many growers plant in "twin rows," which are two rows about seven inches apart, with the middles between the sets of twin rows being about 24 to 36 inches in width. The seed should normally be placed 2 to 4 inches apart in the rows depending on the row pattern and type of peanut. The seed should be placed 1 1/2 to 3 inches deep in light-textured soils and 1 1/2 to 2 inches deep in heavier-textured soils. After planting, the top of the seed rows should be level with or slightly above the middles. Many farmers prefer to plant their peanuts on a ridge in order to provide good water drainage away from the peanuts. If the soil is well drained, it would not be necessary to plant on a ridge. Irrigation may be made ahead of planting in order to make the planting operation easier. If it is dry after planting, be sure to irrigate to get the peanuts up to a good stand. Also, if certain herbicides are applied at or soon after planting and no rain or irrigation occurs, poor results may be obtained. Therefore, an irrigation to move the

pesticides into the top 2 or 3 inches of soil would be beneficial.

Conservation Tillage Methods for Peanuts

Strip tillage has been widely adopted by many commercial peanut producers. With this method of tillage, the land is not turned but is subsoiled, and only a strip of soil in the row is tilled (8-12" wide), while the soil between the rows remains undisturbed. This method of tillage reduces time and costs for land preparation and also helps prevent erosion of the soil by wind and water. Yields of crops planted in conservation tillage systems are often different from conventional plantings. They have been shown to be much higher, much lower and in many cases the same. The residue on the surface lowers disease incidence of leaf spot, TSWV, white mold and other diseases. In conventional tillage systems, the plant residue is incorporated into the soil resulting in more disease problems. Questions arise about strip tilling into fields that have not had crops grown for a few years. There are usually many hard-to-control weed species present including broom sedge and horseweed as well as other broadleaf weeds. Soils in these fields are often mellow in the top few inches due to decomposition of plant residue and not being tilled or compacted by equipment for several years. Normally, if a grower can plant through various plant material there will be no problem with digging. All weeds need to be controlled 4-5 weeks before planting and 2,4-D is often necessary for some of the hard to control weeds like horseweed and evening primrose. Glyphosate or other similar materials can be used to kill many other weeds a few weeks before planting. Peanut can also be planted into old cotton residue without having stalks in the picker when harvesting peanuts.

Yield variability can often be explained due to cover crops and the time the cover crops were killed and fungi associated with decomposition of the cover crops or previous crop residue. Many of the main diseases of cotton and peanut (*Pythium*, *Rhizoctonia*, *Phytophthora*, and *Fusarium*) need carbohydrates and proteins for infections to begin. Both energy and protein can be obtained from newly killed cover crops or plant residue. With cellulose and proteins

being the main food supply for pathogens, green or newly decomposing plant material can set up conditions for disease infections and resulting seedling diseases. Seedling diseases often lower crop yields when cover crops are not killed out soon enough since carbon and other nutrients are adequate for pathogen growth. With legumes, about 50% of the biomass may be lost during the first 4 weeks. Grass cover crops may take several weeks longer than legumes to decompose. Cover crops decompose more slowly when left on the soil surface as compared to incorporation. Generally, decomposition of cover crops is 2-3 times faster for each 10 degrees C increase in temperature. Therefore, cover crops decompose much more slowly under cool conditions as in 2005, when heat units were around 60% of normal by May 15. This resulted in slower decomposition of cover crops than normal leaving more food resources for plant pathogens to infect developing crops. Many root and stem diseases can influence plant growth without severe visual symptoms. This may be one reason that peanut yields were not as good for some people using conservation tillage in 2005. One way to overcome having nutrient sources for the pathogens in the spring is to kill out the cover crop earlier in the year so that the carbohydrates and proteins are expended by micro-organisms leaving inadequate resources for the pathogens. Preliminary research with bahiagrass also shows that peanut yields can be increased 1000-1500 lbs/A if bahiagrass is killed in the fall as compared to the spring prior to planting peanuts.

Pest Management and Monitoring

After planting, peanuts should be monitored continually for proper growth and pest problems. If stands are poor or if the plants are weak and deformed, it may be necessary to replant in order to get a good stand of healthy peanuts. However, determine the reason for the poor stands. If it is poor seed, you may have to find another seed source. If it is disease, you may have to improve the seed treatment. If it is caused by soil compaction, dryness of the soil, or other reasons, then the problems should be corrected before the peanuts are replanted.

Weed control is very important to making high yields. Peanut weed control information can be found

at <http://edis.ifas.ufl.edu/WG008>. Season-long weed control is necessary since weeds present at digging and harvest can disrupt these operations resulting in yield losses. Many growers have ceased to use tillage after planting even if tillage was done to prepare a seedbed. Proper use of herbicides may eliminate the need for cultivation. Researchers have reported that each cultivation can result in loss of 1 inch of soil moisture which can have a major impact on peanuts in a dry year without irrigation.

If peanuts are planted late, it may be necessary to start applying fungicides soon after the peanuts emerge. Foliar disease control information can be found at

<http://plantpath.ifas.ufl.edu/takextpub/ExtPubs/ppp1205.pdf>. If the peanuts have been planted early, fungicide application for leafspot control should begin about 30 days after planting, but no later than June 1. The leafspot disease is much worse in warm, wet weather than in cool, dry weather. Fungicides for leafspot control are one of the major costs in growing peanuts. If leafspot is controlled, higher yields are obtained and there are fewer problems from white mold, which can be caused by the defoliated leaves rotting and providing an inoculum for the white mold fungus. Rust and other pests of peanuts must be monitored closely. The leafspot disease can only be controlled by a spray applied every 10-14 days with this interval being shortened in case of frequent rainfall. Generally all seed are fungicide treated since the peanut shell is very fragile and can be broken easily. Fungicides help prevent diseases from killing the peanuts while still in the soil and before the plants emerge.

Also, monitor insect pests. Thrips are one of the major early-season pests of peanuts (<http://edis.ifas.ufl.edu/IN176>). Lesser corn stalk borers are one of the more difficult pests of peanuts to control. Lesser corn stalk borers can be damaging to peanuts from planting time until harvest. This is primarily a dry weather pest, so plenty of rain and/or irrigation should keep this pest population at low levels. Other insect pests of peanuts include those worms that eat leaves. By using a scouting program to determine which pests are present, only the pesticides that are needed can be applied and only when they are needed.

Nematodes should have been managed by good rotation and use of nematicides (<http://edis.ifas.ufl.edu/NG016>). Certain nematicides can be applied at pegging, but it should be kept in mind that nematodes cannot be effectively controlled in a crop after infestation. In other words, nematodes must be controlled on a preventative basis. Certified seed will have a germination test on the peanut seed that is bought for planting. Generally all seed are fungicide treated since the peanut skin is very fragile and can be broken easily. Fungicides help prevent diseases from killing the peanuts while still in the soil and before the plants emerge. Some nematicides and insecticides are applied before or at planting time. Fumigant nematicides are applied before planting while contact or systemic nematicides may be applied during the planting operation, with additional applications at pegging.

If a systemic insecticide is applied for thrips control, it is placed in the furrow at the time of planting. Herbicides may be applied before and/or at planting. Again, by knowing which pest problems to expect, you can decide which, if any, of these pesticides you want to use on your peanuts.

Irrigation

Peanuts grow well on sandy soils and are somewhat drought resistant at certain growth stages. There are critical stages of growth that require higher levels of soil moisture or yields can be severely limited along with quality. Table 3 below shows four growth stages and the relative response of the plants to moisture during each stage. The period of greatest water use occurs during pegging and is necessary to move Ca through soil solution to pegs and developing pods.

Peanuts are the least susceptible of any of the row crops in the Southeast to drought at planting time since they will emerge from 2-3 inches deep if necessary. However adequate moisture is necessary for good, uniform stands. Mid-season nut development is the most critical time for irrigation if there is a shortage of rainfall since it is the stage of maximum water use by the plant. In the southeast the critical part of the nut development/fruiting period includes the latter part of July, August and the first

week or so of September. Irrigation in the first two months after planting is usually not needed unless extremely dry conditions persist, because excess moisture can trigger excessive vine growth. Irrigation in late September may result in too much moisture during the peanut maturity stage and can increase the severity of CBR, Sclerotinia blight, and leaf-spot diseases. In dry years, irrigation can suppress the outbreak of spider mites and lesser corn stalk borers, make digging operations easier and reduce the threat of aflatoxin.

Determining Maturity

Determining optimum peanut maturity can be done in several ways but pod blasters that strip away the outer pod layer to expose colors indicative of the stage of pod maturity has become one of the primary ways. Blasters are widely used and are available at many county extension offices. Blasters save considerable time, as well as improve the accuracy over hand-scraping for determining the pod maturity profile. A high pressure washer and wire basket can serve as an inexpensive, quick, simple alternative to blasters that use glass beads, water, and compressed air. However, glass beads are still widely used. The key to the pressure washer effectiveness for blasting peanuts is a rotating turbo nozzle. A turbo nozzle takes a 0-degree jet stream which has the highest stripping power of any nozzle, rotates it, and spreads it out over a wide area. It provides superior stripping action compared to a flat fan nozzle at a pressure low enough not to damage the pods. An electric pressure washer providing 1.5 gpm at 1300 psi is quite adequate. In higher capacity, engine-driven models, the pressure should be reduced to approximately 1000 psi with the pressure regulator or by throttling down the engine. Place picked pods in a basket (pick all pods off plants to make about 200 pods from each location in the field that may have different maturities due to soil type, variety differences or other variables). Place the basket in a 5-gallon bucket to prevent splashing. The bucket should be equipped with a drain to prevent water build-up. Hold the pressure washer nozzle approximately 12 inches away and blast while vigorously shaking the basket. Watch the pods carefully. Stop in approximately 30 seconds and remove the immatures (yellows) before they disintegrate. Place the more advanced pods (orange

to black) back into the basket and blast until the entire outer pod layer has been removed. The entire process should be completed within three minutes or less. Blasted pods can be put on a chart to classify and rank the fields for time to harvest.

Peanut Harvest and Grades

One of the most critical parts of growing peanuts is timing of harvest to make maximum grade and yield. There are several things that a peanut grade tells the grower, these are: Two measurements are made from a sample taken directly out of the drying wagon: **FM** or foreign matter which denotes the percent or amount of plant material, rocks or soil that is in a load of peanut. Foreign matter is increased by digging with soils too wet or dry or peanuts vines not being properly cured. Peanut combines can be adjusted to some extent to blow more or less air to help reduce the trash along with having the vines in the proper condition when harvesting. **LSK** or loose shelled kernels is the percent of peanut which have been shelled during the harvest process which may be due to too high picker speed or too much air blowing peanuts against something sharp in the combine or the picker fingers being set too aggressively.

A sample is then taken and shelled and the following are determined on a percentage basis: **SMK** (sound mature kernels) is the percentage of those peanut that are shelled for grading and determined to be mature, while **SS** (sound splits) is the percentage that are good peanuts but are split. **OK** is the percentage of other kernels that may be immature or very small. **ELK** is extra large kernel percentage that ride over a certain screen size that manufacturers may use for special candy purposed. Certain peanuts have a tendency to have larger seed than others but growers seldom are paid for this. **DK** are kernels that are damaged due to disease, insects or other factors. Grades are based mostly on **TSMK** or the total sound mature kernels. Grades in the mid 70s are considered good and the grower probably did a good job timing the digging. If a peanut grade were 73% SMK and 2% SS for a 75 grade, the other 25% would be shells and other kernels. Some peanut varieties have thicker shells and will have lower grades than other varieties. Many growers may start digging too soon if they have large acreages and the

first few loads of peanuts will grade in the upper 60s or low 70s. Low grades can be caused by digging too early or having periods of drought when pegs did not set and there are “two” crops of peanuts, one set early and one set later when moisture returned. This makes it difficult to determine when to dig since you would have some very mature peanuts along with some very immature peanuts that were set later. Irrigated peanuts are usually easier to determine proper digging date since moisture can be supplied for a continuous set of peanuts. However, other factors like disease control and weather conditions can determine digging date and these often cause yield losses. Vines should be kept disease free as much as possible in case weather forces you to delay digging by a week or more and healthy vines will retain mature peanuts better than dying, diseased vines. A week early or late digging peanuts can make 500 lbs/A difference or more in yield and several points in grade. Five hundred pounds of peanuts are worth about \$90 and the difference between a grade of 69 and 75 on a 4000 lb/A peanut crop is worth about \$50. One key for high yields and profit is to keep vines healthy and dig on time.

Management Tips

- Since peanuts require a high level of management, many decisions are required in growing the crop. Fortunately, most of these decisions can be made prior to planting. For example, land selection, variety, and crop rotation are of vital importance in preventing many weed, insect, disease, and nematode problems. If all pest problems in a field are monitored for the crop grown each year, then the selection of pesticides is much easier and many problems can be averted.
- Whether or not to irrigate is a major decision because good yields cannot be maintained during dry years. However, for the small acreage producer, costs for irrigation equipment may make the practice uneconomical. If irrigation is utilized, a grower must be knowledgeable and judicious in use of supplemental water. White mold and pod rots are diseases that can be made more severe with poor irrigation management than the other problems would be with no irrigation.

- Gypsum can be used to supply calcium to peanuts. Calcium is needed by the peanuts to ensure well-filled pods, reduce pod rots caused by imbalances of other nutrients, and to improve viability of peanuts produced for seed. The need for gypsum can be determined by soil tests and, if needed, the gypsum should be applied no later than early flowering.
- Insecticide applications to the peanut foliage should be determined by scouting procedures that include insect counts of various species. Other insect pests, such as lesser corn stalk borers and cutworms, may attack below-ground peanut parts and thus require scouting techniques peculiar to peanuts.
- Fungicide applications are generally required for high yields of peanuts. Leafspot is usually the major disease of the crop. Information is available on selecting fungicide programs for peanuts. Normally, fungicide applications must be made on 10-14 day intervals, or on a weather-based procedure. Crop rotations and other factors will determine which fungicide should be selected.
- Some weed problems in the growing crop can be controlled by herbicide applications. Timing is of critical importance in these herbicide applications. In some instances, cultivation, hoeing, and/or hand pulling would be the preferred means of weed control. If cultivating, it is important not to throw dirt on the peanut vines because white mold problems can be made more severe if the vines are partially covered with soil.
- Because all pods do not mature at the same time, one of the most difficult problems in producing peanuts is determining when to harvest. Runner peanuts normally mature in approximately 135-155 days after planting. However, if Valencia peanuts are grown for boiling purposes, they may mature in as little as 75 days from planting in mid-summer. The same variety may take from 15 to 30 days longer to mature when planted in early or late season. Therefore, it is not possible to accurately predict at planting when a peanut crop will be ready to harvest. There are reliable and objective procedures that can be used to determine peanut maturity and the best time to harvest. For the commercial grower, the peanut maturity profile is preferred while others may pull vines occasionally to check pod maturity. There are many locations where pod blasters are available and charts to determine proper digging dates.
- After harvest, peanuts must be dried unless they are for consumption as boiled or green peanuts. Natural and artificial drying is normally used. It is important to use care in harvesting and drying peanuts as the fungus that causes aflatoxin development is favored by poor drying techniques.
- Storage of peanuts should be in an insect-free area. An insecticide is normally sprayed on the stored peanuts to prevent insect infestations. The moisture content of the peanuts should be below 10 percent for safe, long-term storage. The storage facility should be one that prevents moisture accumulation in or around the peanuts.

Table 2. Peanut varieties evaluated for disease in the southeast.

Variety ¹	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White mold	Limb rot
SunOleic 97R ²	50	unknown	unknown	unknown
Flavorrunner 458 ²	50	unknown	unknown	unknown
NC-V 11	35	30	25	25
NC12C	35	unknown	unknown	unknown
AT-201 ²	35	30	20	unknown
Georgia Green	30	20	20	15
Virugard	30	20	20	unknown
Gregory	30	30	20	25
VC2	30	unknown	unknown	unknown
Anorden ²	25	20	20	20
Andru II ²	25	30	20	25
C-99R ⁴	20	15	15	25
Hull ²	20	10	15	25
Carver ³	20	30	20	25
GA03L ⁵	15	15	10	20
GA02C ^{2,3,5}	15	20	10	20
GA01R ³	10	10	15	15
DP1 ⁴	10	5	10	25
AP3 ⁴	10	25	10	25
Tifrunner	10	15	25	25

¹Adequate research data is not available for all varieties with regards to all diseases. Additional varieties will be included as data to support the assignment of an index value are available. High numbers indicate more disease.

²High oleic variety.

³Varieties Carver, GA-02C, and GA-01R have increased resistance to *Cylindrocladium* black rot (CBR) than do other varieties commonly planted in Georgia.

⁴Varieties AP3, DP1, and C-99R are less resistant to CBR and are not recommended for fields where this disease is a problem.

⁵The malady referred to as “funky” or “irregular” leaf spot tends to be more severe in GA02C and GA03L than in other varieties. Although this condition can look like early leaf spot (*Cercospora arachidicola*), the cause “funky” leaf spot is unknown. Disease losses are not typically associated with funky leaf spot.

Table 3. Critical periods for soil moisture for peanuts.

Week of Growth	Growth Stage	Water Requirements
Week 1 to 2	Planting to emergence	Moderate
Weeks 3-8	Emergence to pegging/ flowering	Low
Weeks 8-15	Flowering/pegging and pod formation	High
Weeks 16-22	Pod formation to maturity	Moderate