

Saw Palmetto Control: Individual Plant and Broadcast Application¹

Brandon Fast, Jason Ferrell, and Brent Sellers²

Saw palmetto [*Serenoa repens* (Bartram) Small] is a shrubby palm species native to Florida and common throughout the state. Plants are typically 2 to 8 feet tall. They arise from subterranean or prostrate stems and may grow in clumps up to 18 feet in diameter. The palmate leaves of saw palmetto are 2 to 3 feet in diameter and are borne on sharply saw-toothed petioles (leaf stems) that are approximately 2 feet long (Figure 1). Flowering occurs between February and April on branches that are approximately 2 feet long, and flowers produce round fruits called “drupes” that are 0.5 inches in diameter and ripen from September to October. Refer to EDIS publication WEC109 (<http://edis.ifas.ufl.edu/uw110>) for further information concerning saw palmetto biology and ecology.



Figure 1. Palmate leaves on saw palmetto.

Saw palmetto leaves and drupes provide cover and food for over 211 species of wildlife. Drupes can also be consumed by humans, and they have reportedly been used for the treatment of 63 commonly occurring medical conditions. Despite its beneficial uses, saw palmetto is a serious weed problem in pastures, forests, and non-cropland areas throughout Florida (Figure 2). Saw palmetto competes with desirable native plant species in forests, prevents access to power poles and other utility structures in non-cropland areas, reduces forage yield and quality in pastures, and hinders ranching operations by reducing visibility and accessibility to livestock (Figure 3). For these reasons, control of this common native plant is often necessary.



Figure 2. Saw palmetto infesting native rangeland.

1. This document is SS-AGR-341, one of a series of the Agronomy Department, UF/IFAS Extension. Original publication date March 2011. Revised September 2012. Reviewed October 2015. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. Brandon Fast, former graduate student assistant; Jason Ferrell, professor, Agronomy Department; and Brent Sellers, associate professor, Agronomy Department, Range Cattle Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.



Figure 3. Saw palmetto can interfere with access to highway structures.

Mechanical Control

The massive underground stems and waxy leaves of saw palmetto make it an extremely difficult plant to control. Double roller-chopping (chopping twice in perpendicular directions) can effectively control saw palmetto; however, this procedure is extremely labor intensive, costly, and non-selective, which means that it causes damage to desirable plant species. Consider carefully before beginning a roller-chop program. For more information on roller-chopping and other mechanical treatments for saw palmetto, please see EDIS publication SS-AGR-105 (<http://edis.ifas.ufl.edu/ag173>).

Chemical Control

Control of saw palmetto can be accomplished by either spot-treating individual clumps or by broadcast spraying larger infestations. Spot-treatments can be made using either a backpack sprayer or a motorized sprayer mounted on a tractor or ATV. When making spot-treatments, treat the entire plant and achieve good spray coverage, but do not allow spray runoff.

For spot-treatment applications, using metsulfuron alone (the active ingredient in MSM 60, Cimarron, Ally, and others) will rarely be effective (Table 1). However, using triclopyr ester alone (triclopyr ester is the active ingredient in Remedy Ultra, Garlon 4, and many other products) is not a great deal better. With triclopyr ester, even at a 4% solution, you will observe rapid brown-out of all treated foliage and the plant will appear to be dead, but within a single year of the application, new sprouts will form at the base of the plant, and the plant will need to be retreated. With metsulfuron, total brown-out rarely occurs, but no new sprouts will emerge. Applying metsulfuron alone will

cause saw palmetto to slowly decline with yellow/green leaves remaining on the plant for the rest of the natural life of the plant. Triclopyr ester and metsulfuron mixed together usually provide more effective palmetto control. It seems that the triclopyr ester provides the leaf browning, while the metsulfuron helps to reduce resprouting. Regardless of whether you choose triclopyr ester, metsulfuron, or the combination, you will not likely achieve 100% control with one application. It is usually necessary to make a second herbicide application 8 to 12 months after the first.

When broadcasting herbicides on palmetto, it is important to increase the application volume to between 20 and 40 gallons of water per acre with either a crop oil or methylated seed oil adjuvant. Adequate spray coverage and proper adjuvant are essential for good control. For broadcast applications, results were similar to that of the individual plant treatments. Triclopyr ester plus metsulfuron combinations were required for maximum control. Replacing triclopyr with fluroxypyr (Vista XRT or Cleanwave) or aminopyralid (one of the components of Chaparral) was less effective. In broadcast applications as in individual plant treatments, none of the herbicide combinations tested provided 100% control. Follow the broadcast application with either an additional broadcast or spot-spray to control palmetto resprouts.

Metsulfuron is useful for broadcast application to pastures, with some precautions. Metsulfuron is highly effective on blackberry and many other weeds, but will also severely injure or kill Pensacola bahiagrass. If a sufficient population of bahiagrass is present in the palmetto-infested area, metsulfuron should only be used in spot-spraying or limited broadcast applications. Additionally, the maximum use rate for a metsulfuron product (with 60% active ingredient) in pastures is 1.66 oz of product per acre per year, or 0.06 lb active ingredient per acre. Use rates for forestry or industrial sites can exceed 1.66 oz/A/yr, but consult the product label before you apply.

For additional information on individual plant treatment and broadcast herbicide applications, see the following publications:

Herbicide Application Techniques for Woody Plant Control, EDIS Publication No. SS-AGR-260. (<http://edis.ifas.ufl.edu/ag245>)

Broadcast Boom Sprayer Calibration, EDIS Publication No. PI24. (<http://edis.ifas.ufl.edu/pi016>)

Table 1. Saw palmetto control 12 months after treatment with individual plant and broadcast treatments.

Treatment Type ¹	Herbicide ²	Rate	Saw palmetto control (%)
Individual plant	Metsulfuron 60DF	1 oz/100 gal	32
Individual plant	Metsulfuron 60DF	2 oz/100 gal	45
Individual plant	Metsulfuron 60DF	3 oz/10 gal	52
Individual plant	Triclopyr ester	2% solution	62
Individual plant	Triclopyr ester	4% solution	72
Individual plant	Metsulfuron 60DF + Triclopyr ester	0.5 oz/100 gal + 0.5 % v/v	78
Individual plant	Metsulfuron 60DF + Triclopyr ester	1.0 oz/100 gal + 0.5% v./v	78
Individual plant	Metsulfuron 60DF + Triclopyr ester	0.5 oz/100 gal + 1.0% v/v	87
Individual plant	Metsulfuron 60DF + Triclopyr ester	1.0 oz/100 gal + 1.0% v/v	88
Broadcast	Chaparral ³	3.3 oz/ac	9
Broadcast	Chaparral + Metsulfuron 60DF	3.3 oz + 0.5 oz/ac	6
Broadcast	Chaparral + Vista XRT ⁴	3.3 oz + 23 oz/ac	47
Broadcast	Chaparral + Triclopyr ester	3.3 oz + 1 qt/ac	88
Broadcast	Chaparral + Triclopyr ester	3.3 oz + 2 qt/ac	87
Broadcast	Triclopyr ester + Metsulfuron 60DF	2 qt + 2 oz/ac ⁵	91

¹ Individual plant treatments included 2% v/v methylated seed oil and broadcast treatments included 1% v/v crop oil concentrate.
² Always consult the manufacturer's label and follow all label directions when applying herbicides.
³ A dry-flowable herbicide containing aminopyralid (62%) + metsulfuron (9.5%).
⁴ Vista XRT at 23 oz/ac is equivalent to Cleanwave at 52 oz/ac.
⁵ Metsulfuron 60DF use rates for pastures cannot exceed 1.66 oz/A/yr, but higher rates can be used for forestry or industrial sites.