

# *Antirrhinum majus* Snapdragon<sup>1</sup>

Edward F. Gilman, Ryan W. Klein, and Gail Hansen<sup>2</sup>

## Introduction

A wide range of snapdragon selections is available. The tall types are 2 to 3 feet tall, the intermediates are 1 to 2 feet tall, the bedding types are 6 to 15 inches tall, and the rock garden hybrids are about 6 inches tall. The flowers come in a wide range of colors from reds, orange, yellow, and maroon. Plants with dark-colored flowers have dark green or reddish stems and those with white or pale flowers have pale green stems.



Figure 1. Flower—*Antirrhinum majus*: snapdragon.  
Credits: Edward F. Gilman, UF/IFAS



Figure 2. Full form—*Antirrhinum majus*: snapdragon.  
Credits: Edward F. Gilman, UF/IFAS

## General Information

**Scientific name:** *Antirrhinum majus*

**Pronunciation:** an-tur-RYE-num MAY-jus

**Common name(s):** snapdragon

**Family:** *Scrophulariaceae*

**Plant type:** herbaceous; annual

**USDA hardiness zones:** all zones (Figure 3)

**Planting months for zone 7:** April; September

**Planting months for zone 8:** February; March; October; November; December

**Planting months for zone 9:** February; October; November; December

**Planting months for zone 10 and 11:** February; November; December

**Origin:** not native to North America

1. This document is FPS-44, one of a series of the Environmental Horticulture Department, UF/IFAS Extension. Original publication date October 1999. Revised August 2018. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.

2. Edward F. Gilman, professor, Environmental Horticulture Department; Ryan W. Klein, graduate assistant, Environmental Horticulture Department; and Gail Hansen, associate professor, Environmental Horticulture Department; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

**Invasive potential:** not known to be invasive

**Uses:** mass planting; container or above-ground planter; cut flowers; edging

**Availability:** generally available in many areas within its hardiness range




Figure 3. Shaded area represents potential planting range.

## Description

**Height:** .5 to 3 feet

**Spread:** 1 to 2 feet

**Plant habit:** upright

**Plant density:** moderate

**Growth rate:** fast

**Texture:** medium

## Foliage

**Leaf arrangement:** alternate

**Leaf type:** simple

**Leaf margin:** entire

**Leaf shape:** oblong; spatulate

**Leaf venation:** none, or difficult to see

**Leaf type and persistence:** not applicable

**Leaf blade length:** 2 to 4 inches

**Leaf color:** green

**Fall color:** not applicable

**Fall characteristic:** not applicable

## Flower

**Flower color:** yellow; white; pink; orange; salmon; lavender; purple

**Flower characteristic:** showy

## Fruit

**Fruit shape:** no fruit

**Fruit length:** no fruit

**Fruit cover:** no fruit

**Fruit color:** not applicable

**Fruit characteristic:** inconspicuous and not showy

## Trunk and Branches

**Trunk/bark/branches:** not applicable

**Current year stem/twig color:** green

**Current year stem/twig thickness:** thick

## Culture

**Light requirement:** plant grows in part shade/part sun

**Soil tolerances:** acidic; slightly alkaline; clay; sand; loam

**Soil salt tolerance:** unknown

**Plant spacing:** 6 to 12 inches

## Other

**Roots:** not applicable

**Winter interest:** not applicable

**Outstanding plant:** not particularly outstanding

**Pest resistance:** long-term health usually not affected by pests

## Use and Management

Snapdragons grow in any slightly acidic, garden soil; however, they do not grow well in unamended clay. The plants require full sun and moist soil. A second crop of flowers may be obtained from plants that have finished flowering. Cut them back to within 5 or 6 nodes of the ground when the first flowers fade. Fertilize when the second crop of flower buds become visible.

Snapdragons may be propagated by seeds, or by cuttings which root readily. The seed germinates in 10 to 14 days at 70°F. Do not cover the seed with soil. Prechilled seeds germinate best. Seedlings with two to three sets of leaves are pinched, however, dwarf forms do not need pinching. Set plants in the ground after the danger of frost has passed. Plant in the fall for winter color in USDA hardiness zones 9 to 11. Plants sometimes survive and flower throughout the winter in zone 8b and south. Set the plants six to ten inches apart.

Dwarf cultivars include: 'Floral Carpet', 'Floral Showers', 'Kolibri', 'Royal Carpet', and 'Tahiti'. Intermediate cultivars include 'Princess', 'Liberty', 'Sonnet', 'Pixie', 'Sprite', and 'Cinderella'. Tall cultivars include 'Panorama', 'Burpee's Topper', 'Spring Giant', and 'Rocket'.

## Design Considerations

The warm-colored flowers of the snapdragon are a striking feature that can be used to bring color to a plant bed. Warm

colors show best in full sun when paired with other plants with white flowers to make the red, orange, and maroon look more intense. Plants with dark green glossy leaves would also contrast well with the leaves of the snapdragon. Companion plants could include bold, large leaf textures in a tight clumping form or large-leaved small shrubs.

## **Pests and Diseases**

Aphids feed on terminal growth and the underside of the leaves. The insects suck juices, and heavy infestations seriously weaken the plants. The greenhouse leaf tier chews irregular-shaped areas in leaves and webs the leaves together. Pesticides are seldom effective after the insect rolls the leaves. Mites cause a bronzed or stippled appearance on the foliage, especially in hot weather.

Rust causes brown pustules surrounded by yellowed tissue on the leaves. Plants may bloom prematurely, have small flowers, and die early. Use proper plant spacings and resistant varieties.

Anthraxnose attacks the leaves and stems in late summer. On older stems the spots are sunken, oblong, yellowish-green to gray with a narrow brown border. On the leaves, the spots are yellowish-green turning dirty white with a narrow brown border. When the stem is girdled the plant dies. Destroy infected plants and use wider spacings.

Gray mold causes flower spikes to wilt, and light brown areas form on the lower stem of the flower cluster. Infected plants break over below the flowers. The disease is worse in wet weather. Cut off infected flower stalks and keep beds free of debris.

Stem rot can be detected by the presence of cottony growth on stems of infected plants near the soil line. Infected plants die and should be destroyed.