

Lagerstroemia x 'Tuscarora': 'Tuscarora' Crapemyrtle¹

Edward F. Gilman and Dennis G. Watson²

Introduction

A long period of striking flower color, attractive fall foliage, fabulous, exfoliating bark and good drought-tolerance all combine to make this Crape-Myrtle a favorite small tree for either formal or informal landscapes. But this cultivar appears to be less cold hardy than some of the other mildew-resistant selections. It is recommended for planting in urban areas.

General Information

Scientific name: *Lagerstroemia* x

Pronunciation: lay-ger-STREE-mee-uh

Common name(s): 'Tuscarora' Crapemyrtle

Family: *Lythraceae*

USDA hardiness zones: 7A through 9A (Fig. 2)

Origin: not native to North America

Invasive potential: little invasive potential

Uses: tree lawn 3-4 feet wide; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; street without sidewalk; parking lot island < 100 sq ft; parking lot island 100-200 sq ft; parking lot island > 200 sq ft; container or planter; trained as a standard; deck or patio; specimen; urban tolerant; highway median

Availability: not native to North America

Description

Height: 18 to 25 feet

Spread: 15 to 18 feet

Crown uniformity: symmetrical

Crown shape: vase




Figure 1. Young *Lagerstroemia* x 'Tuscarora': 'Tuscarora' Crapemyrtle

Crown density: moderate

Growth rate: moderate

Texture: medium

Foliage

Leaf arrangement: opposite/subopposite (Fig. 3)

Leaf type: simple

Leaf margin: entire

Leaf shape: oblong, elliptic (oval), obovate

1. This document is ENH-508, one of a series of the Environmental Horticulture, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date November 1993. Revised December 2006. Reviewed May 2011. Visit the EDIS website at <http://edis.ifas.ufl.edu>.

2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, former associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.



Figure 2. Range

Leaf venation: pinnate
Leaf type and persistence: deciduous
Leaf blade length: less than 2 inches, 2 to 4 inches
Leaf color: green
Fall color: orange, red, yellow
Fall characteristic: showy

Flower

Flower color: red
Flower characteristics: very showy

Fruit

Fruit shape: oval, round
Fruit length: less than .5 inch
Fruit covering: dry or hard
Fruit color: brown
Fruit characteristics: does not attract wildlife; showy; fruit/leaves not a litter problem

Trunk and Branches

Trunk/bark/branches: branches droop; showy; typically multi-trunked; thorns
Pruning requirement: little required
Breakage: resistant
Current year twig color: brown, green
Current year twig thickness: thin
Wood specific gravity: unknown

Culture

Light requirement: full sun
Soil tolerances: sand; loam; clay; acidic; alkaline; well-drained
Drought tolerance: high
Aerosol salt tolerance: moderate

Other

Roots: not a problem
Winter interest: yes
Outstanding tree: yes
Ozone sensitivity: unknown

Verticillium wilt susceptibility: resistant
Pest resistance: resistant to pests/diseases



Figure 3. Foliage



Figure 4. Flower

Use and Management

The 6- to 12-inch-long clustered coral pink blooms appear on the tips of branches during the summer. The individual flowers are ruffled and crinkly as to appear made of crepe paper. The smooth, peeling bark and multi-branched, open habit of Crape-Myrtle make it ideal for specimen planting where its bright red to orange-colored fall leaves add further interest. The tree is upright-spreading, or vase-shaped, with branches spreading out as they ascend. The tree probably grows 16 to 20 feet tall with a 15 to 18-foot spread. Lower branches droop as they grow older, and they will need to be removed to show off the bark and interesting trunk form.

Pruning should be done in late winter or early in the spring before growth begins because it is easier to see which branches to prune. New growth can be pinched during the growing season to increase branchiness and flower number. Pruning methods vary from topping to cutting Crape-Myrtle nearly to the ground each spring to the removal of dead wood and old flower stalks only. Topping creates several long, thin branches from each cut which droop down under the weight of the flowers. This practice disfigures the nice trunk and branch structure. Lower branches are often thinned to show off the trunk form and color. You can remove the spent flower heads to encourage a second flush of flowers and to prevent formation of the brown fruits. Since cultivars are now available in a wide range of growth heights, severe pruning should not be necessary to control size. Severe pruning can stimulate basal sprouting which can become a constant nuisance, requiring regular removal. Some Crape-Myrtle trees sprout from the base of the trunk and roots even without severe heading.

Crape-Myrtle grows best in full sun with rich, moist soil but will tolerate less hospitable positions in the landscape just as well, once it becomes established. It grows well in limited soil spaces in urban areas such as along boulevards, in parking lots, and in small pavement cutouts if provided with some irrigation. They tolerate clay and alkaline soil well. However, the flowers of some selections may stain car paint. Insect pests are few and 'Tuscarora' is resistant to powdery mildew. There are other new cultivars (many developed by the USDA) available which are resistant to powdery mildew.

Many cultivars of Crape-Myrtle are available: hybrid 'Acoma', 14 to 16 feet tall, white flowers, purple-red fall foliage, mildew resistant; hybrid 'Biloxi', 25 feet tall, pale pink blooms, orange-red fall foliage, hardy and mildew resistant; 'Cherokee', 10 to 12 feet, bright red flowers; 'Powhatan', 14

to 20 feet, clear yellow fall foliage, medium purple flowers. The hybrid cultivars 'Natchez', 30 feet tall, pure white flowers, and 'Muskogee', 24 feet tall, light lavender flowers, are hybrids between *Lagerstroemia indica* and *Lagerstroemia fauriei* and have greater resistance to mildew. The cultivar 'Crape Myrtlettes' have the same color range as the species but only grow to three to four feet high. The National Arboretum releases are generally superior because they have been selected for their disease resistance. These releases may prove more resistant to powdery mildew in the Deep South, although further testing needs to be done to confirm this.

Propagation is by cuttings or seed.

Pests

Aphids often infest the new growth causing an unsightly but harmless sooty mold to grow on the foliage. Heavy aphid infestations cause a heavy black sooty mold which detracts from the tree's appearance.

Diseases

Powdery mildew can severely affect Crape-Myrtle but 'Tuscarora' is resistant.