

Errata

On the first page of the Table of Contents in the September 2020 issue, 103 (3), the research paper by Houlihan et al. should show the page range as 369–375. We regret the error.

We have been informed that there was a misplacement of figures in the article beginning on page 619 in the September 2019 issue, 102 (3). We regret the error and have made the correction below.

Natural enemies of *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae), a recent invasive pest on maize in South India

Sharanabasappa, C. M. Kalleshwaraswamy, J. Poorani, M. S. Maruthi, H. B. Pavithra, and J. Diraviam

The correct captions are:

Fig. 1. Larval parasitoid of *Coccygidium melleum* on *Spodoptera frugiperda*.

Fig. 2. Larval parasitoid of *Odontepyris* sp. on *Spodoptera frugiperda*.

We have been informed that there was an incorrect figure in the article beginning on page 376 in the September 2020 issue, 103 (3). Figure 1 should be the image shown below. We regret the error.

Coloration patterns of the tegmina of *Mahanarva spectabilis* (Hemiptera: Cercopidae): biological, morphological and genetic bases

Ricardo Aquino Borges, Alexander Machado Auad, Marcy das Graças Fonseca, Cristiano Amancio Vieira Borges, Ana Luisa Sousa Azevedo, Tiago Teixeira de Resende, Gervásio Silva Carvalho, and Fernando Rodrigues de Oliveira




Fig. 1. Relative frequency of 484 adults collected at the Embrapa Dairy Cattle experimental field in the town of Coronel Pacheco, Minas Gerais, Brazil, from which 242 couples were formed, with their 1,484 offspring, with regard to wing color patterns, regardless of sex.