- Butler, G. D., Jr., and A. L. Wardecker. 1971. Temperature and the development of eggs and nymphs of Lygus hesperus. Ann. Ent. Soc. Amer. 64:144-5.
- Fye, R. E., R. Pantana, and W. C. McAda. 1969. Developmental periods for boll weevils reared at several constant and fluctuating temperatures. J. Econ. Ent. 62:1402-5.
- Stitt, L. L. 1940. Three species of the genus Lygus and their relation to alfalfa seed production in southern Arizona and California. USDA Tech. Bull. 741, 19 p.



BURSTING WORKERS: A NEW MEANS OF DEFENCE IN SOCIAL HYMENOPTERA—(Notice.) When disturbed mechanically, minor workers of Camponotus (Colobopsis) saundersi and of C. sp. near saundersi contract their gaster until it bursts at an intersegmental fold. The mandibular glands, which extend throughout the whole body, also burst releasing large quantities of a whitish yellow (C. saundersi) or bright yellow (C. sp. near saundersi) secretion. The secreted fluid is very sticky and attacking ants are unable to move when contaminated with it. The term "autothysis" (greek.: self sacrifice) is proposed for the phenomenon. Oecologia (Berl.), 1974, 14, 289-294; U. and E. Maschwitz, Univ. Frankfurt/M.