

CORRECTION TO NEW STATE RECORD AND NORTHERNEASTERN
RANGE EXTENSION FOR *CAENOCHOLAX FENYESI* SENSU LATO
(STREPSIPTERA: MYRMECOLACIDAE)

JEYARANEY KATHIRITHAMBY¹, STEVEN J. TAYLOR² AND RICHARD LAREAU³

¹Department of Zoology, South Parks Road, Oxford OX1 3PS, United Kingdom

²Illinois Natural History Survey, Institute of Natural Resource Sustainability, University of Illinois at Urbana-Champaign, 1816 South Oak Street (MC-652), Champaign, IL 61820-6953

³1393 Thayer Hall Dr., Mt. Pleasant, SC 29466-9478

The females of *Caenocholax fenyesi waloffi* Kathirithamby & Johnston were said in Kathirithamby et al. (2007) to stylopize *Hapithus agitator* Uhler (Orthoptera: Gryllidae: Eneopterinae) in Mexico. This is incorrect. Female *C. f. waloffi* parasitize *Macroanaxipha macilenta* (Saussure) (Orthoptera: Gryllidae: Trigonidiinae) (Kathirithamby & Johnston 2004).

REFERENCES CITED

- KATHIRITHAMBY, J., AND JOHNSTON, J. S. 2004. The discovery after 94 years of the elusive female of a myrmecolacid (Strepsiptera), and the cryptic species of *Caenocholax fenyesi* Pierce *sensu lato*. Proc. Royal Soc. London, B (3) 271: S5-S8.
- KATHIRITHAMBY, J., TAYLOR, S. J., AND LAREAU, R. 2007. New state record and northeastern range extension for *Caenocholax fenyesi* *sensu lato* (Strepsiptera: Myrmecolacidae). Florida Entomol. 90(4): 762.