

Community Structure and Intertidal Zonation of the Macroinfauna on the Atlantic Coast of Uruguay

Omar Defeo†, Eduardo Jaramillo‡ and Anibal Lyonnet*

† Instituto Nacional de Pesca
P.O. Box 1612
Montevideo, Uruguay

‡ Instituto de Zoología
Universidad Austral de Chile
P.O. Box 567
Valdivia, Chile

* Pereira de la Luz St.
Montevideo, Uruguay

ABSTRACT

DEFEO, O.; JARAMILLO, E., and LYONNET, A., 1992. Community structure and intertidal zonation of the macroinfauna on the Atlantic coast of Uruguay. *Journal of Coastal Research*, 8(4), 830-839. Fort Lauderdale (Florida), ISSN 0749-0208.

Five sandy beaches, covering a complete range of dissipative-reflective categories, were studied at the Atlantic coast of Uruguay. The objective of this study was to analyze the species richness, abundance and zonation of the intertidal macroinfauna in relation to beach textural characteristics. The most distinctive features of the macroinfauna studied were: (a) maximum species richness (17), densities and biomass occurred at a long dissipative beach (Barra del Chuy) of fine sands and a flat slope, while the lowest values of community parameters occurred at a truly reflective beach (Manantiales), with coarse sands and a steep slope; (b) polychaete worms were mostly collected from the long dissipative beach; (c) cirrolanid isopods showed a distinct pattern of distribution, with *Excirolana braziliensis* as the dominant cirrolanid in coarse sands and *Excirolana armata* in fine sands; and (d) faunistical zones were not easy to distinguish: they were recognized at Arachania and Punta del Diablo (two belts), and Barra del Chuy (three belts).

ADDITIONAL INDEX WORDS: Community structure, intertidal zonation, sand beach, Uruguay coast, macroinfauna.



INTRODUCTION

Substrate characteristics (*i.e.* grain size, sorting), which generally result from wave conditions and the geological history of an area, have been widely documented as one of the major factors influencing distribution and abundance of sandy beach macroinfauna (see review by MCLACHLAN, 1983). Generally, in a gradient of grain size and wave exposure, crustaceans dominate towards coarser and more exposed areas, while the proportion of polychaetes and bivalves increase towards the other side of that gradient (*e.g.* SEED and LOWRY, 1973; CROKER *et al.*, 1975; ELEFTheriou and NICHOLSON, 1975; MCLACHLAN *et al.*, 1981; DEXTER, 1984).

Until recently, very little was known about animal-substrate relationship and macroinfaunal community structure of sandy beaches of Uruguay; earlier studies were mainly descriptives like those of SCARABINO *et al.* (1974) and ESCOFET *et al.* (1979). Reflective and dissipative beaches (*sensu* SHORT, 1979, 1983; SHORT and WRIGHT, 1983) alternate along nearly 200 km of Atlantic coast providing a wide between-site variability in

textural characteristics, a situation which in turn suggests a macroinfaunal variability. Thus, the objective of this study was to analyze the zonation and community structure of the macroinfauna along this coast by comparing five beaches with different physical characteristics.

MATERIAL AND METHODS

Five sandy beaches on the Atlantic coast of Uruguay, at Manantiales, Solari, Arachania, Punta del Diablo and Barra del Chuy (Figure 1) were sampled during October 1982. The beaches were selected to represent a wide variability in grain size characteristics. Triplicate 0.11 m² sediments samples were collected at 5 m intervals on transects normal to the shoreline and spaced 3 m apart. Each transect was extended between the lower limit of the swash zone, determined by the minimum tide advance during the sampling time, and the beach face (defined by foredunes, cliff or drift lines). Depending on the width of each site, those beach levels varied between five and twelve stations. The sediment samples were sieved through a 1 mm screen and macroinfauna fixed in 5% formalin for later analyses. These analyses included species identification, counting and dry weight determinations.

91057 received 23 June 1991; accepted in revision 12 April 1992.

† Present address: CINVESTAV-Unidad Mérida. A.P. 73 Cordemex, 97310 Mérida, Yucatán, México.




Figure 1. Map of the Atlantic coast of Uruguay showing the location of the beaches analyzed in this study.

The beach face slope of each site was measured by the coefficient $a/l \times 100$, where a is the difference in height between the highest and lowest station (lower water line) and l is the distance between these two points. One sediment sample (20 cm deep) for grain size analyses was collected at each station. These samples were passed through a series of sieves of $-1.0, 0.0, 1.0, 2.0, 3.0$ and 4.0 phi ($\text{phi} = -\log_2 \text{mm}$); a mechanical shaker was used for 15 minutes on each sample. Graphic mean and standard deviation were determined by Folk's method (1980). Results were expressed in phi units and mm.

The rating system proposed by McLachlan (1980a) was used to categorize the five sandy beaches in relation to exposure. McLachlan's rating system of exposure takes into account observations of wave action, surf zone width, percentage of fine sand, median particle diameter, slope, depth of reduced layers and the presence of stable burrows. On a 20 point scale of exposure, beaches scoring 5 to 10 are considered sheltered, 11 to 15 exposed, and higher than 15 define very exposed beaches.

Density values per 0.11 m^2 were calculated and used to draw kite diagrams and describe zonation patterns. To analyze the zonation of species as well as abundance variability amongst sites, macrofaunal samples were subjected to classification and ordination procedures as follows:

(1) The taxonomic similarity between paired samples within and between beaches was calculated with the quantitative Bray-Curtis coefficient (Field *et al.*, 1982), and dendrograms were obtained after the Pair Group Method (Sokal and Sneath, 1973). In order to analyze clustering, the original abundances were root-root transformed (Field *et al.*, 1982; Raffaelli *et al.*, 1991).

(2) Ordination techniques were carried out by means of: (a) Principal Component Analysis (PCA), performed on substrate composition data (standardized correlation matrix), to search for patterns of substratum heterogeneity and to interpret macrofaunal variability in terms of grain size (Green, 1979). (b) Nonmetric Multidimensional Scaling (MDS), based on root-root transformed abundances, was applied in order to characterize the among-sites variability of the macrofauna (McLachlan, 1990; Raffaelli *et al.*, 1991).

RESULTS

The Beaches

The geographic location and several physical characteristics of the beaches studied are given in Table 1. Length and width showed a high between-site variability; Barra del Chuy being the longest and widest beach (22 km in length and 60 m in width). In general, the steepest beaches

Table 1. *Physical and biological characteristics of Uruguay sandy beaches.*

Beach Features	Manantiales	Solari	Arachania	Punta Diablo	Barra Chuy
Latitude	34°58'S	34°40'S	34°36'S	34°07'S	33°45'S
Longitude	54°48'W	54°12'W	54°06'S	53°44'W	53°27'W
Length (km)	2.50	12.00	3.50	2.50	22.00
Width (m)	20.00	30.00	35.00	25.00	60.00
Textural group*	coarse	fine	medium	fine	fine
Beach face slope (%)	11.00	5.16	7.56	3.00	3.17
Grain size** Φ	0.31	2.46	1.00	2.32	2.31
μm	807	182	500	200	202
Sorting Φ	0.73	0.46	1.11	0.63	0.59
μm	603	727	463	646	664
Exposure***	very exposed	exposed	very exposed	sheltered	exposed
Morphodyn. type	reflective	dissipative	intermediate		dissipative
Number of species	4	2	5	4	17
Density: Ind \cdot m ⁻² **	26 \pm 2	249 \pm 98	65 \pm 16	202 \pm 81	536 \pm 137
Dry biomass (g \cdot m ⁻²)**	0.18 \pm 0.03	1.16 \pm 0.48	11.59 \pm 6.26	4.00 \pm 1.77	20.32 \pm 8.24

* Type of sand after Folk (1980); ** overall mean based on the mean values calculated for the individual stations at each size; *** after McLACHLAN (1980a)

(Manantiales and Arachania) had the coarsest sediments and vice versa (Table 1). Grain size increased seaward, except at Barra del Chuy which presented the coarsest grains at both, the high and low beach levels (Figure 2). Sediments of Manantiales were represented by coarse sands (0.1–0.5 phi) all over the transect studied. The same holds true for the mid and low beach levels of Arachania (station 4–8), while those of the high beach levels (station 1–3) were made up of medium sands (1.2–1.7 phi). On the other hand, the sediments of Solari, Punta del Diablo and Barra del Chuy were fine sands (2.1–2.6 phi). Stations with the coarsest grains had the highest sorting values, and *vice versa* (Figure 2).

The rating system proposed by McLACHLAN (1980a) defined the beaches of Manantiales and Arachania as very exposed (score = 16); Solari (10) and Barra del Chuy (13) were defined as exposed, whereas Punta del Diablo, partly protected by rocky sharps, is relatively sheltered (9) from wave action (Table 1).

The Macrofauna

The highest values of overall mean density and biomass were found at Barra del Chuy beach, which also had the highest number of species. The lowest mean densities were observed from Manantiales and Arachania, being also the former the site with the lowest biomass (Table 1).

Mean grain size and slope values were significantly correlated ($r = -0.94$; $p < 0.02$) with macrofaunal densities (Figure 3). In both cases, a

monotonically decreasing exponential function of the form

$$\text{DENSITY} = ae^{-bx}$$

was successfully fitted, X being either slope or mean grain size.

Table 2 shows mean densities and abundances calculated for all species at each beach. The cirrolanid isopods *Excirolana braziliensis* and *Excirolana armata* numerically dominated all the sites, except Barra del Chuy. *Excirolana braziliensis* reached densities up to 44 ind m⁻² (Arachania), while the highest value for *Excirolana armata* was 264 ind m⁻² (Solari); both cirrolanids had a percentage representation higher than 60%. Barra del Chuy was numerically dominated by *Euzonus furciferus*, a polychaete with a mean density of 567 ind m⁻² and 44% of the whole macrofauna. In terms of dry biomass, cirrolanid isopods dominated at Manantiales (0.13 g m⁻²) and Solari (0.90 g m⁻²), while bivalves (*Donax hanleyanus* and *Mesodesma mactroides*) made up the bulk of the biomass at the other beaches. *Donax hanleyanus* dominated at Arachania (mean = 9.8 g m⁻²), while *Mesodesma mactroides* was the major contributor to biomass at Punta del Diablo (mean = 1.8 g m⁻²) and Barra del Chuy (mean = 8.6 g m⁻²).

The zonation of the species showed that high beach levels were mainly occupied by the talitrid amphipod *Orchestoidea brasiliensis* and the cirrolanid isopods *Excirolana braziliensis* and *Excirolana armata* (Figure 4). *Orchestoidea brasiliensis* and *Excirolana braziliensis* made up the




Figure 2. Mean grain size and sorting of the intertidal sediments sampled at the beaches studied.

main component of the mid beach levels at Manantiales and Arachania, the beaches with the coarsest grains. On the other hand, the mid beach levels of Solari and Punta del Diablo were numerically dominated by *Excirolana armata*. This species and the polychaete *Euzonus furciferus* were the dominant organisms at similar beach levels of Barra del Chuy. The low levels of this beach had the highest species richness of all the similar beach levels, and included the amphipods *Bathyporeiapus ruffoi*, *Stephensenia haematopus* and *Metarpinia* sp., the isopod *Macrochiri-*




Figure 3. Scatter diagrams and relationship between overall mean densities and (a) slope and (b) mean grain size (Mz).

dotea giambiagiae, the bivalves *Donax hanleyanus* and *Mesodesma mactroides*, the anomuran crab *Emerita brasiliensis*, the polychaetes *Spio gaucha* and *Sigalion cirriferum*, and the gastropods *Olivancillaria vesica auricularia* and *Buccinanops duartei* (Figure 4).

Multivariate Analysis

The first two components obtained from PCA based on percentage substrate size fractions for all sampling stations, accounted for 75% of the variation. The first component explained 59% of the variance and showed a dominant trend from fine to medium-coarse sands. Three main groups were delineated: (1) group G1, including stations with finer sands of Barra del Chuy, Solari and Punta del Diablo; (2) coarse grain stations of Manantiales and higher-mid levels of Arachania defined group G2; and (3) group G3, mainly represented by stations 6, 7 and 8 of Arachania (lower level), which presented the coarsest grain sizes.

The classification analyses discriminated by

Table 2. Species, taxonomic composition, dry biomass and density of the fauna on five Uruguay beaches.

Species (and taxon)		Manantiales		Solari		Arachania		Punta Diablo		Barra Chuy	
		B	D	B	D	B	D	B	D	B	D
<i>Orch. brasiliensis</i>	(A)	50	7			80	15			20	17
<i>Exciorolana armata</i>	(I)			901	264			1,636	216	580	169
<i>Exc. brasiliensis</i>	(I)	134	18			410	44	14	6	23	13
<i>Bathyporeiapus ruffoi</i>	(A)									14	41
<i>Phoxoceph. zimmeri</i>	(A)	0.3	3							22	100
<i>Macrochiridotea giam.</i>	(I)									6	11
<i>Euzonus furciferus</i>	(P)									1,548	567
<i>Donax hanleyanus</i>	(B)					9,780	24			3,052	32
<i>Hemipodus olivieri</i>	(P)					91	10			68	12
<i>Metarpinia</i> sp.	(I)	1	7							16	70
<i>Mesodesma mactroides</i>	(B)							1,768	6	8,589	34
<i>Stephensenia haemat.</i>	(A)									0.2	10
<i>Emerita brasiliensis</i>	(An)			262	87	1,129	19			1,508	22
<i>Spio gaucha</i>	(P)									1	20
<i>Buccinanops duartei</i>	(G)									4,433	40
<i>Olivancillaria v.a.</i>	(G)							586	7	411	17
<i>Sigalion cirriferum</i>	(P)									28	20

B = dry biomass (mg·m⁻²), D = density (ind·m⁻²). Under taxon A = Amphipoda, I = Isopoda, P = Polychaeta, B = Bivalvia, G = Gastropoda, An = Anomura

beach (Figure 5) showed that four of the five stations sampled at Manantiales (in station 5 no macrofauna was found) had a high value of taxonomic similarity (> 60%); i.e. no major groups were detected. At Solari, stations 1 to 6 formed a compact group (with *Exciorolana armata* as the single species), while the lowest intertidal level (station 7) remained apart. Stations sampled at Arachania were clearly classified in two groups; one with *Orchestoidea brasiliensis* and *Exciorolana brasiliensis* (stations 1 to 5), and another one with *Donax hanleyanus*, *Emerita brasiliensis* and *Hemipodus olivieri* as characteristic species (stations 6 to 8). At Punta del Diablo, two groups of stations were also identified: the co-occurrence of cirrolanid isopods *Exciorolana brasiliensis* and *Exciorolana armata* defined the first group (stations 1 to 3), while stations 4 to 6 showed a high value of taxonomic similarity (> 75%) due to the presence of the bivalve *Mesodesma mactroides* and its major predator, the gastropod *Olivancillaria vesica auricularia*. Finally, the dendrogram obtained for Barra del Chuy showed that the intertidal levels of this beach can be classified in three groups which are linked at very low values of taxonomic similarity with station 1 (Figure 5).

Examination of the cluster analysis by pooling the sampling stations for all beaches (similarity level greater than 40%), and ordination plot by means of MDS, suggested the same affinities and association of stations with similar physical char-

acteristics (e.g. grain size). Thus, only the MDS plot is shown (Figure 6). Four groups were recognized: (1) group G1, clustering the majority of stations of Barra del Chuy, except those of higher beach levels; (2) group G2, comprising all stations with finer sands and flatter slopes (i.e. Solari and Punta del Diablo) and the common occurrence of the cirrolanid isopod *Exciorolana armata*; (3) group G3, corresponding to Manantiales and high-mid levels of Arachania. These stations with coarse grain had in common the numeric dominance of the cirrolanid isopod *Exciorolana brasiliensis* and the amphipod *Orchestoidea brasiliensis*. Finally, group G4 clumped stations pertaining to lower beach levels of Arachania, (i.e., group G3 in the PCA), with the presence of the tidal migrators *Donax hanleyanus* and *Emerita brasiliensis*. Stations 2 and 3 of Barra del Chuy, with a mixed macroinfauna of coarser (e.g. *Orchestoidea brasiliensis*, *Exciorolana brasiliensis*) and finer (e.g. *Exciorolana armata*) sands, and station 7 of Solari, with the unique presence of *Emerita brasiliensis*, departed from groups G2 and G3 respectively.

DISCUSSION

Community Structure and Ecology of Uruguayan Beaches

The five sandy beaches analyzed in this study covered a wide range of beaches with differences in their morphodynamic type and exposure. Even though morphodynamic type was not accurately




Figure 4. Intertidal distribution of the macroinfauna at five Uruguay sandy beaches.

quantified by means of Dean's parameter (SHORT and WRIGHT, 1983; McLACHLAN, 1990), it is clear that coarser grains, a steep slope and a narrow surf zone width found at Manantiales chiefly de-

fining its reflective character, while the finer sands and flatter slopes demarcated the dissipative character of Solari and Barra del Chuy. Arachania, with intermediate values in grain sizes and slopes, may be categorized as an intermediate type of beach but closer to the reflective category.

The results of this study agree with others in which grain size and slope are assumed to be among the dominant factors controlling sandy beach community structure (e.g. COLMAN and SEGROVE, 1955; JONES, 1970; CROKER, 1977; DEXTER, 1979; McLACHLAN *et al.*, 1981; JARAMILLO, 1987). The most obvious difference between the macroinfauna was the marked contrast in abundance of the species collected in the reflective and intermediate beaches of Manantiales and Arachania, respectively as compared with the other beaches. The paucity of macroinfauna at those sites agrees with the findings of GAULD and BUCHANAN (1956), DYE *et al.* (1981) and McLACHLAN (1985), who reported that reflective beaches of Ghana, South Africa and Australia were characterized by an impoverished macroinfauna. As shown by several authors (e.g. McLACHLAN *et al.*, 1981; DEXTER, 1983), species richness and macroinfaunal abundance were negatively correlated with particle size. But grain size characteristics can not explain the low diversity of Solari, a dissipative beach with sands similar in size to that of Punta del Diablo and Barra del Chuy, which had a higher species richness. The sands of Solari differed from that of the other beaches by its higher water content, due to the fact that an effluent of Laguna de Rocha flows through this beach. Thus, the higher water content (and hence, a different sediment fabric) affects the macroinfaunal community of this beach.

Another important difference along the Atlantic beaches of Uruguay was related to the abundance of cirrolanid isopods. While *Excirolana brasiliensis* characterized the upper and middle intertidal of coarsest beaches (Manantiales and Arachania), *Excirolana armata* dominated the fine sands of Solari, Punta del Diablo and Barra del Chuy. This pattern suggests preferences or adaptive abilities to different environmental characteristics such as wave exposure and grain size (DE ALAVA, 1989; DEFEO *et al.*, 1989), as observed for some closely related sandy beach peracarids (JONES, 1970). The distributional pattern shown by *Excirolana* spp. on the beaches studied here, represents a common example of closely related species partitioning available space as a way of reducing or avoiding potential competition (JONES,




Figure 5. Cluster analysis of Uruguayan fauna discriminated by beach. Stations are numbered as in Figure 4.

1979; CROKER, 1967; CROKER and HATFIELD, 1980; DEFE0 *et al.*, 1989).

Abundance and diversity of polychaetes were other differences found along Uruguayan beaches. As reported by several authors (*e.g.* MCINTYRE, 1970; SEED and LOWEY, 1973; CROKER, 1977; DEXTER, 1983; MCLACHLAN, 1983), the abundance of sandy beach polychaetes increases along a gradient of decreasing wave exposure. In con-

trast, in our study most of the polychaetes were collected at the dissipative beach of Barra del Chuy, being the opheliid *Euzonus furciferus* the macroinfaunal species which reached the highest abundance, whereas no polychaetes were collected on Punta del Diablo, a relatively sheltered beach with similar beach slope and grain sizes to Barra del Chuy.

The above mentioned facts suggest that factors other than those related to wave energy could explain the observed differences in abundance and species richness among the dissipative beaches studied. Such differences may be related to the amount of food supply, usually provided by surf phytoplankton (MCLACHLAN, 1980b; MCLACHLAN and LEWIN, 1981; GIANUCA, 1983; LEWIN and SCHAEFER, 1983). High productivity and dense blooms of surf diatoms (particularly *Asterionella glacialis*) have been frequently registered at Barra del Chuy (BAYSSE *et al.*, 1989; DEFE0 and SCARABINO, 1990). This situation does not occur at Solari and Punta del Diablo, with lower wave energy and surf diatom productivity. Such differences in food availability may also explain the higher biomass of filter feeders such as *Mesodesma mactroides*, *Donax hanleyanus* and *Emerita brasiliensis* at Barra del Chuy, as reported for similar beach/surf zone ecosystems around the world (MCLACHLAN, 1983, 1990).




Figure 6. Ordination of stations using Multi-dimensional Scaling. Groups identified in the dendrogram are encircled here.

Zonation

Several zonation schemes have been proposed for sandy beach habitats. While DAHL (1952), PICHON (1967) and TREVALLION *et al.* (1970) based their schemes on biological factors (*i.e.* the intertidal distribution of the macrofauna), SALVAT (1964) used physical characteristics; *i.e.* the water content of the sediments (see detailed discussion of these schemes in McLACHLAN, 1990 and RAFFAELLI *et al.*, 1991). Recently, classification and ordination techniques have been used to examine the number of such faunistical zones: four zones have been identified in sandy beaches of South Africa (BALLY, 1983; WENDT and McLACHLAN, 1985), Namibia (DONN and COCKCROFT, 1989) and United States (McLACHLAN, 1990), while in northern Chile just three zones were determined (CLARKE and PEÑA, 1988). However, the recent paper of RAFFAELLI *et al.* (1991) highlighted the difficulties in establishing major biological zones, and suggested that the only zonation scheme universally applicable might be the one's of BROWN and McLACHLAN (1990), which recognizes a high-shore assemblage of air-breathers and a low zone of water-breathers.

The kite diagrams representing the intertidal distribution of the macrofauna and the multivariate analysis carried out, showed that a general pattern of zonation can not be presented for the Atlantic beaches of Uruguay; *i.e.*, the number of faunistical belts varied between one (Manantiales) and three (Barra del Chuy). However, it was possible to distinguish species which characterize different tidal levels. Thus, *Orchestoidea brasiliensis* and *Excirolana brasiliensis* can be considered as typical species of the subterrestrial fringe of DAHL (1952) or drying zone of SALVAT (1964). Both peracarids also characterize the high beach levels of the midlittoral zone (retention zone, *sensu* SALVAT, 1964) in which *Excirolana armata* and *Euzonus furciferus* also occurred. Mid and low levels of the midlittoral zone may be characterized by cirrolanid isopods and polychaetes, while the sublittoral fringe or saturation zone is chiefly characterized by *Emerita brasiliensis*, *Mesodesma mactroides* and *Donax hanleyanus*. It should be emphasized, however, that seasonal migration of some species such as *Mesodesma mactroides* (DEFEO *et al.*, 1986) and *Excirolana armata* (DE ALAVA and DEFEO, 1991) might obscure the observed zonation schemes.

In summary, three main conclusions can be drawn:

(1) The sandy beach macrofauna of the Atlantic coast of Uruguay increased in number of species and abundances from reflective to dissipative beaches. A quantitative relationship between biological (*i.e.* total densities) and physical (slope, grain size) variables was observed: total abundance of macrofauna was negative and exponentially correlated with grain size and slope. Multivariate analysis also showed the significance of particle size in structuring sandy beach communities.

(2) It was possible to recognize typical organisms that inhabit beaches with different grain size, such as *Excirolana armata* in finer sands and flatter slopes, and *Excirolana brasiliensis* in coarser sands and steep slopes.

(3) Even though macrofauna increased from higher to lower beach levels and, in some beaches, discrete assemblages coincided with Salvat's and Dahl's zones, zonation patterns were not easily distinguished.

ACKNOWLEDGEMENTS

We particularly thank Drs. A. McLachlan, A. de Ruyck and L. Capurro for critical reading and valuable suggestions to the final manuscript. We also thank Dr. J.C. Seijo for useful comments. The first author is grateful for the financial support granted by UNESCO, through fellowships (Grants 241.669.7 and 267.047.9) at Chile and CINVESTAV-IPN Unidad Mérida, Mexico.

LITERATURE CITED

- BALLY, R., 1983. Intertidal zonation on sandy beaches of the west coast of South Africa. *Cahiers de Biologie Marine*, 24, 85-103.
- BAYSSE, C.; ELGUE, J.C., and BURONE, F., 1989. Variaciones en la distribución y relaciones interespecíficas del fitoplancton en una playa arenosa de la costa atlántica uruguaya. *Frente Marítimo*, 5, 95-114.
- BROWN, A.C. and McLACHLAN, A., 1990. *Ecology of Sandy Shores*. Amsterdam: Elsevier.
- CLARKE, M. and PEÑA, R., 1988. Zonación de la macrofauna en una playa arenosa del norte de Chile. *Estudios Oceanológicos (Chile)* 7, 17-31.
- COLMAN, J.S. and SEGROVE, F., 1955. The fauna living in Stoupe Beck Sand, Robin Hood's Bay (Yorkshire, North Riding). *Journal of Animal Ecology*, 24, 426-444.
- CROKER, R.A., 1967. Niche diversity in five sympatric species of intertidal amphipods (Crustacea: Hausto-riidae). *Ecological Monographs*, 49, 746-751.
- CROKER, R.A., 1977. Macrofauna of northern New England marine sand: Long-term intertidal community structure. In: COULI, B.C. (ed.), *Ecology of Marine Benthos*. Belle W. Baruch Library in Marine Science

- #6, Columbia, SC: University of South Carolina Press, pp. 439-450.
- CROKER, R.A.; HAGER, R.P., and SCOTT, K.J., 1975. Macrofauna of northern New England marine sand. II. Amphipod-dominated intertidal communities. *Canadian Journal of Zoology*, 53, 42-51.
- CROKER, R.A. and HATFIELD, E.B., 1980. Space partitioning and interactions in an intertidal sand-burrowing amphipod guild. *Marine Biology*, 61, 79-88.
- DAHL, E., 1952. Some aspects of the ecology and zonation of the fauna on sandy beaches. *Oikos*, 4, 1-27.
- DE ALAVA, A., 1989. Dinámica poblacional y estructura espacio-temporal de *Excirolana armata* y *Excirolana braziliensis* (Isopoda: Cirolanidae) en playas del litoral atlántico uruguayo. B.S. Thesis, Universidad de la República, Montevideo, Uruguay.
- DE ALAVA, A. and DEFEO, O., 1991. Distributional pattern and population dynamics of *Excirolana armata* (Isopoda: Cirolanidae) in an uruguayan sandy beach *Estuarine, Coastal and Shelf Science*, 33, 333-344.
- DEFEO, O. and SCARABINO, V., 1990. Preliminary observations about the ecological significance of a possible-deposit feeding strategy in *Mesodesma mactroides* (Deshayes, 1854) (Mollusca: Pelecypoda). *Atlantica*, 12(1), 55-65.
- DEFEO, O.; LAYERLE, C., and MASELLO, A., 1986. Spatial and temporal structure of the yellow clam *Mesodesma mactroides* (Deshayes, 1854) in Uruguay. *Medio Ambiente* (Chile), 8, 48-57.
- DEFEO, O.; RIESTRA, G.; BRAZEIRO, A., and DE ALAVA, A., 1989. Ecología experimental en playas arenosas del Uruguay: Selección del sustrato e interacciones entre *Excirolana armata* y *Excirolana braziliensis* (Isopoda: Cirolanidae). I Congreso Latinoamericano de Ecología. Abstract.
- DEXTER, D.M., 1979. Community structure and seasonal variation in intertidal Panamanian sandy beaches. *Estuarine and Coastal Marine Science*, 9, 543-558.
- DEXTER, D.M., 1983. Community structure of intertidal sandy beaches in New South Wales, Australia. In: McLachlan, A. and Erasmus, T. (eds.), *Sandy Beaches as Ecosystems*. The Hague: W. Junk, pp. 461-472.
- DEXTER, D.M., 1984. Temporal and spatial variability in the community structure of the fauna of four sandy beaches in South-eastern New South Wales. *Australian Journal of Marine and Freshwater Research*, 35, 663-672.
- DONN, T.E. and COCKCROFT, A.C., 1989. Macrofaunal community structure and zonation of two sandy beaches on the central Namib coast, South West Africa/Namibia. *Madoqua*, 16, 129-135.
- DYE, A.H.; McLachlan, A., and Woolridge, T., 1981. The ecology of sandy beach in Natal, South Africa. *South African Journal of Zoology*, 16, 200-209.
- ELFETHERIOU, A. and NICHOLSON, M.D., 1975. The effects of exposure on beach fauna. *Cahiers de Biologie Marine*, 16, 695-710.
- ESCOFET, A.; GIANUCA, N.; MAYTIA, S., and SCARABINO, V., 1979. Playas arenosas del Atlántico Sudoccidental entre los 29° y 43° Lat. Sur: Consideraciones generales y esquema biocenológico. In: *Memorias del seminario sobre ecología bentónica y sedimentación de la plataforma continental del Atlántico Sur*. UNESCO Seminar, Montevideo, 9-12 May, 1978: 245-258.
- FIELD, J.G.; K.R. CLARKE, and WARWICK, R.M., 1982. A practical strategy for analysing multispecies distribution patterns. *Marine Ecology Progress Series*, 8, 37-52.
- FOLK, R., 1980. *Petrology of Sedimentary Rocks*. Austin TX: Hemphill, 182p.
- GAULD, D.T. and BUCHANAN, J.B., 1956. The fauna of sandy beaches in the Gold Coast. *Oikos*, 7, 293-301.
- GIANUCA, N.M., 1983. A preliminary account of the ecology of sandy beaches in southern Brazil. In: McLachlan, A. and Erasmus, T. (eds.), *Sandy Beaches as Ecosystems*. The Hague: W. Junk, pp. 413-420.
- GREEN, R.H., 1979. *Sampling Design and Statistical Methods for Environmental Biologists*. New York: Wiley.
- JARAMILLO, E., 1987. Community ecology of Chilean sandy beaches. Dissertation, University of New Hampshire, Durham, U.S.A.
- JONES, D.A., 1970. Factors affecting the distribution of the intertidal isopods *Eurydice pulchra* Leach and *Eurydice affinis* Hansen in Britain. *Journal of Animal Ecology*, 39, 455-472.
- JONES, D.A., 1979. The ecology of sandy beaches in Penang, Malaysia, with special reference to *Excirolana orientalis* (Dana). *Estuarine and Coastal Marine Science*, 9, 677-682.
- LEWIN, T. and SCHAEFER, C.T., 1983. The role of phytoplankton in surf ecosystems. In: McLachlan, A. and Erasmus T. (eds.), *Sandy Beaches as Ecosystems*. The Hague: W. Junk, pp. 381-390.
- McINTYRE, A.D., 1970. The range of biomass in intertidal sand, with special reference to the bivalve *Tellina tenuis*. *Journal of the Marine Biological Association of the United Kingdom*, 50, 561-575.
- McLACHLAN, A., 1980a. The definition of sandy beaches in relation to exposure: A simple rating system. *South African Journal of Science*, 76, 137-138.
- McLACHLAN, A., 1980b. Exposed sandy beaches as semi-closed ecosystems. *Marine Environmental Research*, 4, 59-63.
- McLACHLAN, A., 1983. Sandy beach ecology. A review. In: McLachlan, A. and Erasmus, T. (eds.), *Sandy Beaches Ecosystems*. The Hague: W. Junk, pp. 321-380.
- McLACHLAN, A., 1985. The biomass of macro- and interstitial fauna on clean and wrack-covered beaches in Western Australia. *Estuarine, Coastal and Shelf Science*, 21, 587-599.
- McLACHLAN, A., 1990. Dissipative beaches and macrofauna communities on exposed intertidal sands. *Journal of Coastal Research*, 6, 57-71.
- McLACHLAN, A. and LEWIN, J., 1981. Observations on surf phytoplankton blooms along the coast of South Africa. *Botanica Marina*, 24, 553-557.
- McLACHLAN, A.; WOOLDRIDGE, T., and DYE, A., 1981. The ecology of sandy beaches in Southern Africa. *South African Journal of Zoology*, 16, 219-232.
- PICHON, M., 1967. Contribution a l'étude des peuplements de la zone intertidale sur sables fins et sables vaseux non fixés dans la région de Tular. *Recueil Travaux Station Marine Endoume*, Supplement, 7, 57-100.
- RAFFAELLI, D.; KARAKASSIS, I., and GALLOWAY, A., 1991. Zonation schemes on sandy shores: A multivariate

- approach. *Journal of Experimental Marine Biology and Ecology*, 148, 241–253.
- SALVAT, B., 1964. Les conditions hydrodynamiques interstitielles de sédiments meubles intertidaux et la repartition verticale de la faune endogée. *Cahiers de Recherche de la Academie de Science de Paris*, 259, 1576–1579.
- SCARABINO, V.; MAYTIA, S., and FAEDO, J.C., 1974. Zonación biocenológica de playas arenosas del Departamento de Rocha (Uruguay) con especial referencia a la presencia de *Ocypode quadrata Fabricius*, 1967; (Decapoda: Brachyura). *Boletín Comisión Nacional de Oceanografía* (Uruguay), 1, 42–52.
- SEED, R. and LOWRY, B.J., 1973. The intertidal macrofauna of seven sandy beaches of County Down. *Proceedings of the Royal Irish Academy*, 73, 217–230.
- SHORT, A.D., 1979. Three dimensional beach stage model. *Journal of Geology*, 87, 553–571.
- SHORT, A.D., 1983. Sediments and structures in beach-nearshore environments, South East Australia. In: MCLACHLAN, A. and ERASMUS, T. (eds.), *Sandy Beaches as Ecosystems*. The Hague: W. Junk Publishers, pp. 145–155.
- SHORT, A.D. and WRIGHT, L.D., 1983. Physical variability of sandy beaches. In: MCLACHLAN, A. and ERASMUS, T. (eds.), *Sandy Beaches as Ecosystems*. The Hague: W. Junk Publishers, pp. 133–144.
- SOKAL, R.R. and SNEATH, P.H.A., 1973. *Numerical Taxonomy*. San Francisco, CA: W.F. Freeman.
- TREVALLION, A.; ANSELL, A.D.; SIVADAS, D., and NARAYANAN, B., 1970. A preliminary account of two sandy beaches in South West India. *Marine Biology*, 6, 268–279.
- WENDT, G.E. and MCLACHLAN, A., 1985. Zonation and biomass of the intertidal macrofauna of two sandy beaches in South West India. *Marine Biology*, 26, 1–14.

□ RÉSUMÉ □

On a étudié cinq plages de la côte atlantique uruguayenne qui se classent entre les types à dissipation et à réflexion. On a pu ainsi analyser en relation avec les caractères texturaux, la richesse des espèces, l'abondance et la zonation de la macroendofaune. Les principales caractéristiques de la macroendofaune sont: (a) l'abondance maximale en espèces (17), densités et biomasse se produit sur une longue plage de sable fin et plate, à dissipation (Barra del Chuy). Les plus faibles valeurs des paramètres se produisent sur une vraie plage à dissipation, au sable grossier et en pente raide (Manantiales). (b) Les vers polychètes se récoltent le plus fréquemment sur les plages à dissipation. (c) Les isopodes cirolanides ont une distribution bien distincte, *Excirrolana braziliensis* domine dans les sables grossiers, *Excirrolana armata* dans les sables fins. (d) La zonation faunistique n'est pas facile à faire. On a pu la reconnaître à Archania, à Punta del Diablo (deux ceintures), et à Barra del Chuy (trois ceintures).—Catherine Bousquet-Bressolier, Géomorphologie E.P.H.E., Montrouge, France.

□ RESUMEN □

En la costa Atlántica del Uruguay se han estudiado cinco playas arenosas, que cubre el rango completo desde playas disipativas a reflectivas. El objetivo de este estudio fue analizar la abundancia de las especies, la abundancia y la zonación de la macrofauna intermareal en relación a las características texturales de la playa. Las características principales de la macrofauna estudiada eran: (a) máxima abundancia de las especies (17), densidades y biomasa a lo largo de una playa disipativa (Barra del Chuy) compuesta de arena fina y poca pendiente, mientras que los valores más bajos de los parámetros de la comunidad se presentaron en una verdadera playa reflectiva (Manantiales), con arenas gruesas y una fuerte pendiente; (b) los poliquetos fueron recogidos sobre una larga playa disipativa; (c) los isopodos cirolanidos presentaban un esquema de distribución diferente, con *Excirrolana braziliensis* con cirolanidos dominantes en las arenas gruesas y *Excirrolana armata* en las arenas finas; (d) no fue sencillo distinguir las zonas faunísticas: ellas eran reconocidas en Arachania y Punta del Diablo (dos fajas), y en la Barra del Chuy (tres fajas).—Néstor W. Lanfredi, CIC-UNLP, La Plata, Argentina.

□ ZUSAMMENFASSUNG □

An der Atlantikküste von Uruguay wurden fünf Strände untersucht, welche das gesamte Energiespektrum von vollständig reflektierten Wellen bis vollständigem Wellenenergieverlust umfassen. Das Ziel dieser Studie war die Analyse von Artenvielfalt, Reichhaltigkeit und Zonierung der Makrofauna im Gezeitenbereich in ihrer Beziehung zum Aufbau und dem Gefüge des Strandes. Die untersuchte Makrofauna zeichnet sich durch folgende besondere Charakteristika aus: A) Maximales Artenreichtum (17), Dichte und Biomasse traten an einem langen Strand auf, der Wellenenergieverlust aufwies (Barra del Chuy). Der Strand setzte sich aus Feinsand zusammen und besaß eine geringe Neigung. Die niedrigsten Werte für sog. Tiergesellschaftsparameter fanden sich aber an einem rein reflektierenden Strand (Manantiales), welcher sich durch Grobsand und eine starke Neigung auszeichnete. B) Polychätenwürmer wurden hauptsächlich an dem langen Strand mit Wellenenergieverlust gesammelt. C) Die Vertreter der Isopodengattung *Cirrolana* zeigten ein bestimmtes Verbreitungsmuster, in dem *Excirrolana braziliensis* als die dominante Art im Grobsand vorherrschte und *Excirrolana armata* im Feinsand. D) Es war nicht so einfach, zwischen faunistischen Zonen zu unterscheiden, aber bei Arachania und Punta del Diablo waren es zwei und bei Barra del Chuy drei Gürtel zwischen denen unterschieden werden konnte.—Ulrich Radtke, Geographisches Institut, Universität Düsseldorf, Germany.