

Observations on the Distribution of *Meloidogyne* spp. in a Vineyard Soil

H. FERRIS¹

In an investigation on nematode distribution in a vineyard (1), difficulty arose in determining the *Meloidogyne* spp. present. Chromosomal analysis indicated the presence of at least six populations, representing four species (2). The objective of this study was to determine any differences among the *Meloidogyne* populations in their distribution in the soil and between vines.

Tomato seedlings (c.v. 'Pearson') were planted in seventy 500-cm³ soil samples [from different regions of the root zone of six vines (1)] and grown in a greenhouse for 6 weeks. Perineal patterns were cut from *Meloidogyne* females removed from the root systems of these plants. A minimum of six patterns was obtained from each plant. The patterns were of the *M. incognita* and *M. arenaria* types. None of the patterns resembled *M. hapla* or *M. javanica*, which were observed in the original study (2).

The percentage of *M. incognita* patterns was greater than that of *M. arenaria* (Fig. 1) when data were analyzed after arcsin transformation (3). The ratio of *M. incognita* to *M. arenaria* patterns throughout the vine root zone did not vary with depth (Fig. 1). The percentage of *M. incognita* in patterns taken from soil samples from each of the six vines did not differ (range 60-




FIG. 1. Percentage of *Meloidogyne incognita* and *M. arenaria* females in samples taken throughout the root zone of six grapevines. The height of each bar represents the percentage occurrence of a species at that position.

92%). At least half (50-67%) of the 500-cm³ soil samples from five of the vines had both *M. incognita* and *M. arenaria*. One-third of the samples from the sixth vine had both species.

There was no evidence of any differential distribution of the two *Meloidogyne* spp., either among vines or within the root zones of individual vines. Although this study did not determine whether the two species occupied the same galls or feeding sites, they frequently occurred in the same region of the vine root system. I am unable to explain the absence of *M. hapla* and *M. javanica* in these samples, unless their occurrence in the vineyard was more sporadic.

Received for publication 24 November 1975.

¹Assistant Nematologist, Department of Nematology, University of California, Riverside 92502. Supported in part by USDA-CSRS grant 316-15-63.

LITERATURE CITED

1. FERRIS, H., and M. V. MCKENRY. 1974. Seasonal fluctuations in the spatial distribution of nematode populations in a California vineyard. *J. Nematol.* 6:203-210.
2. HACKNEY, R. W. 1974. The use of chromosome number to identify *Meloidogyne* spp. on grapes. *J. Nematol.* 6:141-142 (Abstr.).
3. SNEDECOR, G. W., and W. G. COCHRAN. 1967. *Statistical Methods*. Iowa State University Press, Ames. 593 p.

BACK ISSUES OF JOURNAL OF NEMATOLOGY
AVAILABLE AT REDUCED PRICES

Single volumes 1 through 6 are available at \$6.00 per volume or \$30.00 for complete set of 6 volumes (plus \$2.00 for postage and handling). Please send checks, including bank charges, or money orders, and volume numbers desired to: Dr. J. M. Ferris, Department of Entomology, Purdue University, West Lafayette, Indiana 47907, USA.