

Molecular Characterization and Phylogenetic Evaluation of the Hsp90 Gene from Selected Nematodes

ANDREA M. SKANTAR¹ AND LYNN K. CARTA²

Abstract: While multiple genes are optimal for corroborating nematode phylogenies, only a few are commonly used. Here we examine the phylogenetic potential of the nuclear Hsp90 chaperone gene. We used degenerate primers to obtain partial Hsp90 sequences from several plant-parasitic and free-living nematodes. Hsp90 was single-copy in *Heterodera glycines* and *Meloidogyne javanica*, similar to the situation for *Caenorhabditis elegans*. The full-length *H. glycines* Hsp90 protein sequence showed homology to sequences from *C. elegans* and *Brugia pahangi* and to other eukaryotes, and contains several functionally important regions common to cytoplasmic Hsp90 proteins. The Hsp90 amino acid phylogeny supported the Coelomata hypothesis for metazoan evolution. Phylogenetic trees, substitution scatter plots, and statistics for phylogenetic signal were made for Hsp90, 18S small subunit (SSU), and 28S large subunit (LSU) over a limited but broad sampling of nematode taxa. Only the LSU data set failed to recover any of the expected topology and showed extensive substitution saturation. In an intensive sampling of plant-parasitic nematode taxa, the Hsp90 tree topologies were generally congruent with rDNA results and alignments were unambiguous. Hsp90 sequences may help strengthen branch support or clarify tree topologies when other molecules show ambiguous alignments, greater branch-length heterogeneity, or codon bias in certain taxonomic groups.

Key words: DNA, Hsp90, phylogenetics, protein, rDNA, saturation.

A number of molecules have been tested for their phylogenetic utility with varying degrees of success (Graybeal, 1994; Regier et al., 1998; Tang and Lewontin, 1999). The ribosomal genes are especially attractive for nematodes because their multi-copy nature facilitates amplification of the small quantity of DNA that may be available from single valuable specimens and they lack the introns that can sometimes complicate data analysis. However, length and sequence variability may require extensive optimization of sequence alignment procedures to ensure that the resulting trees are accurate (Morrison and Ellis, 1997). Heterogeneity in the ribosomal genes within individual nematodes has been reported (Blok et al., 1998; Cherry et al., 1997; Subbotin, et al., 2000; Zijlstra et al., 1995). Such variation may, in some instances, confound phylogenetic analysis.

In spite of the need for additional molecules, only a few other nuclear genes have been employed in nematode phylogeny, including RNA polymerase II (Baldwin et al, 1997), major sperm protein (MSP) (Setterquist et al., 1996), and Hsp70 (Beckenbach et al., 1992; Hashmi et al., 1997). Both Hsp70 and Hsp90 heat shock protein genes have been used extensively for phylogenetic analysis in other animal and bacterial systems (Gupta, 1995, 1998; Krishna and Gloor, 2001; Landais et al., 2001; Pepin et al., 2001; Welch and Meselson, 2000, 2001a,b). A study of Hsp70 from sharks highlighted the

limitations of using paralogous genes for phylogeny (Martin and Burg, 2002), but when true orthologs were distinguished from paralogs and then combined with morphological information, this gene family was demonstrated to be phylogenetically useful.

Hsp90 is an attractive candidate for use in nematode phylogenetics because it is likely present in a single copy and is not easily confused with distantly related genes. While many vertebrates (Pepin et al., 2001) and plants (Krishna and Gloor, 2001) have more than one nearly identical cytoplasmic Hsp90 gene, Birnby et al. (2000) reported a single Hsp90 gene in the *Caenorhabditis elegans* genome, located on chromosome V (cosmid C47E8.5). Other *C. elegans* genes that have been loosely classified as Hsp90 family members share limited structural homology with this sequence. The most closely related sequences include a putative endoplasmic reticulum heat shock protein GRP94 on chromosome IV (cosmid T05E11.3; 46% identical aa) and a TRAP1 homolog on chromosome III (cosmid R151.7; 42% identical aa, limited to the ATP-binding domain). The only other nematode species for which a full-length Hsp90 gene has been described is the filarial nematode *Brugia pahangi* (Thompson et al., 2001). Although information regarding Hsp90 copy number in other nematodes is currently lacking, single copy Hsp90 genes have been demonstrated in several *Drosophila* species (Konstantopoulou and Scouras, 1998) and in the Lepidoptera *Bombyx mori* and *Spodoptera frugiperda* (Landais et al., 2001).

Our goals in this study were to establish that Hsp90 sequences can be readily obtained from bulked nematode DNA or single nematode specimens and to determine the copy number of Hsp90 from selected plant-parasitic nematodes. Once we met these objectives, we set out to determine the extent of Hsp90 sequence variation at the protein and DNA levels and to test the phylogenetic utility of Hsp90 in comparison to the 28S LSU and 18S SSU ribosomal DNA genes commonly

Received for publication 11 February 2004.

¹Research Molecular Biologist and ²Research Plant Pathologist, USDA-ARS Nematology Laboratory, Bldg. 011A BARC West, Rm. 165B, Beltsville, MD 20705.

The authors thank Zafar Handoo for specimens and confirmation of their identification, Charles Opperman for *Heterodera glycines* strain OP50, and Eric L. Davis for the *H. glycines* J2 cDNA library. They also thank Maria Hult, Sharon Ochs, and Matt Lewis for excellent technical assistance and David Carta for help with Matlab.

Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

E-mail: skantara@ba.ars.usda.gov

This paper was edited by J. L. Starr.

used for nematode phylogeny. Here we examine the amount of variation and types of substitutions in a conserved partial region of the Hsp90 gene and compare the branching patterns in phylogenetic trees generated by various methods.

MATERIALS AND METHODS

Nematode isolates, culture methods, and DNA extracts: *Meloidogyne javanica* and *M. arenaria* (both from North Carolina) were cultured on *Lycopersicon esculentum* cv. Big Boy. *Heterodera glycines* inbred strain OP50 (inbred from a North Carolina isolate) was maintained on *Glycine max*, cv. Lee 68. In these cases, nematode eggs were harvested by sugar flotation (Jenkins, 1964), snap-frozen in liquid nitrogen, and stored at -80°C . Bulk nematode genomic DNA was prepared from *M. javanica*, *M. arenaria*, and *H. glycines* as described (Bird and Riddle, 1989). The following species used for single-nematode PCR were obtained from various locations in Maryland: *Anguina* sp., *Ditylenchus* sp., *Meloidogyne hapla*, and *Pratylenchus crenatus*. Nematodes from other locations included *Acrobeloides nanus* PS1959 (France), *Heterodera goldeni* (Egypt), *M. floridensis* (Florida), *M. sasserii* (Delaware), *Oscheius myriophila* DF5020 (California), *P. penetrans* (Arkansas), *P. teres* (South Africa), and *Zeldia punctata* PS1145 (California). Extracts were prepared from single juveniles of the plant-parasitic or bacterial-feeding nematodes as previously described (Thomas et al., 1997). In some cases, extracts were made from a pool of two to nine juveniles to increase the amount of template DNA.

Gene amplification, cloning, and sequencing: Hsp90 was amplified using a modification of RAN-PCR (Skantar and Carta, 2000) and primers U831 [5'-AA(T/C)AA(A/G)AC(A/C)AAGCC(A/C/G/T)T(T/C)TGGAC-3'] and L1110 [5'-TC(A/G)CA(A/G)TT(G/A/C)TCCATGAT(A/G)AA(G/A/C)AC-3']. Reactions were assembled in PCR tubes by first adding a cocktail containing dNTP's and primers. This mixture was overlaid with a drop of paraffin wax that was allowed to harden, separating it from a top layer that contained the DNA template and Eppendorf MasterTaq (Brinkmann, Westbury, NY) in reaction buffer supplied by the manufacturer. PCR reactions were performed in an Eppendorf MasterCycler (Brinkmann, Westbury, NY). Cycling conditions consisted of a pre-heat step of 94°C for 10 minutes, to allow the layered components to mix in a hot start, followed by 35 cycles of 94°C for 20 seconds, 65°C for 5 seconds, 60°C for 5 seconds, 55°C for 5 seconds, 50°C for 5 seconds, and 68°C for 1 minute. A final step of 68°C for 15 minutes allowed for complete extension of products. Each 25- μl PCR reaction was analyzed on 1.5% SeaKem GTG agarose (BMA, Rockland, ME) in 1X Tris-Acetate-EDTA (TAE). PCR products were visualized with UV illumination after ethidium bromide staining. Bands containing

single PCR products were excised from the gel and, if necessary to increase yield, 2 μl was subjected to re-amplification as described above. Otherwise, excised PCR products were purified with the QIAquick Gel Extraction Kit (Qiagen, Valencia, CA), cloned into pCR2.1, and transformed into *Escherichia coli* TOP10 cells according to the manufacturer's instructions (Invitrogen, Carlsbad, CA). Plasmid DNA was prepared with a Wizard Plus miniprep kit (Promega, Madison, WI) and digested with EcoRI to verify the correct insert. Double stranded DNA was sequenced with either Big Dye 3.0 Terminator cycle sequencing reagents and analyzed with an ABI 310 Gene Analyzer (Perkin Elmer Applied Biosystems, Foster City, CA) or with the CEQ DTCS Quick Start Kit and analyzed on the CEQ8000 Genetic Analysis System (Beckman Coulter, Fullerton, CA). Sequence was determined on both strands from all clones using M13 forward and M13 reverse primers. Whether template was from bulked nematodes or single specimens, no less than two clones (typically four to six) were sequenced from each PCR reaction.

Heterodera glycines Hsp90 cDNA cloning: An *H. glycines* cDNA library in the Uni-ZAP vector was screened by PCR for the presence of the Hsp90 gene. Preparation of phage and mass-excision of the pBluescript phagemid DNA containing nematode cDNA inserts were performed according to the manufacturer's instructions. Excised phagemid DNA was used as template in PCR reactions at 30 ng per 25- μl reaction. Primer L1110 was paired with the M13 reverse primer and primer U831 was paired with the M13 forward primer to amplify the 5' and 3' ends of the *H. glycines* Hsp90 (*Hg-Hsp90*) cDNA, respectively. An additional 5' end clone was obtained by pairing the primer INV-3, [5'-GAGCGTCAGAGGAGTTGGAG-3'] with the M13 reverse primer. All PCR products were TA-cloned as described above and sequenced.

Southern hybridization: Ten μg of *H. glycines* or *M. javanica* genomic DNA was digested with BamHI in 40- μl reactions overnight at 37°C . The digests were electrophoresed overnight in 0.8% agarose/1X TBE, stained with ethidium bromide, and photographed. The DNA was transferred to nylon by standard methods (Ausubel et al., 2001). The Hsp90 probe was labeled with digoxigenin-UTP using established PCR conditions, and a chemiluminescent Southern hybridization was performed using the Genius kit (Roche Applied Science, Indianapolis, IN) as described by the manufacturer. The probe was hybridized overnight at 65°C . The blot was washed twice for 5 minutes at room temperature in 2X SSC/1% SDS and followed by two 15-minute washes at 65°C in 0.1X SSC/0.1% SDS. Hybridized products were detected with the chemiluminescent substrate CDP-Star and visualized by autoradiography.

Hsp90 amino acid sequence alignments: The deduced amino acid sequence of *Hg-Hsp90* was determined in Sequencher 4.1 (Genecodes, Ann Arbor, MI) using

standard codon tables. The organisms and GenBank accession numbers for full-length Hsp90 amino acid sequences are listed in Table 1. Hsp90 amino acid sequence alignments were performed with ClustalW (vers. 1.4) (Thompson et al., 1994) and manually adjusted in GeneDoc (Nicholas et al., 1997).

Nucleic acid sequence alignments: The phylogenetic properties of Hsp90 were examined in comparison to LSU and SSU rDNA genes commonly used in nematode systematics. Six nematode taxa were selected to broadly represent expected nematode clades (Blaxter, 2001): *B. malayi*, *Strongyloides ratti*, *C. elegans*, *M. javanica* or *M. incognita* (SSU), *H. glycines*, and either *P. teres* (Hsp90, LSU) or the closely related *Nacobbus aberrans* (SSU) (Carta et al., 2001). DNA sequence alignments were generated for the Hsp90 gene (213 bp), the LSU D3 expansion segment (178 nucleotides excluding gaps), and the SSU gene (including 832 bp, corresponding to positions 1015 through 1846 in the *C. elegans* reference sequence, Accession # X03680). Accession numbers for DNA sequences (from this study or GenBank) from these taxa are listed in Table 2. *Schistosoma mansoni* was selected as the closest outgroup for which all genes were available. Based upon results from large SSU rDNA datasets (Aguinaldo et al., 1997), *D. melanogaster*

TABLE 1. Hsp90 protein sequences included in this study.

Organism	Group, common name	Accession No.
<i>Escherichia coli</i>	enterobacteria	NP_286214
<i>Ajellomyces capsulatus</i>	fungi	AAA33383
<i>Aspergillus niger</i>	fungi	CAA72292
<i>Candida albicans</i>	fungi	CAA56931
<i>Neurospora crassa</i>	fungi	CAC28765
<i>Podospora anserina</i>	fungi	AAB97626
<i>Saccharomyces cerevisiae</i>	fungi	AAA02743
<i>Guillardia theta</i>	cryptomonads	AAF24209
<i>Dictyostelium discoideum</i>	slime molds	AAA69917
<i>Trypanosoma brucei</i>	kinetoplastids	AAC48281
<i>Leishmania amazonensis</i>	kinetoplastids	P27741
<i>Tetrahymena thermophila</i>	ciliates	AAD41357
<i>Cryptosporidium parvum</i>	apicomplexa	AAC24767
<i>Eimeria tenella</i>	apicomplexa	AAB97088
<i>Plasmodium falciparum</i>	apicomplexa	P20147
<i>Brugia pahangi</i>	filarial nematode	CAA06694
<i>Caenorhabditis elegans</i>	bacterial-feeding nematode	CAA99793
<i>Heterodera glycines</i>	soybean cyst nematode	AA014563 ^a
<i>Arabidopsis thaliana</i>	eudicots	AAA32822
<i>Lycopersicon esculentum</i>	eudicots, tomato	AAB01376
<i>Triticum aestivum</i>	monocots, wheat	CAA67191
<i>Zea mays</i>	monocots, corn	AAB26482
<i>Anopheles albimanus</i>	insects, mosquito	AAB05638
<i>Bombyx mori</i>	insects, silkworm	BAB4120
<i>Drosophila melanogaster</i>	insects, fruit fly	P02828
<i>Spodoptera frugiperda</i>	insects, armyworm	AAG44630
<i>Danio rerio</i>	bony fishes, zebrafish	Q90474
<i>Cricetulus griseus</i> , α	mammals, hamster	AAA36992
<i>Gallus gallus</i> , α	mammals, chicken	CAA30251
<i>Homo sapiens</i> , α	mammals, human	CAA33259
<i>Mus musculus</i> , α	mammals, mouse	AAA53068
<i>Sus scrofa</i> , α	mammals, pig	AAC48718

^a This study.TABLE 2. Nucleotide sequences used for seven taxon comparisons of Hsp90, 28S, and 18S.^a

Species Name	Hsp90	28S	18S
<i>Brugia malayi</i>	AW179950	AF100625	AF036588
<i>Caenorhabditis elegans</i>	Z75530	X03680	X03680
<i>Heterodera glycines</i>	AF449485 ^b	AF133304	AF216579
<i>Meloidogyne javanica</i>	AF201338 ^b	U47559	U81578 ^c
<i>Pratylenchus teres</i>	AF457583 ^b	AF196353	AF442190 ^d
<i>Strongyloides ratti</i>	BG893447	AF438342	U81581
<i>Schistosoma mansoni</i> ^e	J04017	Z46503	M62652

^a Unless noted otherwise, sequences obtained from Genbank.^b This study.^c *Meloidogyne incognita* or ^d*Nacobbus aberrans* 18S sequences were used from the closest relative for which sequence was available.^e Outgroup taxon

also was tested as an outgroup in the Bayesian analysis. Both outgroups were used for rooting trees. Introns in the Hsp90 partial genomic sequences were determined by comparing the three-frame conceptual translations to the *C. elegans* Hsp90 sequence; these regions were removed prior to further analysis. Sequences obtained from GenBank were trimmed as necessary to remove regions that did not match in all aligned taxa.

Alignments were made in ClustalW with default parameters and subsequently checked for consistency of conserved positions among sequences. Manual adjustments to remove all gaps from the ribosomal DNA alignments were performed in GeneDoc (Nicholas et al., 1997). Our objective was not to produce a definitive phylogenetic estimate using relatively sparse data from each of the three genes but to demonstrate their general substitution properties relative to one another, using unambiguous alignments. However, due to the sensitivity of tree topology to alternative alignments for SSU rDNA (Morrison and Ellis, 1997), various other trees were constructed for comparison with the conservative, non-gapped alignment and the topology abstracted from a large SSU Tree (Blaxter et al., 1998). These alignments and their corresponding trees (not shown) were made with and without gaps for both Clustal default and minimal gap extension penalty alignments. Some alignments included an extra segment represented by an additional 679 bp in the raw alignment for six taxa that was missing from *H. glycines*. For higher-density taxon sampling of nematodes, we generated partial Hsp90 sequences or used EST sequences from GenBank (Table 3). Alignments are available at the authors' Web site (<http://www.barc.usda.gov/psi/nem/staff.htm>).

Phylogenetic analysis: Maximum Parsimony (MP) analyses (Swofford, 1998; Swofford et al., 1996) were performed in PAUP vers. 4.0b10. Character state optimization was made with ACCTRAN and TBR branch swapping on 3,000 pseudoreplicates. Tree lengths, measures of consistency index (CI) for homoplasy (Kluge and Farris, 1969), and retention index (RI) for synapomorphy (Farris, 1989) were determined (Table

TABLE 3. Nematode Hsp90 sequences used for clade-specific analyses.^a

Species name	Accession #	Species name	Accession #
<i>Ancylostoma ceylanicum</i>	CB276335	<i>M. hapla</i>	AY528417 ^b
<i>Acroboloides nanus</i>	AY603518 ^c	<i>M. incognita</i>	AF459026 ^c
<i>Anguina</i> sp.	AY603510 ^c	<i>M. javanica</i>	AF201338 ^c
<i>Ascaris lumbricoides</i>	BU568768	<i>M. sasserii</i>	AF457581 ^c
<i>Brugia malayi</i>	AW179950	<i>Oscheius myriophila</i>	AY603515 ^c
<i>B. pahangi</i>	AJ005784	<i>Parastrongyloides trichosuri</i>	BI863596
<i>Caenorhabditis elegans</i>	Z75530	<i>Pratylenchus crenatus</i>	AF457582 ^c
<i>C. briggsae</i>	CBG04560	<i>P. penetrans</i>	AY603513 ^c
<i>Ditylenchus</i> sp.	AY603512 ^c	<i>P. teres</i>	AF457583 ^c
<i>Heterodera glycines</i>	AF449485 ^c	<i>Strongyloides ratti</i>	BG893447
<i>Heterodera goldeni</i>	AY603511 ^c	<i>S. stercoralis</i>	BF014947
<i>M. arenaria</i>	AF457580 ^c	<i>Trichinella spiralis</i>	BG520497
<i>M. chitwoodi</i>	CB830188	<i>Zeldia punctata</i>	AY603516 ^c
<i>M. floridiae</i>	AY603514 ^c		

^a Unless noted otherwise, sequences obtained from Genbank.

^b Described in another study.

^c This study.

4). Retention Index describes the proportion of synapomorphy from the data that is retained on a tree, with larger values representing more consistency of the characters with the cladogram. Unlike the CI, empty information from autapomorphies is not included. A test for data structure was also generated in PAUP, with *gI* skewness scores from a million random trees compared to a table of critical values (Hillis and Huelsenbeck, 1992). For Hsp90, SSU, and LSU, an entropy-based statistical test for saturation of alignment positions (Xia et al., 2003) was made with DAMBE (Xia and Xie, 2001). Scatter plots of successive pairwise transi-

tions and transversions as a function of distance were constructed from base composition data in PAUP and entered in MatLab vers. 6.5.1 (The Mathworks, Natick, MA). The evolutionary models of sequence substitution for each alignment were determined by Model Test (Posada and Crandall, 1998); parameters are listed in Table 4. The hLRT criterion was selected over the AIC criterion for consistency because some of its parameters are used within MrBayes. These parameters were employed in distance settings in PAUP. Neighbor-joining (NJ) distance trees with 3,000 pseudoreplicates were generated with among-site rate variation parameters and with log-determinant distance minimum evolution (Log-Det ME) options (Lockhart et al., 1994). Parsimony was originally conceived as an abbreviated version of maximum likelihood (ML) without as much computational complexity (Page and Holmes, 1998). Maximum likelihood is the phylogenetic estimate method least affected by sampling error and generally provides better tree inference than distance or parsimony methods (Swofford et al., 1996) when the appropriate model of substitution is not violated (Siddall, 1998). Phylogenetic likelihood, which is proportional to the probability of the data conditional on a tree, underlies both ML and Bayesian inference. We performed Bayesian phylogeny estimations (MB) using MrBayes vers. 3.0b4 (Huelsenbeck et al., 2000). MrBayes is a statistical program that permits incorporation of prior tree information. Because it employs the more computationally efficient Markov Chain Monte Carlo (MCMC) method for generating posterior probabilities (Huelsenbeck et al., 2002), MrBayes can search several hundred orders of magnitude of tree space more rap-

TABLE 4. Properties from ModelTest performed for equivalent seven-taxon nucleotide alignments of Hsp90, 18S SSU, and 28S LSU, and for a 16-taxon alignment of Hsp90.

Parameter	Hsp90 (seven-taxon)	SSU rDNA (seven taxa)	LSU rDNA (seven taxa)	Hsp90 (16 taxa)
hLRT Criteria	TrNef + γ	TIM + γ	K80	GTR + γ
nst	6	6	2	6
γ shape	0.32	0.83	—	0.29
Prop. invariant sites	0	0	0	0
Ti/Tv	—	—	1.31	—
–lnL	1047.52	4372.92	2070.47	1392.72
AIC Criteria	SYM + γ	TIM + γ	K81	SYM + I
nst	6	6	6	6
γ shape	0.31	0.87	—	1.30
Prop. invariant sites	0	0	0	0.45
Ti/Tv	—	—	1.31	—
AUC	2094.56	4959.13	4139.84	2796.14
Alignment characters	213	831	178	213
Variable	45%	60%	94%	44%
Parsimony informative	27%	24%	56%	36%
Tree length	200	830	219	319
CI (homoplasy)	0.74	0.83	0.78	0.45
RI (synapomorphy)	0.39	0.50	0.63	0.39

AIC = Akaike Information Criteria; CI = Consistency Index; GTR = General Time Reversible Model (Tavaré, 1986); γ = gamma distribution; hLRT = hierarchical Likelihood Ratio Test; I = Invariable sites; K80, K81 = Models of Kimura (Kimura, 1980; Kimura, 1981); L = Likelihood; nst = number of substitution parameters; RI = Retention Index; SYM = Symmetrical Model (Zharkikh, 1994); Ti/Tv = ratio of transitions to transversions; TIM = Transitional model (rAC = rGT \neq rAT = rCG = rAG \neq rGT, Posada and Crandall, 2001); TrNef = Tamura-Nei model, equal base frequencies (Tamura and Nei, 1993).

idly than traditional maximum likelihood methods. Beginning with a random tree, MCMC was run at least twice, for 250,000 to 10^6 cycles with each 100th cycle sampled. Branch lengths were recorded to allow generation of a consensus phylogram. The first 10% of samples were excluded as burn in and examined for convergence before creating 50% majority-rule consensus trees from the remaining trees in PAUP. The numbers reported on the consensus trees represent the probability a clade was supported among sampled trees, analogous to bootstrap support using ML parameters (Huelsenbeck et al., 2000).

RESULTS

Amplification and cloning of Hsp90 from plant-parasitic nematodes: Degenerate PCR primers U831 and L1110 were used previously in a ramped-annealing PCR reaction (RAN-PCR) to amplify the Hsp90 gene from *H. glycines*. These primers and PCR conditions were designed to specifically amplify the gene for cytoplasmic Hsp90, thus reducing the chance of obtaining unwanted Hsp90 paralogs. Using genomic DNA from either bulked nematodes or crude single-nematode DNA extracts for template, we generated Hsp90 PCR products from a broad range of plant-parasitic and free-living species (Tables 2 and 3). Each PCR reaction typically generated a single major band in the size range predicted from the *C. elegans* Hsp90 gene (not shown). Excision and purification of Hsp90 PCR products from agarose gels were routinely performed to remove any minor reaction products or primer dimers prior to cloning and sequencing.

To determine whether *H. glycines* Hsp90 was present in a single genomic location, the 350-bp *Hg-Hsp90* PCR product was hybridized to restriction enzyme-digested *H. glycines* genomic DNA. The probe detected a single band in the Hind III-digested DNA and two bands in the Eco RI-digested DNA (Fig. 1A). DNA sequencing revealed a single Eco RI site in the middle of the *Hg-Hsp90* PCR probe, so two bands would be expected in the Southern hybridization to Eco RI-digested DNA. Sequencing also showed a Hind III site located 39 bp from the 3' end of the amplified probe. Thus, the lack of detectable hybridization to a second band in the Hind III digested lane is not surprising and may have been due to insufficient pairing between the probe and the blotted restriction products. The *Hg-Hsp90* probe also hybridized to a blot of Bam HI-digested *M. javanica* genomic DNA (Fig. 1B). A single band was observed, consistent with the presence of a single copy of Hsp90 in this species. Taken together, these results mirror the single Hsp90 gene that was found in *C. elegans*, whose entire genome sequence is known.

To isolate the Hsp90 cDNA, an excised *H. glycines* second-stage juvenile cDNA library was used as PCR template for PCR. In these reactions, the U831 and




FIG. 1. Evidence that Hsp90 represents a single gene in the genomes of *Heterodera glycines* and *Meloidogyne javanica*. A Southern blot of restriction enzyme-cleaved genomic DNA extracted from plant-parasitic nematode second-stage juveniles was hybridized with a probe obtained by PCR amplification of the *H. glycines* Hsp90 gene, and detected by chemiluminescence. Bands that hybridized to the Hsp90 probe are indicated by arrows. Numbers at left indicate molecular size markers (1 kb ladder, New England Biolabs, Beverly, MA). A) *Heterodera glycines* DNA digested with Hind III (lane 1) or Eco RI (lane 2). B) *Meloidogyne javanica* DNA digested with Bam HI.

L1110 primers were paired separately with either the M13 forward or reverse vector primers. The 5' and 3' PCR products were cloned and sequenced, and the contiguous cDNA sequence was reconstructed. Upon sequencing the 5' end clone obtained from the pairing of L1110 with M13 reverse, we found that this cDNA lacked the initiator methionine codon. We then performed an additional PCR amplification of the library cDNA's, using the internal *Hg-Hsp90* primer (INV-3) paired with the M13 reverse primer. The PCR product of this reaction was cloned and sequenced, revealing the genuine 5' end of the cDNA. Attempts to obtain the 5' end by pairing INV-3 or other internal primers with an SL1 spliced leader primer were unsuccessful, suggesting that unlike *C. elegans daf-21*, *Hg-Hsp90* does not undergo trans-splicing (not shown).

Hg-Hsp90 sequence analysis: BLAST analysis of the *Hg-Hsp90* amino acid sequence revealed a high level of similarity to other members of this protein family. The alignment with two other available full-length nematode Hsp90 protein sequences and representative

In *H. glycines*, this 45 amino acid domain was 14 residues longer than the corresponding region from *C. elegans*, five residues longer than *B. pahangi*, but comparable in length to *D. melanogaster* and *M. musculus*. This region was followed by a highly conserved domain, which contains the amino acid residues used for design of the degenerate PCR primers, U831 and L1110 (boxed in black). A third highly variable region was located between residues 679–707, just upstream of the C-terminal MEEVD motif found only in cytoplasmic Hsp90 sequences. The one known *C. elegans* Hsp90 mutation, *daf-21(p673)*, appears at position 293, buried within a 38 amino acid region that was highly conserved across all taxa. Amino acids that were identical in *H. glycines* and one or more of the other nematodes, but not in the other species, were dispersed throughout the sequence (Fig. 2, shaded in gray).

Amino acid phylogeny: There was sufficient phyloge-

netic information in the full protein sequence to provide strong support for most branches (Fig. 3). The nematode clade branched basal to the insects. The clade including insects plus vertebrates was supported at 70%. Although *S. mansoni* would have been a desirable outgroup to the nematodes, a full-length amino acid sequence was not available for comparison.

Nucleic acid phylogeny: To determine the phylogenetic value of Hsp90 relative to the LSU and SSU rDNA genes traditionally used for nematodes, substitution parameters resulting from ModelTest were determined for each molecule and subjected to MB and ME analyses. Hsp90, SSU, and LSU each exhibited unique data set patterns derived from MP analyses of the seven-taxon alignments (trees not shown), Bayesian probability trees (Fig. 4), and a scatter plot of sequence variations (Fig. 5). Each data set resulted in a single bootstrapped consensus tree. MP analysis results are shown

	V	
HGL	NCELMPEYLNFKGVVDS EDLPLNISRETLQQSKILKVKIRKNLVKKCMDLFEI SEDKDNEK KFYEQFAKNIKL	434
BPA	..D.....M.....LE..D..A.....S.....	429
CELM.....E..ID.VA.....G..L..	418
MMUI.....R.....M.....LE..T.LA...E.Y.....S.....	447
DDI	..ADII.....VR.I.....N..T.....IE..N..A.NSEDY...A.S..L..	415
DME	..D.I.....M.....M..N.V.....T.E.I..LT...E.Y...D..S..L..	432
ATHI...S.V.....D.....N.....IEM.N..A.N.EDYT...A.S..L..	419
HGL	GIHEDSVNRK KLSDFL RYITSASG EP CSFKQDYVSRMKENQ IC IYYITGESKDVVQNSSFVERVKKRGFEVIYM	509
BPAT.....E...F...S..MT.L.-.....KQ..F...REA.AS.A.....R.....	503
CELT.....S.....D..T.L.-E.....Q.....AA.A.....S.....L..	491
MMUQ.....EL.....D.MV.L.-..CT.....KH..F...T..Q.A.A...LR.H.L....	521
DDI	.V...Q..E.FA.L...Q..K..D.LVTL.-E..G...G.NE.....KA.E..P.I.GL..KNL....	489
DME	.V...N..A..A...FH...DDF..LA-.....D..KHF.F.....Q.S..A...A...V..	506
ATHQ..G.IA.L...HSTK..D.MT...-..T...G.KD.F.....KA.E..P.L..L...Y..L..	493
HGL	VDPID EY CVQQLKEYDGKKLVSVTK EGLELP -E SEEE KKK FEED KVKFEKLCKVIKDI-DKKVQKVSNSRLVSS	582
BPA	T.....N.....M...LE...E..A.....	577
CEL	C.....-T.....AY.N.....LE...E..G.....	565
MMU	IE.....FE..T.....D.....Q..K.T..N..IM..LE...E..V.....T..	595
DDI	C.....A.....I.....K.D-T.D...A.Q..AAN.E.L.QV..VLGD..E..VL.T..AN..	563
DME	TE...VI.H...K..Q.....D.S...R...A...S...LM.S.L.N..E..V.....D..	580
ATH	..A...A.G.....A...K.ED.T.....R..K.KS..N...T..E.LGD..E..V..D.I.D..	568
HGL	PCCI VTGEY GWSANMERIMRAQALRDSSTMGY MASK KNLEINPDHSIIKSLRDRVEKEQDDKTAKDLVLLYETS	657
BPAS.....K.....A..H.....V..A..E...ADKN...V.....F..A	652
CELS.....K.....A..H.....A.M.T.....VDKN...V.....F..A	640
MMUST..T.....K.....N.....A..H.....ET..QKA.ADKN..SV...I.....A	670
DDI	..VL..S.....K.....MSS..S..T..L...P.VRD.AKKAEE--KS..F..F.Y...A	636
DMESQF.....K.....TA.....G..Q.....P.VET..QKADADKN..AV...I..F...	655
ATH	..L.....T.....K.....MS...S...TM....NG.MEE..K.A.ADKN..SV...M....A	643
	VI	
HGL	LLTSGFSLEDPQ QH ASRIYRMV KGLG LIPDE EP AEQ QP STSG EP TIAEKI AGAE E EA SRMEEVD	722
BPA	..S.....L.....I.....TED..EEAIASVS GK EDCV PN LV...D.....	717
CEL	..A.....E..S.....I.....G.D.IEDSAV..SC---AEA..E...D.....	702
MMU	..S.....T..N...I...G.DEDDPTVDDTSAAVT---EEMPPLEGDDDT.....	732
DDIDE.SSF...H..I...S.Q.DSSSTTEESTNTTTSDDIPPLEEN-D.P.E..K..	700
DME	..S.....DS..V.....I...G.DED.PMTTDDAQSA.---D.PSLVEDT.D..H....	717
ATHDE.NTF.A..H..L...S.DED.NVE.D--G-----DMPELEEDAA.E.K.....	700

FIG. 2. Continued




FIG. 3. Hsp90 protein cladograms of Eukarya with *E. coli* as outgroup. Maximum parsimony cladogram. Branch lengths representing absolute character changes are positioned below branches, and bootstrap values from 1,000 replications positioned above. Heuristic search on an 834 amino acid character alignment with branch-swapping, character-state optimization and ACCTRAN options selected.

at the bottom of Table 4. In the LSU and SSU alignments, about 20% of the characters consist of gaps for one or more taxon, compared to only 1% for Hsp90 (due to a single codon insertion in outgroup *S. mansoni*). The MP tree length for Hsp90 showed 87% support for a clade including *C. elegans* as an outgroup to the three plant parasites. The SSU tree had 81% bootstrap support for *Brugia* as an outgroup taxon to the three plant parasites. This topology was also present in the alternative alignments tested. The LSU tree showed 65% support for *C. elegans* as an outgroup to *M. javanica* + *P. teres*, and 73% support for *H. glycines* as an outgroup to *B. malayi* + *S. ratti*. Outgroup comparisons revealed that *S. mansoni* provided shorter branch length and higher support values for in-group clades than did *D. melanogaster*. Because the *Schistosoma* results were consistent with those expected from intensively sampled SSU nematode trees (Blaxter et al., 1998), this taxon was used as the outgroup for trees shown in Figure 4. The Hsp90 tree topology (Fig. 4a) was consistent with SSU rDNA-based trees (Blaxter et al., 1998), except for

the reversal in position of *Strongyloides ratti* and *C. elegans*. A similar topology was generated with *D. melanogaster* as the outgroup, but with much lower branch support for the clade of *C. elegans*, *H. glycines*, *P. teres*, and *M. javanica* (64% vs. 99%) (tree not shown). The SSU tree (Fig. 4b) was constructed with model TIM + γ (Table 4), resulting in a clade of (*B. malayi* + (*S. ratti* + *C. elegans*, 100%) 64%) and (*H. glycines* + (*N. aberrans* + *M. incognita*) 100%). The LSU tree (Fig. 4c) was constructed with K80 model. For LSU, an unusually long branch for *H. glycines* persistently joined the more divergent animal parasites at the tree base, although the support among various tree searches was not strong. This pattern occurred even when a heuristic ML tree with the expected topology was presented to MrBayes, although support values for the unexpected grouping of *H. glycines* + *B. malayi* + *S. ratti* were notably reduced (73% to 64%). Other than this difference, the support values from duplicate MrBayes runs generally varied by as little as two to four units for any branch.

The *gI* parsimony tree scores for the three genes each were derived from a distribution of a million tree lengths that indicated significant signal in these data sets. *G1* scores (expressed as the observed score relative to a set critical value; Hillis and Huelsenbeck, 1992) were: Hsp90 = $-0.52 < -0.45$; SSU = $-0.83 < -0.49$; 28S LSU = $-1.036 < -0.46$ ($p < 0.01$ for all). We also employed a more robust, entropy-based statistical test that takes into account tree topology, sequence length, and the number of taxa to arrive at critical values for determining phylogenetic signal (Xia et al., 2003). This method of determining substitution saturation (Xia et al., 2003) showed significant signal for two of the seven-taxon data sets from Figure 4, i.e., Hsp90 (Iss = $0.30 < 0.73$ for symmetrical tree, < 0.63 for asymmetrical tree, $p < 0.0001$) and SSU (Iss = $0.35 < 0.78$ for symmetrical tree, < 0.69 for asymmetrical tree, $p < 0.0001$) but not for LSU (Iss = $0.70 < 0.73$ insignificant for symmetrical tree, $p = 0.22$, and significantly > 0.63 for asymmetrical tree, $p = 0.01$). Significant signal was present in the 16-taxon plant parasite Hsp90 alignment for Figure 6 (Iss = $0.37 < 0.68$ for symmetrical tree, $p < 0.0001$, and < 0.49 for asymmetrical tree, $p = 0.01$).

Saturation scatter plots with curve fitting of transitions and transversions for all taxon pairs were created for each data set using DAMBE (data not shown). Information from these plots was abstracted in Figure 5, which shows a snapshot of possible signal saturation in Hsp90, SSU, or LSU, in pairwise comparisons of the taxa represented in Figure 4. The data in Figure 5 are expressed as the number of faster-accumulating transitions and rarer transversions as a function of distance. For all Hsp90 taxon pairs except for *Brugia-Strongyloides*, the number of transitions was greater than the number of transversions, which may be attributed to the relatively high AT content of *Strongyloides* (60% to 62%) compared to the average for other taxa (50% to 55%).

The inverse pattern of more transversions appeared uniformly in the LSU rDNA cluster, with the single exception of the least divergent *Meloidogyne-Pratylenchus*





FIG. 5. Scatter plot of nucleotide transitions and transversions from alignments used in Fig. 4 as a function of distance. Substitution patterns are plotted for Hsp90, 18S, and 28S genes, for taxon pairs specified as follows: 1 = *Schistosoma mansoni* = *Brugia malayi*; 2 = *Brugia malayi* - *Strongyloides ratti*; 3 = *Strongyloides ratti* = *Caenorhabditis elegans*; 4 = *Caenorhabditis elegans* = *Heterodera glycines*; 5 = *Heterodera glycines* = *Meloidogyne javanica*, *M. incognita* (18S); 6 = *Heterodera glycines* = *Pratylenchus teres*, *Nacobbus aberrans* (18S); 7 = *Meloidogyne javanica*, *M. incognita* (18S) = *Pratylenchus teres*, *Nacobbus aberrans* (18S).

pair (Fig. 5, point 7). The SSU data points showed an intermediate pattern, where sequences became more divergent as a function of distance, and transversions approached and exceeded transitions in the four most divergent pairs. The distance between the outgroup and the most basal nematode taxa, between numbered data points 1 and 2, was at least twice as great in SSU compared to Hsp90. The Bayesian inference phylogram for the Hsp90 matrix was obtained for a select group of plant-parasite taxa and rooted with outgroup *Acrobeloides nanus*, with branch support shown for ME distance bootstrap values, MP bootstrap values, and MB posterior probabilities (Fig. 6). A range of variation in simple sequence divergence between species included a 1% difference between *M. javanica* and *M. incognita*, 8% between *M. hapla* and *M. javanica*, 14% between *P. penetrans* and *P. crenatus*, and 16% between *H. goldeni* and *H. glycines*. The ME values were always higher than MP values but lower than MB probabilities, except for the 11% higher value of 100% for the clade of *M. javanica* + *M. floridensis*, + *M. incognita* + *M. arenaria*. The MP tree resolved *Ditylenchus* and *Anguina* (92%) and all

FIG. 4. Comparison of seven taxon consensus phylograms rooted with *Schistosoma mansoni*, derived from MrBayes program for Bayesian phylogeny inference using the hLRT criterion for evolutionary model parameters derived from ModelTest as implemented in PAUP, with log likelihood tree scores (-lnL). A. Hsp90 tree from a 216 nt alignment. Tree score -lnL = 1039.85. Model: TrNef + γ . B. 18S SSU rDNA tree from an 831 nt alignment. Tree score -lnL = 4372.93. Model: TIM + γ . C. 28S LSU rDNA tree from a 178 nt alignment. Tree score -lnL = 1219.26. Model: K80. Curved lines and numerals to the right of species names indicate taxon pairs in the scatter plots shown in Figure 5.




FIG. 6. MrBayes consensus phylogram for plant-parasitic nematode Hsp90, based upon a 213 bp ClustalW alignment of 16 taxa, with *Acroboloides nanus* designated as the outgroup. Tree generated in PAUP with Log likelihood score $-\ln L = 828.11$, and consensus phylogram tree scores $TL = 281$, $CI = 0.52$, $RI = 0.54$. Support values are positioned over and under branches in sequential order: bootstrap values for Minimum Evolution distance (ME, Log-Det), maximum parsimony (MP), and posterior probabilities (MB) from MrBayes. Single values on branches represent posterior MB probabilities in instances when MP and distance values were less than the branch support threshold of 50%. MB probability values derived from 50% majority-rule consensus of 2,875 trees. Substitution model parameters generated by ModelTest for GTR + γ and variable rates entered into Among Site Rate Variation Distance setting in PAUP. Maximum parsimony bootstrap analysis employed character state optimization and TBR branch swapping with 3,000 pseudo-replicates, resulting in tree scores of $TL = 319$, $CI = 0.45$, $RI = 0.39$, $gl = -0.48$.

seven *Meloidogyne* (82%), with *M. hapla* (53%) outside four other *Meloidogyne* spp. (98%). In this Hsp90 alignment, there were 77 parsimony informative characters (36% of total) where 78% of these characters were in third codon positions, plus 17 variable but uninformative characters (8%), 119 constant characters (56%), and no gaps. The best evolutionary model for this alignment was GTR + γ (Table 4). In the Bayesian tree, the position of the *Pratylenchus teres* branch was equivocal between the position shown and a position basal to other *Pratylenchus* and *Heterodera* spp. Among the sparsely sampled animal parasites, trees including *S. ratti* and *S. stercoralis* (not shown) never joined a clade with a partial sequence of *Parastrongyloides trichosuri*, whether they were rooted with *Brugia* spp. or were placed at the root of other taxa from clade V (Blaxter et al., 1998).

DISCUSSION

There is a clear need for multiple genes of different types to construct reliable phylogenies (Baldauf et al.,

2000; Kroken and Taylor, 2001; Sidow and Thomas, 1994). The nuclear gene Hsp90 possesses sufficient variation in its DNA and protein sequences to be useful for nematode phylogeny at different taxonomic levels. The Southern blots detected single Hsp90 genes in both *H. glycines* and *M. javanica*, a result we expect would be true for most, if not all, nematodes. While it would be impractical to perform the Southern analysis for every species, it is noteworthy that the U831 and L1110 primers specifically amplified single PCR products representing only the cytoplasmic form of Hsp90 from all nematodes examined thus far. These results are consistent with the single-copy nature of Hsp90 in *C. elegans*. That our primers specifically amplified only the gene for the cytoplasmic form of Hsp90 allowed us to avoid the complications of paralogy that can sometimes occur with multiple gene families. As more nematode Hsp90 genes are obtained and knowledge of taxon-specific codon usage increases, PCR primers and cycling conditions can be tailored to improve the amplification of Hsp90 from specific taxa of interest. The degenerate primers presented here serve as a useful entry point for sampling Hsp90 from diverse species, particularly when codon usage information is unknown.

Hsp90 protein sequence analysis: The *Hg-Hsp90* amino acid sequence exhibits a high level of overall conservation with the sequences from two other nematodes for which full-length sequences were available, *C. elegans* and *B. pahangi*, and lower levels of identity to other organisms. The high degree of sequence conservation in Hsp90 is advantageous for phylogeny because sequence alignments are straightforward. We have located positions of amino acid variation that may represent signature sequences for nematode Hsp90, several of which occur in or near functionally significant domains (reviewed in Buchner, 1999; Caplan, 1999; Pearl and Prodromou, 2000; and Prodromou et al., 1997).

As expected, *H. glycines* contains all of the consensus sequences previously defined by Gupta (1995, 1998). The high levels of conservation observed in regions I-IV reflect the involvement of these regions in ATP binding to Hsp90, which is essential for the protein's chaperone function (Csermely and Kahn, 1991). Hsp90-mediated regulation of steroid hormone receptor folding and activation has been widely established (Pratt, 1997), and region V is known to be critical for receptor binding (Carrello et al., 1999). The presence of a steroid-binding domain in nematode Hsp90 is fascinating given that *C. elegans daf-12*, which encodes a nuclear hormone receptor (Antebi et al., 2000), coexists in the dauer pathway with *daf-21/Hsp90*. Although an interaction between *C. elegans daf-21/Hsp90* and *daf-12* has not been directly addressed experimentally, the known biochemical features of both molecules would be consistent with Hsp90 regulation of DAF-12-mediated effects on nematode development and lifespan.

The C-terminal MEEVD sequence (region VI) in *H. glycines* also agreed with the consensus defined by Gupta (1995). This domain is required for the binding of tetratricopeptide- or TPR-containing co-chaperones to Hsp90 (Russell et al., 1999) and distinguishes cytoplasmic Hsp90 from the endoplasmic reticulum-specific paralogs identified from many eukaryotes. Examination of the region just upstream of MEEVD reveals several additional residues that appear to be unique to nematodes (Fig. 2, gray-shaded amino acids). Several putative TPR-containing co-chaperones of Hsp90 can be identified from publicly available nematode EST's or from the completed *C. elegans* genome. Thus, the extended "GAEEE/DASRMEEVD" motif may add specificity to the interaction of such co-chaperones with nematode Hsp90's.

Protein phylogeny: The Hsp90 tree (Fig. 3) shows nematodes branching outside insects, in line with the majority of other protein-coding genes that support a traditional Coelomata hypothesis (Wägele et al., 1999) over the newer Ecdysozoa hypothesis for metazoan evolution (Aguinaldo et al., 1997). In a comparison of 36 orthologous protein-coding genes from *Homo sapiens*, *D. melanogaster*, *C. elegans*, and *Saccharomyces cerevisiae*, 24 of them supported trees where insects were a sister group to humans, whereas 11 supported a clade containing both insects and nematodes (Mushegian et al., 1998). When even more genes were analyzed in similar four-taxon comparisons, trees with the highest confidence levels favored the Coelomata hypothesis at a 10:1 ratio over the Ecdysozoa (Blair et al., 2002). A more recent study of more than 500 protein sequences, alone and concatenated, also supported Coelomata (Wolf et al., 2004). While these conflicting hypotheses are still controversial (Mallatt et al., 2003), it is important to note that Hsp90 is one of those genes supporting Coelomata. This difference may increase its value in robustly comparing phylogenetic hypotheses against other genes supporting Ecdysozoa. These include actin, tubulin, and ribosomal genes, which are known to have lineage-specific biases worth testing with a more uniformly evolving character as seen in the full amino acid alignment of Hsp90 (Stechmann and Cavalier-Smith, 2003). Hsp90 protein sequences have proven especially informative in a study comparing four alveolate genes, and supported groupings of animals with fungi (opisthokonts) and plants with red algae (bikonts) more strongly than did Hsp70, despite sparser taxon sampling (Fast et al., 2002). Hsp90 protein comparisons also provided strong support for the monophyly of bikonts with Apusozoa, thus contradicting a less strongly supported position in rRNA trees (Stechmann and Cavalier-Smith, 2003).

Hsp90, SSU, and LSU comparisons: When phylogenetic signal is overwhelmed by multiple changes at each nucleotide alignment site, saturation of the data occurs. With increasing taxon divergence, saturation can be

seen when relatively high t_i/t_v drops below 1 and approaches 0.4 (Hillis, 1991). Saturation in a data set can adversely affect likelihood parameter estimates and the final tree, and long-branch attraction of random sequence similarity may increase (Philippe, 2000). While the *gI* skewness statistic for the seven-taxon alignments indicated the presence of phylogenetic signal for all three molecules, a robust new statistical method for measuring substitution saturation detected significant signal in the Hsp90 and SSU data sets, but not in the LSU data set.

Comparison of the trees and associated scatter plots for the three genes (Figs. 4, 5) shows interesting nucleotide substitution patterns that reflect this range of saturation. The portions of all three genes we examined are small relative to their total sizes. Because fractional data sets combined with sparse taxon sampling can have a large influence on tree topology, these data sets should not be used to infer which molecule is likely to provide the truest tree. However, the data are useful for comparing substitution patterns. In the SSU data, as distance between taxon pairs increases, the number of transversions approached the relative number of transitions. The LSU data were more saturated than those of either Hsp90 or SSU, as reflected in trees with relatively poor phylogenetic resolution and the apparent long-branch attraction of *H. glycines* to the animal parasites at the base of the tree. This is not surprising because the D3 region of LSU region is more commonly used with intermediate-to-shallow phylogenetic distances than the deeper divergences resolved by longer segments of SSU rDNA (Baldwin et al., 1997; Larson, 1991). A similar pattern (Fig. 5) of increased saturation in LSU over SSU rDNA data, and greater divergence among taxa for SSU than for LSU, was previously reported for fish mitochondrial mt-rRNA genes (Bakke and Johansen, 2002). Tree topologies from our Hsp90 and SSU data sets (Fig. 4a,b), both of which were less saturated than LSU, were generally consistent with the patterns seen in trees based upon broader taxon sampling of SSU (Blaxter, 2001; Blaxter et al., 1998).

One possible tree topology for the same taxa as in Figure 4 but derived from prior SSU data sets (Blaxter et al., 1998; DeLey et al., 2002) has a succession of taxa with *B. malayi*, *C. elegans*, and *S. ratti*, followed by the plant parasites *H. glycines*, *P. teres* or *N. aberrans*, and *M. javanica* or *M. incognita*. Both rDNA data sets gave a single parsimony tree in a simple search, but the topology of both trees required major topological rearrangement to approximate this expected tree. These differences probably resulted from saturated positions in this portion of the SSU from the basal taxa of *Brugia*, *Caenorhabditis*, and *Strongyloides*. The partial LSU parsimony tree had *H. glycines* and *C. elegans* reversed from their positions in the SSU trees. The Hsp90 data set was slightly less decisive, giving two trees in a simple heuristic search with surprisingly good topology. Only the

Caenorhabditis Hsp90 sequence was homoplastic within the plant parasites. The only difference in these trees involved whether *Pratylenchus* and *Meloidogyne* were joined in a clade. Likelihood methods that went beyond the observed data to model hidden substitutions were required to improve the resolution of trees from these data sets; thus, the more resolved MB trees are shown in Figure 4 rather than MP trees.

The three tree-building methods used in the Hsp90 trees (Fig. 6) showed no major conflicts in branching order, but MrBayes trees always showed higher branch support than distance or parsimony trees as observed in other studies (Mallatt et al., 2003). Similarly, maximum likelihood methods applied to an Hsp90 amino acid phylogeny of basal eukaryotes also reflected other gene trees better than either distance or parsimony methods (Stechmann and Cavalier-Smith, 2003). The Hsp90 tree topology (Fig. 4a) was nearly congruent with the expected SSU tree (Blaxter et al., 1998), except that *C. elegans* and *Strongyloides* branches were reversed. In this case, Bayesian likelihood could not overcome the questionable grouping of *Brugia* + *Strongyloides* (Fig. 4a). The topology of plant-parasitic taxa within an Hsp90-based tree (Fig. 6) is generally congruent with a previous 12-species SSU tree, which showed that *Globodera* was not the closest lineage to *Meloidogyne* (DeLey et al., 2002). Similarly, the Hsp90 tree placed *Heterodera* spp. closer to *Pratylenchus* than to *Meloidogyne*. In the SSU trees, *M. chitwoodi* and *M. hapla* were basal to the *M. incognita* group, (DeLey et al., 2002) similar to Hsp90 (Fig. 6). However, the placement of *M. javanica* differed in SSU and Hsp90 trees. In SSU, *M. javanica* joined *M. incognita*, with *M. arenaria* outside (DeLey et al., 2002). In the Hsp90 tree, *M. javanica* was basal to the group of *M. incognita* and *M. arenaria*. *Meloidogyne sasserii* was basal to the other *Meloidogyne* species. At the species level, the partial Hsp90 sequence did not resolve *M. incognita*, *M. arenaria*, and *M. floridensis*. However, a longer Hsp90 sequence or its intron sequence variation should be useful for this purpose (not shown). *Heterodera goldeni* was morphologically similar to *H. graminophila* (Handoo and Ibrahim, 2002), which also has affinity to *H. cyperi* (Mulvey and Golden, 1983). Examination of ITS rDNA supported six major species clades within Heteroiderinae and showed that the *Cyperi* group is most similar to the *Schachtii* group, including *H. glycines* (Subbotin et al., 2001). Therefore, the two *Heterodera* species in our Hsp90 tree appear to represent the two closely related clades from ITS trees. The lack of resolution of *P. teres* relative to the other two *Pratylenchus* spp. and *Heterodera* may be expected to improve with more taxa.

The Hsp90 sequence divergence between other nematode species was generally lower than mitochondrial gene differences reported for the same taxa. For instance, the 8% difference in Hsp90 between *M. hapla* and *M. javanica* was lower than the 20% difference seen

in the mitochondrial (mt) COII to cytochrome b interval (Blouin et al., 1998; Hugall et al., 1997). The 6% difference between *C. elegans* and *C. briggsae* was lower than the 10% difference in COII (Thomas and Wilson, 1991). The mean AT content of Hsp90 for 27 nematode taxa was 54%, lower than the approximately 80% AT composition for nematode mt DNA (Blouin et al., 1998; Hugall et al., 1997). Because extremes in base composition can reduce the phylogenetic utility of sequence variations, it appears that Hsp90 is less prone to this limitation than are mitochondrial genes.

The greater sequence divergence in Hsp90 between *P. crenatus* and *P. teres* (12%) compared to LSU rDNA (8%) (Carta et al., 2001) favors the utility of Hsp90 for resolving these species when more taxa are included. At the population level, two morphologically distinct populations of *P. teres* from South Africa had identical LSU D3 rDNA (Carta et al., 2002) and identical partial Hsp90 sequences (not shown). Also, two isolates of *M. hapla* with different pathogenicity (Eastern Shore, MD, and Maui, HI) were nearly identical over a longer (1078 bp) partial Hsp90 sequence, with two variable intron positions and three silent changes at third codon positions (not shown). The full length of Hsp90 may contain sufficient variation to be useful for the molecular discrimination of nematode populations.

The relatively large data set of Figure 6 showed clock-like behavior, where terminal taxa were equally distant from the root when evaluated by the AIC of ModelTest (Table 4). Other Hsp90 data sets with different taxa gave similar results (not shown). This finding is consistent with the clock-like behavior of a large amino acid alignment of eukaryotes (Stechmann and Cavalier-Smith, 2003).

The few basal animal parasite and free-living taxa for which this segment of Hsp90 sequence was available (e.g., *Trichinella spiralis*, *Brugia* spp., *Ascaris lumbricoides*, *Ancylostoma caninum*, *Oscheius myriophila*, and *C. elegans*) were too sparse and divergent to provide reasonable tree topologies with more distant taxa. However, from various data subsets we examined it was clear that *S. stercoralis* and *S. ratti* were not present in a clade with *P. trichosuri* (data not shown). This is consistent with *P. trichosuri* branching outside *Strongyloides* spp. in SSU trees (Dorris et al., 2002).

Conclusion: When variable-length intron sequences are removed, the alignment of single-copy Hsp90 sequences is relatively simple. We also found relatively uniform branch lengths in Hsp90 across nematode taxa, similar to that reported in a broader eukaryotic amino acid phylogeny (Stechmann and Cavalier-Smith, 2003). This uniformity has been attributed to the central role of Hsp90 in regulating many developmentally important proteins. Compared to the rDNA pattern, the smaller divergences of Hsp90 ingroups to outgroups may reduce the opportunity for long-branch attraction. In other organisms, as many as 100 molecules

per taxonomic sampling have been used to improve phylogenetic estimation (Bapteste et al., 2002). The fact that a relatively small part of Hsp90 demonstrates sufficient signal and produces trees generally congruent with previous trees supports the use of this gene to augment nematode phylogenetic hypotheses that are currently based upon only a few molecules.

LITERATURE CITED

- Aguinaldo, A. M. A., J. M. Turbeville, L. S. Linford, L. Hebshi, M. C. Rivera, J. R. Garey, R. A. Raff, and J. A. Lake. 1997. Evidence for a clade of nematodes, arthropods and other moulting animals. *Nature* 387:489–493.
- Antebi, A., W. H. Yeh, D. Tait, E. M. Hedgecock, and D. L. Riddle. 2000. *Daf-12* encodes a nuclear receptor that regulates the dauer diapause and developmental age in *C. elegans*. *Genes and Development* 14:1512–27.
- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl, eds. 2001. *Current protocols in molecular biology*. New York: John Wiley and Sons.
- Bakke, I., and S. Johansen. 2002. Characterization of mitochondrial ribosomal RNA genes in gadiformes: Sequence variations, secondary structural features, and phylogenetic implications. *Molecular Phylogenetics and Evolution* 25:87–100.
- Baldauf, S. L., A. J. Roger, I. Wenk-Siefert, and W. F. Doolittle. 2000. A kingdom-level phylogeny of eukaryotes based on combined protein data. *Science* 290:972–977.
- Baldwin, J. G., L. M. Frisse, J. T. Vida, C. D. Eddleman, and W. K. Thomas. 1997. An evolutionary framework for the study of developmental evolution in a set of nematodes related to *Caenorhabditis elegans*. *Molecular Phylogenetics and Evolution* 8:249–259.
- Bapteste, E., H. Brinkmann, J. A. Lee, D. V. Moore, C. W. Sensen, P. Gordon, L. Durufle, T. Gaasterland, P. Lopez, M. Muller, and H. Philippe. 2002. The analysis of 100 genes supports the grouping of three highly divergent amoebae *Dictyostelium*, *Entamoeba*, and *Mastigamoeba*. *Proceedings of the National Academy of Sciences USA* 99:1414–1419.
- Beckenbach, K., M. J. Smith, and J. M. Webster. 1992. Taxonomic affinities and intra- and interspecific variation in *Bursaphelenchus* spp. as determined by polymerase chain reaction. *Journal of Nematology* 24:140–147.
- Bird, D. Mc., and D. L. Riddle. 1989. Molecular cloning and sequence of *ama-1*, the gene encoding the largest subunit of *Caenorhabditis elegans* RNA polymerase II. *Molecular and Cellular Biology* 9:4119–4130.
- Birnby, D. A., E. M. Link, J. J. Vowels, H. Tian, P. L. Colacurcio, and J. H. Thomas. 2000. A transmembrane guanylyl cyclase (DAF-11) and Hsp90 (DAF-21) regulate a common set of chemosensory behaviors in *Caenorhabditis elegans*. *Genetics* 155:85–104.
- Blair, J. E., K. Ikeo, T. Gojobori, and S. B. Hedges. 2002. The evolutionary position of nematodes. *BioMed Central Evolutionary Biology* 2:1–7.
- Blaxter, M. 2001. Molecular analysis of nematode evolution. Pp. 1–25 in M. W. Kennedy, and W. Harnett, eds. *Parasitic nematodes*. Wallingford, UK: CABI Publishing.
- Blaxter, M. L., P. DeLey, J. R. Garey, L. X. Liu, P. Scheldeman, A. Vierstraete, J. R. Vanfleteren, L. Y. Mackey, M. Dorris, L. M. Frisse, J. T. Vida, and W. K. Thomas. 1998. A molecular evolutionary framework for the phylum Nematoda. *Nature* 392:71–74.
- Blok, V. C., G. Malloch, B. Harrower, M. S. Phillips, and T. C. Vrain. 1998. Intraspecific variation in ribosomal DNA in populations of the potato cyst nematode *Globodera pallida*. *Journal of Nematology* 30:262–274.
- Blouin, M. S., C. A. Yowell, C. H. Courtney, and J. B. Dame. 1998. Substitution bias, rapid saturation, and the use of mtDNA for nematode systematics. *Molecular Biology and Evolution* 15:1719–1727.
- Buchner, J. 1999. Hsp90 and Co. – a holding for folding. *Trends in Biochemical Science* 24:136–141.
- Caplan, A. J. 1999. Hsp90's secrets unfold: New insights from structural and functional studies. *Trends in Cell Biology* 9:262–268.
- Carrello, A., E. Ingley, R. F. Minchin, S. Tsai, and T. Ratajczak. 1999. The common tetratricopeptide repeat acceptor site for steroid receptor-associated immunophilins and hop is located in the dimerization domain of Hsp90. *Journal of Biological Chemistry* 274:2682–2689.
- Carta, L. K., Z. A. Handoo, A. M. Skantar, J. van Biljon, and M. Botha. 2002. A Redescription of *Pratylenchus teres* Khan & Singh, 1974 (Nemata: Pratylenchidae), with the description of a new subspecies from South Africa, and a phylogenetic analysis of related species. *African Plant Protection* 8:13–24.
- Carta, L. K., A. M. Skantar, and Z. A. Handoo. 2001. Molecular, morphological and thermal characters of 19 *Pratylenchus* spp. and relatives using the D3 segment of the nuclear LS rRNA gene. *Nematotropa* 31:195–209.
- Cherry, T., A. L. Szalanski, T. C. Todd, and T. O. Powers. 1997. The internal transcribed space region of *Belonolaimus* (Nemata: Belonolaimidae). *Journal of Nematology* 29:23–29.
- Csermely, P., and C. R. Kahn. 1991. The 90-kDa heat shock protein (hsp-90) possesses an ATP binding site and autophosphorylating activity. *Journal of Biological Chemistry* 266: 4943–4950.
- DeLey, I. T., P. DeLey, A. Vierstraete, G. Karszen, M. Moens, and J. Vanfleteren. 2002. Phylogenetic analyses of *Meloidogyne* small subunit rDNA. *Journal of Nematology* 34:319–327.
- Dorris, M., M. E. Viney, and M. L. Blaxter. 2002. Molecular phylogenetic analysis of the genus *Strongyloides* and related nematodes. *International Journal for Parasitology* 32:1507–1517.
- Farris, J. S. 1989. The retention index and the rescaled consistency index. *Cladistics* 5:417–419.
- Fast, N. M., X. Lingru, S. Bingham, and P. J. Keeling. 2002. Re-examining alveolate evolution using multiple protein molecular phylogenies. *Journal of Eukaryotic Microbiology* 49:30–37.
- Graybeal, A. 1994. Evaluating the phylogenetic utility of genes: A search for genes informative about deep divergences among vertebrates. *Systematic Biology* 43:174–191.
- Gupta, R. S. 1995. Phylogenetic analysis of the 90 kD heat shock family of protein sequences and an examination of the relationship among animals, plants, and fungi species. *Molecular Biology and Evolution* 12:1063–1073.
- Gupta, R. S. 1998. Life's third domain (*Archaea*): An established fact or an endangered paradigm? A new proposal for classification of organisms based on protein sequences and cell structure. *Theoretical Population Biology* 54:91–104.
- Handoo, Z. A., and I. K. A. Ibrahim. 2002. Description and SEM observations of a new species of cyst nematode *Heterodera goldeni* (Nematoda: Heteroderidae) attacking *Panicum coloratum* in Egypt. *Journal of Nematology* 34:312–318.
- Hashmi, G., S. Hashmi, S. Selvan, P. S. Grewal, and R. Gaugler. 1997. Polymorphism in heat shock protein gene (*hsp 70*) in entomopathogenic nematodes (Rhabditida). *Journal of Thermal Biology* 22: 143–149.
- Hillis, D. M. 1991. Discriminating between phylogenetic signal and random noise in DNA sequences. Pp. 278–294 in M. M. Miyamoto, and J. Cracraft, eds. *Phylogenetic analysis of DNA sequences*. New York: Oxford University Press.
- Hillis, D. M., and J. P. Huelsenbeck. 1992. Signal, noise, and reliability in molecular phylogenetic analyses. *Journal of Heredity* 83: 189–195.
- Huelsenbeck, J. P., B. Larget, R. E. Miller, and F. Ronquist. 2002. Potential applications and pitfalls of Bayesian inference of phylogeny. *Systematic Biology* 51:673–688.
- Huelsenbeck, J. P., F. Ronquist, and B. Hall. 2000. MrBayes: A program for the Bayesian inference of phylogeny (software manual). New York: University of Rochester.
- Hugall, A., J. Stanton, and C. Moritz. 1997. Evolution of the AT-rich mitochondrial DNA of the root-knot nematode, *Meloidogyne hapla*. *Molecular Biology and Evolution* 14:40–48.
- Jenkins, W. R. 1964. A rapid centrifugal-flotation technique for separating nematodes from soil. *Plant Disease Reporter* 48: 692.

- Kluge, A. G., and J. S. Farris. 1969. Quantitative phylogenetics and the evolution of anurans. *Systematic Zoology* 18:1–32.
- Konstantopoulou, I., and Z. G. Scouras. 1998. The heat-shock gene hsp83 of *Drosophila auraria*: Genomic organization, nucleotide sequence, and long antiparallel coupled ORFs (LAC ORFs). *Journal of Molecular Evolution* 46:334–343.
- Krishna, P., and G. Gloor. 2001. The Hsp90 family of proteins in *Arabidopsis thaliana*. *Cell Stress and Chaperones* 6:238–246.
- Kroken, S., and J. W. Taylor. 2001. A gene genealogical approach to recognize phylogenetic species boundaries in the lichenized fungus *Letharia*. *Mycologia* 93:38–53.
- Landais, I., J. J. Pomet, K. Mita, J. Nohata, S. Gimenez, P. Fournier, G. Devauchelle, M. Duonor-Cerutti, and M. Ogliaistro. 2001. Characterization of the cDNA encoding the 90 kDa heat-shock protein in the Lepidoptera *Bombyx mori* and *Spodoptera frugiperda*. *Gene* 271:223–231.
- Larson, A. 1991. Evolutionary analysis of length-variable sequences: Divergent domains of ribosomal RNA. Pp. 221–248 in M. M. Miyamoto, and J. Cracraft, eds. *Phylogenetic analysis of DNA sequences*. New York: Oxford University Press.
- Lockhart, P. J., M. A. Steel, M. D. Hendy, and D. Penny. 1994. Recovering evolutionary trees under a more realistic model of sequence evolution. *Molecular Biology and Evolution* 11:605–612.
- Mallatt, J. M., J. R. Garey, and J. W. Shtultz. 2003. Ecdysozoan phylogeny and Bayesian inference: First use of nearly complete 28S and 18S rRNA gene sequences to classify the arthropods and their kin. *Molecular Phylogenetics and Evolution*. In press.
- Martin, A. P., and T. M. Burg. 2002. Perils of paralogy: Using HSP70 genes for inferring organismal phylogenies. *Systematic Biology* 51:570–587.
- Morrison, D. A., and J. T. Ellis. 1997. Effects of nucleotide sequence alignment on phylogeny estimation: A case study of 18S rDNAs of Apicomplexa. *Molecular Biology and Evolution* 14:428–441.
- Mulvey, R. H., and A. M. Golden. 1983. An illustrated key to the cyst-forming genera and species of Heteroderidae in the Western hemisphere with species morphometrics and distribution. *Journal of Nematology* 11:159–176.
- Mushegian, A.R., J. R. Garey, J. Martin, and L. Liu. 1998. Large-scale taxonomic profiling of eukaryotic model organisms: A comparison of orthologous proteins encoded by the human, fly, nematode, and yeast genomes. *Genome Research* 8:590–598.
- Nicholas, K. B., H. B. Nicholas, Jr., and D. W. Deerfield, II. 1997. GeneDoc: Analysis and visualization of genetic variation. *EMBNEW.NEWS* 4:14.
- Page, R. D. M., and E. C. Holmes. 1998. *Molecular evolution, a phylogenetic approach*. Oxford: Blackwell Science Ltd.
- Pearl, L. H., and C. Prodromou. 2000. Structure and in vivo function of HSP90. *Current Opinion in Structural Biology* 10:46–51.
- Pepin, K., F. Momose, N. Ishida, and K. Nagata. 2001. Molecular cloning of horse Hsp90 cDNA and its comparative analysis with other vertebrate Hsp90 sequences. *Journal of Veterinary and Medical Science* 63:115–124.
- Philippe, H. 2000. Long-branch attraction and protist phylogeny. *Protist* 151:307–316.
- Posada, D., and K. A. Crandall. 1998. MODELTEST: Testing the model of DNA substitution. *Bioinformatics* 14:817–818.
- Pratt, W. B. 1997. The role of the Hsp90-based chaperone system in signal transduction by nuclear receptors and receptors signaling via MAP kinase. *Annual Review of Pharmacology and Toxicology* 37:297–326.
- Prodromou, C., S. M. Roem, P. W. Piper, and L. H. Pearl. 1997. A molecular clamp in the crystal structure of the N-terminal domain of the yeast HSP90 chaperone. *Nature Structural Biology* 4:477–482.
- Regier, J. C., Q. Q. Fang, C. Mitter, R. S. Peigler, T. J. Friedlander, and M. Alma Solis. 1998. Evolution and phylogenetic utility of the *period* Gene in Lepidoptera. *Molecular Biology and Evolution* 15:1172–1182.
- Russell, L. C., S. R. Whitt, M.-S. Chen, and M. Chinkers. 1999. Identification of conserved residues required for the binding of a tetratricopeptide repeat domain to heat shock protein 90. *Journal of Biological Chemistry* 274:2682–2689.
- Setterquist, R. A., G. K. Smith, R. Jones, and G. E. Fox. 1996. Diagnostic probes targeting the major sperm protein gene that may be useful in the molecular identification of nematodes. *Journal of Nematology* 28:414–421.
- Siddall, M. 1998. Success of parsimony in the four-taxon case: Long-branch repulsion by likelihood in the Farris zone. *Cladistics* 14:209–220.
- Sidow, A., and W. K. Thomas. 1994. A molecular evolutionary framework for eukaryotic model organisms. *Current Biology* 4:596–603.
- Skantar, A. M., and L. K. Carta. 2000. Amplification of hsp90 homologs from plant-parasitic nematodes using degenerate primers and ramped annealing PCR. *Biotechniques* 29:1182–1186.
- Stechmann, A., and Cavalier-Smith, T. 2003. Phylogenetic analysis of eukaryotes using heat-shock protein Hsp90. *Journal of Molecular Evolution* 57:408–419.
- Subbotin, S. A., A. Vierstraete, P. DeLey, J. Rowe, L. Waeyenberge, M. Moens, and J. R. Vanfleteren. 2001. Phylogenetic relationships within the cyst-forming nematodes (Nematoda, Heteroderidae) based on analysis of sequences from the ITS regions of ribosomal DNA. *Molecular Phylogenetics and Evolution* 21:1–16.
- Subbotin, S. S. L. Waeyenberge, and M. Moens. 2000. Identification of cyst-forming nematodes of the genus *Heterodera* (Nematoda: Heteroderidae) based on the ribosomal DNA-RFLP. *Nematology* 2:153–164.
- Swofford, D. L. 1998. PAUP* Phylogenetic Analysis Using Parsimony (*and other methods), version 4.0, Sunderland, MA: Sinauer Associates.
- Swofford, D. L., G. J. Olsen, P. J. Waddell, and D. M. Hillis. 1996. Chapter 11, Phylogenetic inference. Pp. 407–513 in D. M. Hillis, C. Moritz, and B. K. Mable, eds. *Molecular systematics*, 2nd ed., Sunderland, MA: Sinauer Associates.
- Tamura, K., and M. Nei. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* 10:512–526.
- Tang, H. and R. C. Lewontin. 1999. Locating regions of differential variability in DNA and protein sequences. *Genetics* 153:485–495.
- Tavaré, S. 1986. Some probabilistic and statistical problems on the analysis of DNA sequences. *Lectures on Mathematics in the Life Sciences* 17: 57–86.
- Thomas, W. K., J. T. Vida, L. M. Frisse, M. Mundo, and J. G. Baldwin. 1997. DNA sequences from formalin-fixed nematodes: Integrating molecular and morphological approaches to taxonomy. *Journal of Nematology* 29: 250–254.
- Thomas, W. K. and A. C. Wilson. 1991. Mode and tempo of molecular evolution in the nematode *Caenorhabditis*: Cytochrome oxidase II and calmodulin sequences. *Genetics* 128:269–279.
- Thompson F. J., A. C. Cockroft, I. Wheatley, C. Britton, and E. Devaney. 2001. Heat shock and developmental expression of hsp83 in the filarial nematode *Brugia pahangi*. *European Journal of Biochemistry* 268:5808–5815.
- Thompson J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties, and weight matrix choice. *Nucleic Acids Research* 22:4673–4680.
- Wägele, J. W., T. Erickson, P. J. Lockhart, and B. Misof. 1999. The ecdysozoa: Artifact or monophylum? *Journal of Zoological Systematics and Evolutionary Research* 37:211–223.
- Welch, D. M., and M. Meselson. 2000. Evidence for the evolution of bdelloid rotifers without sexual reproduction or genetic exchange. *Science* 288:1211–1215.

Welch, D. B. M., and M. Meselson. 2001a. A survey of introns in three genes of rotifers. *Hydrobiologia* 446/447:333–336.

Welch, D. M., and M. Meselson. 2001b. Rates of nucleotide substitution in sexual and asexually reproducing rotifers. *Proceedings of the National Academy of Sciences, USA*. 98:6720–6724.

Wolf, Y. I., I. B. Rogozin, and E. V. Koonin. 2004. Coelomata and not ecdysozoa: Evidence from genome-wide phylogenetic analysis. *Genome Research* 14:29–36.

Xia, X., and Z. Xie. 2001. DAMBE: Software package for data analy-

sis in molecular biology and evolution. *Journal of Heredity* 92:371–373.

Xia, X., Z. Xie, M. Salemi, L. Chen, and Y. Wang. 2003. An index of substitution saturation and its application. *Molecular Phylogenetics and Evolution* 26:1–7.

Zijlstra, C., A. E. M. Lever, B. J. Uenk, and C. H. Van Silfhout. 1995. Differences between ITS regions of isolates of root-knot nematodes *Meloidogyne hapla* and *M. chitwoodi*. *Phytopathology*. 85:1231–1237.