

## Effect of Short-Chain Fatty Acids and Soil Atmosphere on *Tylenchorhynchus* spp.<sup>1</sup>

CLAIRE F. McELDERRY,<sup>2</sup> MARSHA BROWNING,<sup>3</sup> AND JOSÉ A. AMADOR<sup>2</sup>

**Abstract:** Short-chain fatty acids can be produced under anaerobic conditions by fermentative soil microbes and have nematicidal properties. We evaluated the effects of butyric and propionic acids on death and recovery of stunt nematodes (*Tylenchorhynchus* spp.), a common parasite of turfgrass. Nematodes in a sand-soil mix (80:20) were treated with butyric or propionic acid and incubated under air or N<sub>2</sub> for 7 days at 25 °C. Amendment of soil with 0.1 and 1.0 μmol (8.8 and 88 μg) butyric acid/g soil or 1.0 μmol (74 μg) propionic acid/g soil resulted in the death of all nematodes. The composition of the soil atmosphere had no effect on the nematicidal activity of the acids. Addition of hydrochloric acid to adjust soil pH to 4.4 and 3.5 resulted in nematode mortality relative to controls (41% to 86%) but to a lesser degree than short-chain fatty acids at the same pH. Nematodes did not recover after a 28-day period following addition of 10 μmol butyric acid/g soil under air or N<sub>2</sub>. Carbon mineralization decreased during this period, whereas levels of inorganic N and microbial biomass-N remained constant. Short-chain fatty acids appear to be effective in killing *Tylenchorhynchus* spp. independent of atmospheric composition. Nematode mortality appears to be a function of the type and concentration of fatty acid and soil pH.

**Key words:** butyric acid, carbon mineralization, fatty acid, nematicide, nematodes, propionic acid, *Tylenchorhynchus* spp.

Nematode parasitism of turfgrass roots impairs ability to absorb water and nutrients, resulting in disease symptoms that include wilting, chlorosis, and stunting. Excessive root branching, galling, and necrosis may occur at feeding sites. The sandy root zone mix employed in golf putting greens construction is conducive to plant-parasitic nematode establishment and population increase (Pérez and Lewis, 2001).

Synthetic nematicides provide short-term control of plant-parasitic nematodes but populations may rebound, and there is a potential for the development of resistant biotypes (Jatala, 1986). These chemicals have proved expensive, hazardous to apply, and persistent and may contaminate groundwater (Jatala, 1986). Fenamiphos, a systemic organophosphate that acts as a nematostatin, remains available for use on golf course greens in some areas, but there have been reports of reduced efficacy due to rapid microbial degradation (Stirling et al., 1992). Recently enacted government regulations (Food Quality Protection Act [FQPA]) direct the manufacturer to cease the production of fenamiphos by 2007 (Crow, 2002). The fumigants methyl bromide and dazomet, which have been used to eliminate nematodes during complete greens renovation (Pérez and Lewis, 2001), provide short-term control. Metam sodium and 1,3-dichloropropene have been approved for limited use on fairways, roughs, driving ranges, and tees in Florida for nematode control (Crow, 2002), and 1,3-dichloropropene is also approved for use in golf courses in South Carolina, Geor-

gia, and Alabama (Dow AgroSciences, 2004). However, their use requires specialized equipment and sophisticated application techniques.

Concerns about environmental risks associated with synthetic nematicides have led researchers to look for alternatives. A wide variety of organic amendments have been tested (Browning et al., 1999; Feder, 1960; Muller and Gooch, 1982; Rodríguez-Kábana and Morgan-Jones, 1987), mainly in crop soils, in an effort to suppress plant-parasitic nematodes. In most instances their mode of action is not well understood. Mechanisms implicating toxic levels of ammonia or other chemicals present or generated in amendments by microorganisms have been proposed. Proliferation of antagonistic organisms has also been suggested.

Rodríguez-Kábana (1965) reported development of nematode-suppressive soils in rice paddies due to flooding. Johnston (1959a) attributed a reduction in *Tylenchorhynchus martini* (now *T. annulatus*) in saturated soil to the activity of the anaerobic bacterium *Clostridium butyricum*. Of the fermentation products identified from the activities of *C. butyricum*, only the short-chain fatty acids—butyric, propionic, acetic, and formic acids—exhibited nematicidal properties. Because short-chain fatty acids are also found in aerobic soils (Küsel and Drake, 1999; Rodríguez-Kábana, 1965), they may provide a safe alternative to synthetic nematicides because they occur naturally in soil and are readily oxidized to carbon dioxide and water by soil microorganisms. The development of alternative agents for control of plant-parasitic nematodes may also impact pest management in crops other than turfgrass because the availability of some synthetic nematicides used in other crop protection applications is expected to become restricted in response to FQPA regulations.

We conducted a series of experiments to evaluate the effectiveness of two short-chain fatty acids—butyric and propionic acids—in controlling plant-parasitic nematodes. Our objectives were to (i) evaluate the effects of butyric and propionic acids on *Tylenchorhynchus* spp.

Received for publication 16 April 2004.

<sup>1</sup> Contribution no. 5003 from the Rhode Island Agricultural Experiment Station, Kingston, RI.

<sup>2</sup> Laboratory of Soil Ecology and Microbiology, University of Rhode Island, Kingston, RI 02881.

<sup>3</sup> Department of Plant Sciences, University of Rhode Island, Kingston, RI 02881.

The authors thank Noel Jackson, Charles Dawson, Steve Alm, Josef Görres, and Lisa Rowley for their help during the course of this study.

E-mail: jam7740u@postoffice.uri.edu

This paper was edited by James A. LaMondin.

under both aerobic and anaerobic soil atmospheres, (ii) distinguish between the effects of pH and the fatty acids themselves, and (iii) evaluate the potential for recovery of nematode and microbial communities following exposure to nematicidal levels of butyric acid. We postulated the nematicidal activity of fatty acids to be concentration dependent and to result from factors other than acidification of the soil. In addition, we expected increased efficacy in an anaerobic soil atmosphere due to reduced mineralization of fatty acids.

## MATERIALS AND METHODS

*Source of test organisms:* Nematode-infested soil, dominated by *Tylenchorhynchus* spp., was acquired from disease sample cores submitted to the University of Rhode Island's Turf Diagnostic Clinic from golf courses throughout New England. Nematodes were extracted with Baermann trays and allowed to reproduce on creeping bentgrass (*Agrostis palustris* cv Penncross) plants grown in a greenhouse. After 6 months the plants were discarded and the nematode-infested soil was combined at a 1:4 ratio with autoclaved 80:20 (sand:soil) mix. The resulting soil had a pH of 5.2 and was stored in the dark at 4 °C.

*Effects of fatty acids and atmospheric composition on Tylenchorhynchus spp.:* Treatments consisted of soil amended with different concentrations of butyric or propionic acid and maintained under aerobic or anaerobic conditions. Butyric acid was purchased from Kodak Laboratory Chemicals (Rochester, NY), and propionic acid was purchased from Mallinkrodt Laboratory Chemicals (Phillipsburg, NJ). Distilled water was used for control treatments. Ten grams fresh weight of nematode-infested soil was placed in a 20-ml glass serum vial and fatty acid solutions (1 ml) added to yield final concentrations of 0, 0.01, 0.1, and 1  $\mu\text{mol/g}$  soil. Treatments were stirred into the soil and the vials were sealed with a rubber septum and an aluminum crimp collar. To create an anaerobic soil atmosphere, headspace gases were removed by vacuuming and replaced with nitrogen gas. All treatments were incubated in the dark for 7 days at 25 °C. Each treatment (fatty acid concentration  $\times$  atmospheric composition) was replicated 10 times. All 10 replicates were analyzed for  $\text{CO}_2$  evolution at the end of the incubation period with a Shimadzu GC-14 gas chromatograph (Shimadzu Scientific Instruments, Columbia, MD) fitted with a flame ionization detector using hydrogen as a carrier gas (Görres et al., 1998). Replicates were subsequently sampled destructively for nematodes ( $n = 6$ ), and soil moisture, and pH ( $n = 4$ ). Soil pH was determined using a 1:2 soil/water ratio and a pH electrode (Hendershot et al., 1993). Nematodes were extracted from soil samples with Baermann funnels initially and 7 days after treatment and counted.

*Effects of inorganic acid (HCl) on nematode survival:* Hydrochloric acid (0.25 and 1.75  $\mu\text{mol/g}$  soil) was added to nematode-infested soil (10 g fresh weight) in 20-ml glass vials to adjust soil pH values to 4.4 and 3.5, respectively. These values are representative of soil pH following treatment with butyric or propionic acid (as described). Vials were sealed and the headspace of replicates in the anaerobic treatment replaced with  $\text{N}_2$ . Soil samples were incubated for 7 days in the dark at 25 °C. Each treatment (pH  $\times$  atmospheric composition) was replicated 10 times and all replicates analyzed for  $\text{CO}_2$  evolution at the end of the incubation period, as described. Replicates were subsequently analyzed for nematodes ( $n = 6$ ), and soil moisture and pH ( $n = 4$ ).

*Recovery of Tylenchorhynchus spp. and microbial activity following treatment with butyric acid:* Glass vials were filled with 10 g of nematode-infested soil and butyric acid (1 ml) added to yield 10  $\mu\text{mol}$  butyric acid/g soil, with control treatments receiving 1 ml distilled water. The headspace gases in half of the vials was replaced with nitrogen gas. Following initial incubation in the dark for 7 days at 25 °C, the headspace in all vials was replaced with air. Microbial activity and nematode populations were monitored over 28 days by evaluating soil pH, nematode density, C mineralization, soil inorganic nitrogen, and microbial biomass N every 7 days.

Each treatment (butyric acid  $\times$  atmospheric composition) was replicated 14 times within an incubation period. All 14 samples were analyzed for  $\text{CO}_2$  evolution every 7 days. Vials were subsequently sampled for nematode density ( $n = 6$ ), microbial biomass N ( $n = 4$ ), and soil pH and inorganic N ( $n = 4$ ).

Levels of inorganic N ( $\text{NH}_4^+$  and  $\text{NO}_3^-$ ) were determined using the KCl extraction method (Keeney and Nelson, 1982). Analysis was performed with an Alpkem Rapid Flow Analyzer (RFA-300, OI Analytical, College Station, TX). Microbial biomass N was determined by the fumigation-extraction method (Vance et al., 1987). Potassium sulfate extracts of both fumigated and unfumigated samples were oxidized using the alkaline persulfate oxidation method (Cabrera and Beare, 1993). Microbial biomass N was calculated by subtracting N in the unfumigated soil extract from N in the fumigated extract. A conversion factor was not used because of the wide variability of values found in the literature (Van Veen et al., 1985).

*Statistical analysis:* Normality of data sets was determined using the Kolmogorov-Smirnov test. Depending on the normality of the data, either a Mann-Whitney rank sum test or a *t*-test was used to test the null hypothesis that two treatments were not drawn from populations with different means or medians.

Data sets that were not normally distributed and included more than two treatments were analyzed using a Kruskal-Wallis analysis of variance on ranks to test the null hypothesis that there was no difference among the population medians. Multiple comparisons vs. a control

were conducted using Dunn's test to compare treatment groups to the control. For normally distributed data sets including more than two treatments, a one-way analysis of variance was conducted, and multiple comparisons vs. a control were conducted using the Bonferroni *t*-test. All tests were evaluated at the 95% confidence level ( $P \leq 0.05$ ) (SigmaStat for Windows, vers. 2.03, SPSS Inc., Chicago, IL).

RESULTS

*Effects of fatty acids on Tylenchorhynchus spp., pH, and C mineralization:* Butyric acid was equally effective in killing all *Tylenchorhynchus* spp. present in soil when added at 0.1 and 1.0  $\mu\text{mol/g}$  soil, regardless of soil atmospheric composition (Table 1). By contrast, addition of 0.01  $\mu\text{mol}$  butyric acid/g soil had no impact on nematode survival relative to the control treatment. Addition of 1  $\mu\text{mol}$  butyric acid  $\text{g soil}^{-1}$  lowered soil pH under both aerobic and anaerobic conditions relative to the control ( $P \leq 0.05$ ) (Table 1).

Propionic acid was less effective than butyric acid, with a reduction in the number of nematodes observed only at 1.0  $\mu\text{mol/g}$  soil ( $P \leq 0.05$ ) (Table 1). As with butyric acid, soil atmosphere had no impact on the nematicidal activity of propionic acid. Soil pH was lower than untreated controls following addition of propionic acid at 1  $\mu\text{mol/g}$  soil under aerobic conditions and ( $P \leq 0.05$ ) 0.1 and 1.0  $\mu\text{mol/g}$  soil under anaerobic conditions ( $P \leq 0.05$ ) (Table 1).

The cumulative amount of  $\text{CO}_2$  evolved in soil treated with 0, 0.01, 0.1, and 1.0  $\mu\text{mol}$  butyric acid/g soil, and incubated in aerobic or anaerobic soil atmospheres for 7 days is shown in Figure 1. Under aerobic conditions, microbial respiration in soil treated with 0.1 and 1.0  $\mu\text{mol}$  butyric acid/g soil (0.25 and 0.37  $\mu\text{mol CO}_2/\text{g soil/day}$ , respectively) decreased relative to the control treatment (0.77  $\mu\text{mol CO}_2/\text{g soil/day}$ ) ( $P \leq 0.05$ ). Under anaerobic conditions, respiration was not reduced in any butyric acid treatment relative to the




FIG. 1.  $\text{CO}_2$  evolved over 7 days in soil amended with different concentrations of butyric or propionic acid under aerobic and anaerobic soil atmospheres. Values are means  $\pm$  SD ( $n = 10$ ). (\*) Different from unamended ( $P \leq 0.05$ ).

control. In contrast, at the lowest level of butyric acid (0.01  $\mu\text{mol/g}$  soil) the amount of  $\text{CO}_2$  evolved was higher than in the control under aerobic conditions ( $P \leq 0.05$ ). Propionic acid inhibited soil respiration under both aerobic and anaerobic atmospheres only at the highest application rate, 1.0  $\mu\text{mol/g}$  soil ( $P \leq 0.05$ ) (0.15 and 0.27  $\mu\text{mol CO}_2/\text{g soil/day}$ , respectively) compared to 0.66 and 0.35  $\mu\text{mol CO}_2/\text{g soil/day}$ , respectively, in the controls.

*Effects of inorganic acid (HCl) on nematode survival and carbon mineralization:* The addition of 0.25 and 1.75  $\mu\text{mol HCl/g}$  reduced soil pH from an initial value of 4.9 to 4.4 and 3.5, respectively (Table 2). Nematode densities declined after incubation for 7 days in acidified soil by 42% at pH 4.3 and 68% at pH 3.5 relative to the control (Table 2). Under anaerobic conditions, soil pH was reduced from 4.9 to 4.4 and 3.5, with nematode

TABLE 1. Effects of butyric and propionic acids on soil pH ( $n = 4$ ) and survival of *Tylenchorhynchus* spp. ( $n = 6$ ) after a 7-day incubation under aerobic or anaerobic conditions. Values shown are medians  $\pm$  SD.

Fatty acid	Amount added ( $\mu\text{mol/g}$ )	Aerobic		Anaerobic	
		Nematodes/10 g	Soil pH	Nematodes/10 g	Soil pH
Butyric	0.00	80 $\pm$ 54	5.3	90 $\pm$ 29	5.3
	0.01	90 $\pm$ 63	3.9	100 $\pm$ 63	4.0
	0.10	0 $\pm$ 0*	3.6	0 $\pm$ 0*	3.5
	1.00	0 $\pm$ 0*	3.1*	0 $\pm$ 0*	3.1*
Propionic	0.00	120 $\pm$ 37	5.3	90 $\pm$ 84	4.4
	0.01	120 $\pm$ 37	4.6	40 $\pm$ 25	4.5
	0.10	120 $\pm$ 84	4.0	110 $\pm$ 42	4.2*
	1.00	0 $\pm$ 0*	3.0*	0 $\pm$ 0*	3.0*

Values followed by (\*) within a column and fatty acid were significantly different from the control ( $P \leq 0.05$ ).

TABLE 2. Effects of hydrochloric acid on soil pH ( $n = 4$ ) and survival of *Tylenchorhynchus* spp. ( $n = 6$ ) after a 7-day incubation under aerobic or anaerobic conditions. Values shown are means ( $\pm$  SD) for number of nematodes and medians for soil pH.

Amount of acid added ( $\mu\text{mol/g}$ )	Aerobic		Anaerobic	
	Nematodes/10 g	Soil pH	Nematodes/10 g	Soil pH
0.00	63 $\pm$ 18	4.9	90 $\pm$ 30	4.9
0.25	37 $\pm$ 18*	4.4*	43 $\pm$ 25*	4.4
1.75	20 $\pm$ 12*	3.5*	13 $\pm$ 19*	3.5

Values followed by (\*) within a column were significantly different from the control ( $P \leq 0.05$ ).

densities declining by 52% at pH 4.4 and 86% at pH 3.5 relative to the control.

The rate of CO<sub>2</sub> evolution declined relative to the control in acidified soil under both aerobic and anaerobic conditions ( $P \leq 0.05$ ) (Fig. 2). Carbon mineralization rates in aerobic soil were 0.18 and 0.27  $\mu\text{mol CO}_2/\text{g soil/day}$  at soil pH of 4.3 and 3.5, respectively, compared to 0.69  $\mu\text{mol CO}_2/\text{g soil/day}$  in the controls (pH 4.9). Under anaerobic conditions C mineralization rates were reduced from 0.57  $\mu\text{mol CO}_2/\text{g soil/day}$  in the controls (pH 4.9) to 0.23 and 0.28  $\mu\text{mol CO}_2/\text{g soil/day}$  at soil pH of 4.4 and 3.5, respectively.

*Recovery of Tylenchorhynchus spp. and the microbial activity following treatment with butyric acid:* We selected butyric acid to study the effects of short-chain fatty acids on recovery of nematodes and of the microbial activity because it was more effective at reducing nematode numbers per g soil at a lower concentration than propionic acid. We used a higher concentration of butyric acid (10  $\mu\text{mol g soil}^{-1}$ ) than used in previous experiments to study recovery of nematodes and microbial activity under severe conditions. No live nematodes were present in treated samples held under aerobic or anaerobic conditions 7 days after treatment (Fig. 3), and no recovery of nematode populations was observed over the following 21 days regardless of headspace composition. Soil pH was lower in treated soils compared to the controls on all sampling dates ( $P \leq 0.05$ ). Under aerobic conditions, the pH of treated soil was initially 3.4, rising to 3.7 after 14 days following treatment, a value maintained throughout the remainder of the experiment. The pH range in the aerobic control treatment ranged from 4.1 to 4.7 over the same period. Soil pH was less consistent under anaerobic conditions, fluctuating from 3.5 to 4.0 in soil amendment with butyric acid and from 4.3 to 4.7 in the control treatment.

The C mineralization rate of soil amended with butyric acid (10  $\mu\text{mol/g soil}$ ) decreased under both aerobic and anaerobic conditions relative to the control on




FIG. 2. CO<sub>2</sub> evolved over 7 days as a function of soil pH under aerobic and anaerobic soil atmospheres. Soil was amended with HCl to adjust the pH. Values are means  $\pm$  SD ( $n = 10$ ). (\*) Different from control ( $P \leq 0.05$ ).




FIG. 3. Soil pH and *Tylenchorhynchus* spp. recovered following amendment of soil with butyric acid (0 and 10  $\mu\text{mol/g soil}$ ) under aerobic and anaerobic soil atmospheres. Soil pH values are means ( $n = 4$ ), and nematode values are medians ( $n = 6$ ). (\*) Different from control ( $P \leq 0.05$ ).

all sampling dates ( $P \leq 0.05$ ) (Fig. 4) and was undetectable 21 days after treatment with butyric acid.

We did not observe differences between soil treated with 10  $\mu\text{mol butyric acid/g soil}$  and untreated soil in terms of inorganic N, NO<sub>3</sub><sup>-</sup>, or NH<sub>4</sub><sup>+</sup> levels after incubation for 28 days, regardless of headspace composition (data not shown). Similarly, microbial biomass-N was not affected by treatment with butyric acid after incubation for 28 days (data not shown) under aerobic or anaerobic conditions.

## DISCUSSION

Our results support the hypothesis that short-chain fatty acids have nematicidal properties. Earlier studies involved the production of organic acids by anaerobic bacteria in saturated soils (Browning et al., 1999; Hollis and Rodriguez-Kábana, 1966; Johnston, 1959a) or immersing nematodes in an aqueous acid solution (Bagnage and Visser, 1965; Dijan et al., 1994; Johnston, 1959b; Stephenson, 1945; Tarjan, 1956). Our study demonstrated that both butyric and propionic acids reduced *Tylenchorhynchus* spp. numbers in unsaturated soil under both aerobic and anaerobic conditions.

We found that butyric acid was more effective than propionic acid in controlling *Tylenchorhynchus* spp. Similarly Johnston (1959b), who exposed *T. annulatus*




FIG. 4. CO<sub>2</sub> evolution following amendment of soil with butyric acid (0 and 10 μmol/g soil) under aerobic and anaerobic soil atmosphere. Values are means ± SD ( $n = 14$ ). (\*) Different from control ( $P \leq 0.05$ ).

to a series of fatty acid solutions, ranked the acids in order of toxicity as follows: butyric > propionic > acetic > formic, a series that correlates with molecular weight of the fatty acid. By contrast, Banage and Visser (1965) reported little difference among fatty acid solutions in toxicity to *Dorylaimus* sp. They hypothesized that as the ionization constants for the fatty acids are similar—resulting in roughly the same fraction of undissociated molecules at the same pH—permeation rates through the nematode cuticle should be similar. The methods employed by Johnston (1959b) and Banage and Visser (1965) involved a short immersion period in organic acid solution followed by a recovery period in tap water. Johnston (1959b) reported an inactivation time for *T. annulatus* in 0.01 M butyric acid (aq) to be approximately 2.5 minutes compared to 5.2 minutes reported for *Dorylaimus* (Banage and Visser, 1965). Differences in test organisms may account for the reported lack of differential toxicity among acids observed by Banage and Visser (1965). The *Dorylaimus* spp., an omnivore/predator, studied by Banage and Visser (1965) is considerably larger than most *Tylenchorhynchus* spp. Browning et al. (2004) reported that bacterivorous, fungivorous, and entomophagous nematodes followed the expected trend of increasing surface area to volume ratios, resulting in an increased surface area through which protonated forms of butyric acid can diffuse.

This resulted in a lower LD<sub>50</sub> with increasing surface-to-volume ratio. In contrast, as the surface-to-volume ratio of plant-parasitic nematodes increased, the LD<sub>50</sub> for butyric acid increased. Browning et al. (2004) also found differential sensitivity of nematodes to butyric acid based on trophic group, with plant-parasitic nematodes more sensitive than free-living nematodes. Dijan et al. (1994) reported differential sensitivity based on trophic groups from a nematode immersion study in maleic and pentanoic acid solutions.

Addition of 0.1 μmol butyric acid/g soil (8.8 μg/g) resulted in the death of 100% of *Tylenchorhynchus* spp. in 7 days in a sand/soil mixture. The lowest effective rate of butyric acid for suppression of *T. claytoni* in sand over 7 days was reported by Browning et al. (2004) as 880 μg/g, which resulted in a 99% reduction in nematode numbers. Differences in results between sand and sand/soil mixtures suggest that butyric acid is more effective at killing *Tylenchorhynchus* spp. when soil organic matter is present. The reasons for greater effectiveness are not immediately clear. Enhanced toxicity of butyric acid may be the result of compounds associated with organic matter that disrupt the integrity of the cuticle, facilitating acid diffusion. Alternatively, the presence of organic matter may result in microsites with localized low pH that would also facilitate acid diffusion.

A 7-day incubation in soil acidified with HCl to achieve a pH of 3.5 reduced *Tylenchorhynchus* spp. by 70% under aerobic conditions and 80% under anaerobic conditions relative to controls. However, incubation in soil-treated with 1 μmol butyric or propionic acid/g soil, resulting in similar pH values, eliminated nematodes completely. This indicates that a decrease in soil pH contributes to nematode death, particularly at lower pH values. Ruess and Funke (1992) reported a decline in bacterivorous nematodes following acidification of soil to pH values below 3.0 through addition of inorganic acids, whereas root and fungal-feeding nematodes actually increased over time. According to McSorley (1998) plant-parasitic nematodes appear to be unaffected by pH changes in the soil within the normal range. The values tested in the present study are below what is considered normal soil pH.

We expected fatty acids to be more effective under anaerobic conditions as mineralization of the organic acids by aerobic soil bacteria would be diminished in the absence of oxygen. Seven days following treatment, 0.1 and 1 μmol butyric acid and 1 μmol propionic acid/g soil had killed 100% of *Tylenchorhynchus* spp., whereas treatments exposed to lower concentrations of acid did not reduce numbers of nematodes significantly relative to the control, regardless of soil atmosphere. Carbon mineralization rates under aerobic conditions increased when butyric acid was added at low levels, which also were not nematocidal. These data suggest that soil microbes are able to metabolize the acids.

Similarly, Browning et al. (unpubl. results) found that growth by some soil fungi increased in the presence of low levels of butyric acid. In the present study, when acids were applied to soil at nematicidal concentrations, carbon mineralization rates declined relative to the controls. It appears that the concentrations of butyric and propionic acids required to kill nematodes also suppress microbial activity.

The high rate of butyric acid (10  $\mu\text{mol/g}$  soil) used in our recovery experiment likely accounts for the persistent nematicidal effect we observed. The decrease in carbon mineralization following application of butyric acid did not correlate with a decrease in microbial N, which remained largely unchanged relative to controls over the course of the experiment. This suggests that treatment with butyric acid suppressed the activity rather than the survival of the microbial population. Recovery of microbial activity may be more likely under field conditions, where fatty acids applied to soil would be subject to losses by volatilization, leaching, and mineralization by microorganisms outside the treated area.

We expected an increase in total extractable N over time from soils treated with butyric acid due to the release of ammonium from dead cells and inactivation of nitrifying bacteria. Nitrifiers are extremely sensitive to soil fumigants and acidic soil pH, and may be inactive for weeks or months after the application of a sterilant (Ebbels, 1971). Inactivation of nitrifying bacteria should lead to accumulation of ammonium. In our study, ammonium did not accumulate in acid-treated soils to levels higher than found in controls, possibly because of a shift in microbial community composition. Bigelow et al. (2002) reported that amendment of greens sand with sphagnum peat moss resulted in a reduction in soil pH, and a concomitant decline in populations of ammonium and nitrite oxidizers as well as denitrifiers. They suggested that acidification of the soil resulted in an increase in fungal biomass due to reduced bacterial competition. Fungi tend to be more acid tolerant than bacteria and can grow well within a pH range of 2.0 to 5.0 (Brock et al., 1994). They also incorporate a greater fraction of C into biomass and respire less than bacteria (Anderson and Domsch, 1974) so soil respiration may not necessarily increase over time.

We have demonstrated the nematicidal properties of butyric and propionic acids in suppressing *Tylenchorhynchus* spp. nematode densities in soil. Organic acids are produced naturally in soil by fermentative anaerobes and are consumed by aerobic and anaerobic microorganisms. Soil atmosphere had no effect on the efficacy of butyric and propionic acids in soil. The observed increase in C mineralization following the addition of low levels of butyric acid suggests that the soil microbial community may be able to use it as a C and energy source. Levels of organic acids added to soil may also decline via leaching and volatilization, reaching levels

that allow for their decomposition. In addition, the introduction of microbes from surrounding areas should facilitate fatty acid degradation, minimizing its persistence once it has had the intended effects. However, the low soil pH associated with nematicidal concentrations of butyric acid likely limits its use to pre-plant applications.

#### LITERATURE CITED

- Anderson, J. P. E., and K. H. Domsch. 1974. Measurement of bacterial and fungal contributions to respiration of selected agricultural and forest soils. *Canadian Journal of Microbiology* 21:314–322.
- Banage, W. B., and S. A. Visser. 1965. The effect of some fatty acids and pH on a soil nematode. *Nematologica* 11:255–262.
- Bigelow, C. A., D. C. Bowman, and A. G. Wollum II. 2002. Characterization of soil microbial population dynamics in newly constructed sand-based root zones. *Crop Science* 42:1611–1614.
- Brock, T. D., M. T. Madigan, J. M. Martinko, and J. Parker. (eds.) 1994. *Microbial ecology*. Pp. 669 in *Biology of microorganisms*. Englewood Cliffs, NJ: Prentice-Hall.
- Browning, M., C. Dawson, S. R. Alm, C. F. McElderry, and J. A. Amador. 1999. Effect of carbon amendment and soil moisture on *Tylenchorhynchus* spp. and *Hoplolaimus galeatus*. *Journal of Nematology* 31:445–454.
- Browning, M., D. B. Wallace, C. Dawson, S. R. Alm, and J. A. Amador. 2004. Differential effects of butyric acid on nematodes from four trophic groups. *Applied Soil Ecology* 27:47–54.
- Cabrera, M. L., and M. H. Beare. 1993. Alkaline persulfate oxidation for determining total nitrogen in microbial biomass extracts. *Soil Science Society of America Journal* 57:1007–1012.
- Crow, W. T. 2002. Nematode, where is thy sting? *Golf Course Management* 70:103–106.
- Dijan, C., M. Ponchet, and J. C. Cayrol. 1994. Nematicidal properties of carboxylic acids and derivatives. *Pesticide Biochemistry and Physiology* 50:229–239.
- Dow AgroSciences. 2004. Soil fumigant—Curfew [Online] [www.dowagro.com/turf/prod/curfew.htm](http://www.dowagro.com/turf/prod/curfew.htm) [Accessed 17 August 2004].
- Ebbels, D. L. 1971. Effects of soil fumigation on soil nitrogen and on disease incidence in winter wheat. *Annals of Applied Biology* 67:235–243.
- Feder, W. A. 1960. Osmotic destruction of plant-parasitic and saprophytic nematodes by the addition of sugars to soil. *Plant Disease Reporter* 44:883–885.
- Görres, J. H., M. J. DiChiaro, J. B. Lyons, and J. A. Amador. 1998. Spatial and temporal patterns of soil biological activity in a forest and an old field. *Soil Biology and Biochemistry* 30:219–230.
- Hendershot, W. H., H. Lalonde, and M. Duquette. 1993. Soil reaction and exchangeable acidity. Pp. 141–146 in M. R. Carter, ed. *Soil sampling and methods of analysis*. Boca Raton, FL: Lewis Publishers.
- Hollis, J. P., and R. Rodríguez-Kábana. 1966. Rapid kill of nematodes in flooded soils. *Phytopathology* 56:1015–1019.
- Jatala, P. 1986. Biological control of plant-parasitic nematodes. *Annual Review of Phytopathology* 24:453–489.
- Johnston, T. M. 1959a. Antibiosis of *Clostridium butyricum* Prazmowski on *Tylenchorhynchus martini* Fielding, 1956 (Nematoda: Phasmidia), in submerged rice soil. PhD dissertation, Louisiana State University, Baton Rouge, LA.
- Johnston, T. M. 1959b. Effect of fatty acid mixtures on the rice stylet nematode. *Nature* 183:1392.
- Keeney, D. R., and D. W. Nelson. 1982. Nitrogen—inorganic forms. Pp. 643–698 in L. Page, R. H. Miller, and D. R. Keeney, eds. *Methods of soil analysis, part 2A*. Madison, WI: Soil Science Society of America.
- Küsel, K., and H. L. Drake. 1999. Microbial turnover of low molecular weight organic acids during leaf litter decomposition. *Soil Biology and Biochemistry* 31:107–118.
- McSorley, R. 1998. Population dynamics. Pp. 109–134 in K. R. Barker, G. A. Pederson, and G. L. Windham, eds. *Plant-parasitic nematode interactions*. Madison, WI: American Society of Agronomy.

- Muller R., and P. S. Gooch. 1982. Organic amendments in nematode control. An examination of the literature. *Nematological Reviews* 12:319–326.
- Pérez, E., and E. Lewis. 2001. Nematodes: Turf's hidden enemy. *Grounds Maintenance* 36:26–28.
- Rodríguez-Kábana, R. 1965. Chemical antibiosis to nematodes in rice fields. PhD dissertation, Louisiana State University. Baton Rouge, LA.
- Rodríguez-Kábana, R., and G. Morgan-Jones. 1987. Biological control of nematodes: Soil amendments and microbial antagonists. *Plant and Soil* 100:237–247.
- Ruess, L. and W. Funke. 1992. Effects of experimental acidification on nematode populations in soil culture. *Pedobiologia* 36:231–239.
- Stephenson, W. 1945. The effects of acid on a soil nematode. *Parasitology* 36:158–164.
- Stirling, A. M., G. R. Stirling, and I. C. McCrae. 1992. Microbial degradation of fenamiphos after repeated application to a tomato-growing soil. *Nematologica* 38:245–254.
- Tarjan, A. C. 1956. Nematicidal value of some fatty acids. Bulletin 332. Rhode Island Agricultural Experiment Station. University of Rhode Island, Kingston, RI.
- Van Veen, J. A., J. N. Ladd, and M. Amato. 1985. Turnover of carbon and nitrogen through the microbial biomass in a sandy loam and clay soil incubated with [ $^{14}\text{C}(\text{U})$ ] glucose and [ $^{15}\text{N}$ ]  $(\text{NH}_4)_2 \text{SO}_4$  under different moisture regimes. *Soil Biology and Biochemistry* 17:747–756.
- Vance, E. D., P. C. Brookes, and D. S. Jenkinson. 1987. An extraction method for measuring soil microbial biomass C. *Soil Biology and Biochemistry* 19:703–707.