

Errata

Correction to Volume 39(2), page 119.

The legend to Figure 1 should read:

Fig. 1. Left: Map of continental Portugal and location of the affected area. Right: Location of *Bursaphelenchus xylophilus* isolates obtained from different blocks within the affected area. Black: the area where PWN is present; dark grey: the buffer area, established for safety reasons (free of PWN).

Correction to Volume 38(4), page 415.

Seven rows were omitted from the bottom of Table 1. This table is reproduced in its entirety below:

TABLE 1. Measurements (in μm) of females ($n = 15$) and males ($n = 14$) of *Stegelletina coprophila* sp. n.

Locality	Sima de la Fuente Negra (Villanueva del Arzobispo)			Sima de los Murciélagos (Hornos)	
Province	Jaén			Jaén	
Habitat	Bat guano			Organic material	
n	Holotype ♀	Paratypes 9♀♀	Paratypes 10♂♂	5♀♀	4♂♂
Body length	473	488.3 ± 41.7 (412–536)	487.4 ± 28.8 (425–521)	409.8 ± 21.9 (386–433)	427.0 ± 27.6 (391–458)
a	19.7	20.5 ± 1.9 (18.2–24.4)	20.9 ± 1.2 (19.3–23.3)	18.5 ± 1.2 (17.3–19.7)	20.6 ± 1.0 (19.6–21.6)
b	4.3	4.4 ± 0.2 (4.0–4.7)	4.5 ± 0.2 (4.1–4.7)	3.6 ± 0.1 (3.5–3.8)	3.8 ± 0.1 (3.7–4.0)
c	15.3	16.2 ± 0.8 (14.7–17.2)	15.4 ± 1.4 (12.1–17.1)	15.1 ± 0.9 (14.2–16.1)	14.7 ± 1.3 (13.5–16.6)
c'	1.8	2.0 ± 0.2 (1.7–2.3)	1.6 ± 0.1 (1.5–1.8)	2.0 ± 0.2 (1.8–2.2)	1.6 ± 0.2 (1.4–1.8)
V/T	64	64.3 ± 1.3 (62–66)	—	65.1 ± 0.5 (65–66)	—
Labial probolae	4	4.7 ± 0.7 (4–6)	4.2 ± 0.4 (4–5)	5.3 ± 0.5 (5–6)	5.3 ± 0.5 (5–6)
Lip region width	10	9.9 ± 0.8 (8–11)	10.0 ± 0.0 (10–10)	8.8 ± 2.5 (5–10)	9.8 ± 0.5 (9–10)
Stoma	9	8.3 ± 1.0 (7–10)	8.2 ± 0.8 (7–10)	9.5 ± 0.6 (9–10)	9.0 ± 0.8 (8–10)
Pharyngeal corpus	54	57.1 ± 3.7 (52–64)	55.7 ± 3.0 (51–62)	53.0 ± 1.2 (52–54)	54.0 ± 2.2 (51–56)
Isthmus	22	24.8 ± 3.2 (20–30)	24.0 ± 1.6 (22–26)	28.8 ± 3.6 (26–34)	26.8 ± 1.0 (26–28)
Bulb	18	17.7 ± 0.9 (16–19)	16.4 ± 1.0 (15–18)	17.0 ± 1.2 (16–18)	18.3 ± 1.9 (17–21)
Pharynx length	109	111.3 ± 7.8 (103–127)	108.4 ± 3.1 (103–112)	112.5 ± 3.8 (107–115)	111.0 ± 7.3 (103–119)
Nerve ring—ant. end	82	78.9 ± 5.4 (71–86)	77.7 ± 4.0 (71–86)	76.0 ± 4.3 (70–80)	80.5 ± 5.7 (76–88)
Excretory pore—ant. end	88	93.0 ± 6.3 (82–102)	93.2 ± 4.9 (82–98)	83.0 ± 6.1 (77–90)	88.0 ± 4.4 (83–91)
Deirid—ant. end	100	107.4 ± 5.2 (103–114)	107.3 ± 4.7 (100–114)	96.8 ± 7.4 (90–105)	102.0 ± 9.5 (96–113)
Annuli width	2	2.1 ± 0.2 (2–3)	2.1 ± 0.2 (2–3)	2.1 ± 0.2 (2–2)	1.7 ± 0.2 (2–2)
Cuticle thickness	1	1.0 ± 0.0 (1)	1.0 ± 0.0 (1)	1.4 ± 0.3 (1–2)	1.4 ± 0.3 (1–2)
Body width					
Neck base	23	22.3 ± 1.7 (19–25)	22.2 ± 0.9 (21–24)	20.0 ± 1.4 (18–21)	20.3 ± 1.5 (19–22)
Midbody	24	24.0 ± 2.5 (19–27)	23.4 ± 1.1 (22–25)	22.3 ± 1.7 (20–24)	20.8 ± 1.5 (20–23)
Anus	17	15.2 ± 1.2 (13–17)	19.1 ± 0.6 (18–23)	13.5 ± 1.3 (12–15)	18.5 ± 0.6 (18–19)
Lateral field	5	4.6 ± 1.1 (3–6)	5.5 ± 0.5 (5–6)	4.8 ± 0.5 (4–5)	4.0 ± 1.0 (3–5)
Vagina	10	9.1 ± 1.9 (7–12)	—	9.5 ± 1.3 (8–11)	—
Ovary/testis	213	118.5 ± 24.3 (88–154)	72.2 ± 16.5 (46–104)	91.0 ± 7.0 (86–99)	61.5 ± 6.0 (56–70)

TABLE 1. *Continued*

Locality	Sima de la Fuente Negra (Villanueva del Arzobispo)			Sima de los Murciélagos (Hornos)	
Province	Jaén			Jaén	
Habitat	Bat guano			Organic material	
n	Holotype ♀	Paratypes 9♀♀	Paratypes 10♂♂	5♀♀	4♂♂
Spermatheca	35	29.8 ± 8.1 (18–43)	—	29.5 ± 4.9 (26–33)	—
Uterus	60	60.2 ± 9.3 (48–71)	245.6 ± 18.2 (212–270)	46.5 ± 10.6 (39–54)	193.0 ± 9.3 (179–198)
Postvulvar sac	18	20.6 ± 7.8 (10–36)	—	15.0 ± 3.6 (12–19)	—
Rectum	18	16.0 ± 1.6 (14–18)	8.2 ± 1.3 (6–10)	14.09 ± 1.4 (13–16)	8.7 ± 1.5 (7–10)
Tail	31	30.2 ± 2.5 (26–34)	31.8 ± 2.4 (28–35)	27.3 ± 2.5 (24–30)	29.3 ± 3.6 (26–34)
Vulva-anterior end	303	314.0–26.9 (264–345)	—	267.0 ± 16.4 (249–285)	—
Phasmid-anus distance	9	10.5 ± 1.9 (8–13)	11.6 ± 1.9 (10–15)	9.7 ± 0.6 (9–10)	11.8 ± 1.7 (10–14)
Vulva-anus/tail	4.5	4.8 ± 0.3 (4.3–5.2)	—	4.3 ± 0.4 (3.9–4.7)	—
Spicules	—	—	24.0 ± 0.9 (22–25)	—	20.8 ± 1.0 (20–22)
Gubernaculum	—	—	12.0 ± 0.9 (10–13)	—	11.0 ± 0.8 (10–12)