INSECTA MUNDI

A Journal of World Insect Systematics

0042

Preliminary Checklist of the Mecoptera of Florida: Earwigflies, Hangingflies, and Scorpionflies

> Louis A. Somma Department of Botany and Zoology PO Box 118525 220 Bartram Hall, University of Florida Gainesville, FL 32611-8525

James C. Dunford Department of Entomology and Nematology PO Box 110620, IFAS Natural Area Drive, University of Florida Gainesville, FL 32611-0620

Date of Issue: August 22, 2008

Louis A. Somma and James C. Dunford Preliminary Checklist of the Mecoptera of Florida: Earwigflies, Hangingflies, and Scorpionflies Insecta Mundi 0042: 1-9

Published in 2008 by

Center for Systematic Entomology, Inc. P. O. Box 147100 Gainesville, FL 32614-7100 U. S. A. http://www.centerforsystematicentomology.org/

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod taxon. Manuscripts considered for publication include, but are not limited to, systematic or taxonomic studies, revisions, nomenclatural changes, faunal studies, book reviews, phylogenetic analyses, biological or behavioral studies, etc. **Insecta Mundi** is widely distributed, and referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc.

As of 2007, **Insecta Mundi** is published irregularly throughout the year, not as quarterly issues. As manuscripts are completed they are published and given an individual number. Manuscripts must be peer reviewed prior to submission, after which they are again reviewed by the editorial board to insure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Managing editor: Paul E. Skelley, e-mail: insectamundi@gmail.com Production editor: Michael C. Thomas, e-mail: thomasm@doacs.state.fl.us Editorial board: J. H. Frank, M. J. Paulsen

Printed copies deposited in libraries of:

CSIRO, Canberra, ACT, Australia Museu de Zoologia, São Paulo, Brazil Agriculture and Agrifood Canada, Ottawa, Ontario, Canada The Natural History Museum, London, England Muzeum I Instytut Zoologii Pan, Warsaw, Poland National Taiwan University, Taipei, Taiwan California Academy of Sciences, San Francisco, CA, USA Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA Field Museum of Natural History, Chicago, IL, USA National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Electronic copies in PDF format:

Printed CD mailed to all members at end of year. Florida Center for Library Automation: purl.fcla.edu/fcla/insectamundi University of Nebraska-Lincoln, Digital Commons: http://digitalcommons.unl.edu/insectamundi/

Author instructions available on the Insecta Mundi page at: http://www.centerforsystematicentomology.org/insectamundi/

Printed Copy	ISSN 0749-6737
On-Line	ISSN 1942-1354
CD-ROM	ISSN 1942-1362

Preliminary Checklist of the Mecoptera of Florida: Earwigflies, Hangingflies, and Scorpionflies

Louis A. Somma^{1,2} Department of Botany and Zoology PO Box 118525 220 Bartram Hall, University of Florida Gainesville, FL 32611-8525 somma@ufl.edu, sommal@doacs.state.fl.us

James C. Dunford^{1,2} Department of Entomology and Nematology PO Box 110620, IFAS Natural Area Drive, University of Florida Gainesville, FL 32611-0620 dunford@ufl.edu

Abstract. We provide the first species checklist of Mecoptera indigenous to Florida, based upon preliminary data gathered primarily from specimens housed in the Florida State Collection of Arthropods. There are 11 described and one undescribed species of mecopterans, representing three families, inhabiting the state of Florida. These include the recently discovered meropeid (earwigfly), *Merope tuber* Newman, four species of bittacids (hangingflies), represented by the genus *Bittacus* Latreille, and 7 species (one undescribed) of panorpids (scorpionflies), represented by the genus *Panorpa* Linnaeus. We are not certain if one of these, *Bittacus texanus* Banks, is indigenous, represents a nonindigenous occurrence, or is simply erroneous. Two of the species on our list, *Bittacus stigmaterus* Say and *Panorpa venosa* Westwood, represent first state records and one species, *Panorpa floridana* Byers, is endemic. Six of the species which have been recorded in Florida, *M. tuber*, *Bittacus punctiger* Westwood, *Panorpa rufa* Gray, *Panorpa pachymera* Byers, *Panorpa lugubris* Swederus and *P. venosa*, represent the southernmost records for these species in the continental United States. Perceived diversity and abundance of mecopteran species in Florida are limited by climate, biogeography, and collection bias.

Introduction

Mecoptera is a small, ancestral order of holometabolous insects. At present there are more than 600 known, extant species arranged in 34 genera and 9 families worldwide (Enderlein 1910; Esben-Petersen 1915, 1921; Tillyard 1918, 1935; Lameere 1936; Grassé 1951; Brues et al. 1954; Willmann 1987, 1989; Grimaldi and Engel 2005; Palmer et al. 2007; Penny 2008). Five families are widely distributed throughout much of the world, including North America, while the other four are restricted to South America and the Australian region (Kaltenbach 1978; Byers 2005; Dunford and Somma [in press]). Two families, Panorpidae and Bittacidae, comprise most of the species, and in North America they are represented by more than 54 species (1 genus) of panorpids and 10 species (4 genera) of bittacids (Hine 1898; Byers 2005; Cheung et al. 2006; Dunford and Somma [in press]). Fossil Mecoptera are well represented in sedimentary rocks of the lower Permian geological period, approximately 270 million years ago (Novokschonov 1998, 2002; Grimaldi and Engel 2005). There are nearly 400 known fossil species in approximately 87 genera and as many as 34 families (although researchers disagree on the exact number of extinct families), exhibiting a greater diversity than extant mecopteran taxa (Grimaldi and Engel 2005). Traditional "Mecoptera" lack defining, uniquely derived, unifying characters, and may not represent a monophyletic grouping. A combination of recent DNA and older morphological data suggests Mecoptera are closely

¹Florida State Collection of Arthropods, Division of Plant Industry - Entomology Section, Florida Department of Agriculture and Consumer Services, PO Box 147100, 1911 SW 34th Street, Gainesville, FL 32614-7100

 $^2\rm McGuire$ Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, PO Box 112710, University of Florida, Gainesville, FL 32611-2710

related to siphonapterans (Richards 1965; Richards and Richards 1969; Rothschild 1975; Whiting 2002; Grimaldi and Engel 2005; Marshall 2006).

Mecopterans have life histories as diverse as their morphologies. Many species inhabit mesic, forested environments and feed on decaying vegetation or dead, dying, or living soft bodied arthropods as larvae and/or adults; some groups are reported to feed on nectar, pollen, carrion, mosses, or are kleptoparasites of spider webs (Setty 1940; Byers 1963; Webb et al. 1975; Byers and Thornhill 1983; Bockwinkel and Sauer 1993; Palmer and Yeates 2005; Marshal 2006; Dunford and Somma [in press]). Larvae are generally eruciform or caterpillar-like, having a well developed head capsule with chewing mouthparts, and pupae are exarate and decticous, and typically located in soil (Setty 1940; Byers 1963, 1987; Boese 1973); however, these stages remain undiscovered in many mecopterans, including the meropeids (Byers 1987; Abbott et al. 2007; Dunford et al. 2007a, b, [in press]). Since many mecopterans inhabit moist forests, any disturbance reducing moisture levels (i.e., deforestation, global warming) could adversely affect or extirpate those species which are present. In general, Mecoptera are weak fliers, and dispersal could be severely limited if suitable habitat is reduced. This order is of little economic significance in North America but represents a systematically and biologically important, ancient insect lineage.

Materials and Methods

The Mecoptera collection at the Florida State Collection of Arthropods (FSCA) contains 20 drawers of mostly pinned adults, representing more than 4100 specimens (Dunford et al. 2007b). Herein, we present a brief summary of the 302 adult and 7 larval specimens, representing three families of Mecoptera, collected from Florida and housed within FSCA. We have augmented this information with literature records, and additional specimen data from mecopterans gifted to the National Museum of Natural History, Smithsonian Institution (NMNH), from FSCA in 2007. We only present detailed collection data for species representing state and county records of particular significance. We produced all of our illustrations from specimens housed in FSCA. Our overall goal is to provide initial information in order to stimulate further detailed studies on mecopterans inhabiting Florida.

Meropeidae-Earwigflies

The North American earwigfly, *Merope tuber* Newman, was recently (1996) collected in the Florida panhandle, representing first records for the state and southernmost records for North America (Dunford et al. 2007a). All 40 of these specimens, from Leon and Liberty counties, are currently housed in FSCA (Table 1; Dunford et al. 2007a, [in preparation]). This secretive meropeid is indigenous to eastern deciduous forests of North America. Its only other extant confamiliar is *Austromerope poultoni* Killington, an earwigfly restricted to extreme southwestern Western Australia (Byers 1973a; Abbott et al. 2007; Dunford et al. 2007a, [in press]; Somma and Dunford 2007).

Bittacidae-Hangingflies

Four species of bittacids, all in the genus *Bittacus* Latreille, have been collected in Florida. Two of these, *Bittacus pilicornis* Westwood and *Bittacus punctiger* Westwood, are somewhat common and previously recorded from the state (Carpenter 1935; Byers 1954, 1973b; Webb et al. 1975). *Bittacus pilicornis*, the hairy-horned hangingfly, is found in northern Florida, as far west as Jackson County in the eastern panhandle and as far south as Alachua County; while *B. punctiger* (Fig. 1), the spotted hangingfly, is found statewide (Table 1). The two specimens of *B. punctiger* from Miami-Dade County (Fuch's Hammock, near Homestead, 11-12 May 1980, one male collected by T. S. Dickel and H. V. Weems, Jr.; and Everglades National Park, 5 May 1961, one female collected by F. W. Mead), and perhaps the single specimen from an undescribed locality in Monroe County (25 June 1964, one male collected by K. J. Stone), represent the southernmost records for this species in the continental United States. Say's hangingfly, *Bittacus stigmaterus* Say, is only represented by a single specimen from the Florida panhandle, Liberty County (Torreya State Park, 20 August 1987, one male collected by C. Porter and L. A. Stange), and two other specimens from the northern peninsula, Alachua County (Gainesville, 1 October 1940, one individual missing an abdomen collected by T. Hubbell; and unspecified locality, 16 October



Figure 1-4. 1) Habitus of male *Bittacus punctiger*, spotted hangingfly, from Alachua County, Florida. Image by James C. Dunford. **2)** Habitus of male *Panorpa lugubris*, black scorpionfly, from unknown locality, Florida. Image by Gary J. Steck. **3)** Habitus of male *Panorpa floridana* holotype, Florida scorpionfly, from Alachua County, Florida. Image by James C. Dunford. **4)** Habitus of male *Panorpa americana*, American scorpionfly, from Liberty County, Florida. Image by James C. Dunford. **5**

1951, one female collected by "G. E. N.") (Table 1). These are first state records for *B. stigmaterus* in Florida. A single specimen of *Bittacus texanus* Banks, the Texas scorpionfly, was collected in Citrus Center west of Lake Okeechobee, Glades County, Florida (Carpenter 1931; Byers 1993), but no specimens exist in FSCA. As this record is greatly disjunct from the more westerly range of this species (Carpenter 1931; Webb et al. 1975), and no other records for Florida exist of which we are aware, we are not clear whether *B. texanus* is indigenous to the state or if this represents a single nonindigenous record. Byers (1993) suspects that the label locality data for this specimen are erroneous.

Panorpidae - Scorpionflies

Six described and one undescribed species of panorpids, represented by the genus *Panorpa* Linnaeus, occur in Florida. Five of these, *Panorpa lugubris* Swederus, *Panorpa rufa* Gray, *Panorpa floridana*

Byers, *Panorpa pachymera* Byers and *Panorpa americana* Swederus, were previously recorded from the state. *Panorpa lugubris*, the black scorpionfly (Fig. 2), is clearly the most widespread and abundant scorpionfly indigenous to Florida (Glover in Hagen 1861; Hine 1901a, b; Carpenter 1931, 1935; Byers 1954, 1993). Specimens of this species in FSCA represent 15 counties that include the panhandle, northern peninsular and portions of central peninsular Florida, as far south as Orange and Pinellas counties (Table 1). A larval *Panorpa* collected from Glen St. Mary, Baker County, Florida, on 27 January 1961 by E. W. Holder was putatively assigned to *P. lugubris* by G. W. Byers (Table 1). Additionally, a map provided by Byers (1993) includes Escambia, Jackson, and Manatee counties; the lattermost being the southernmost record for *P. lugubris* in Florida and the continental United States. Lastly, on 29 March 2007, FSCA gifted 24 pinned adult *P. lugubris* to NMNH that were collected from Leon (N = 10), Baker (N = 7), and Alachua (N = 7) counties, Florida.

Panorpa rufa, the red scorpionfly, is indigenous to northern Florida. Byers (1954, 1993) and FSCA specimens (Table 1) limit this species to the panhandle in addition to disjunct records from Alachua County in the northern peninsula. The Alachua County records are the southernmost in the United States. FSCA gifted 2 pinned adult specimens from Gulf and Okaloosa counties, Florida, respectively, to NMNH on 29 March 2007. In addition to the 5 counties listed in Table 1, Byers (1954, 1993) includes Bay and Holmes counties in the Florida panhandle.

Panorpa floridana, the Florida scorpionfly, is endemic to Florida and is only known from Alachua and Clay counties (Byers 1993; Table 1). The two specimens in FSCA are the holotype from Alachua County, Florida (near the San Felasco Hammock, Gainesville, 10 November 1970, one male collected by L. A. Hetrick; Fig. 3), and a paratype from Clay County, Florida (Orange Park, 25 December 1936, one male collected by K. Wheeler) (Byers 1993). Additionally, Byers (1993) lists an allotype from Clay County, Florida (Gold Head Branch State Park, 20 November 1982, one female collected by S. W. Gross); and two more paratypes from Florida (Alachua County, 8 November 1974, one female collected by D. Richman; and Gold Head Branch State Park, Clay County, 20 November 1982, one male collected by S. W. Gross).

Byers (1993) tentatively assigned a *Panorpa* specimen from Santa Rosa County, in the Florida panhandle, to *P. pachymera*, the stout-genitaled scorpionfly. Additionally, Carpenter's (1931) record for *Panorpa rufescens* Rambur, the rufous scorpionfly, from Madison County, northern Florida, before this species group was split by Byers (1993), potentially could be referable to *P. pachymera*. Other than these two presumptive records, there exist two *P. pachymera* from Gainesville, Alachua County, Florida (4 December 1918, one male and one female collected by G. B. Merrill) and four from Leon County, Florida (Tall Timbers Research Station, 6-7 May 1968, one male and two females collected by H. V. Weems Jr.; and 6 May 1968, one male collected by G. H. Heinrich). These FSCA specimens verify the existence of *P. pachymera* in Florida (Table 1). The Alachua County specimens are the southernmost records from the United States.

Panorpa americana (Fig. 4), the American scorpionfly, is limited to the Florida panhandle (Carpenter 1931; Byers 1954, 1993; Table 1). Additionally, 12 pinned adult *P. americana* from Okaloosa County, Florida, were gifted from FSCA to NMNH on 29 March 2007.

We report the first state records for *Panorpa venosa* Westwood, the veined scorpionfly, in Florida; they include Baker (Glen St. Mary, 17-28 November 1960, two males and two females collected by H. V. Weems Jr.) and Leon counties (Tall Timbers Research Station, 4-17 May 1996, one female collected by P. W. Kovarik) (Table 1). These specimens are the southernmost records for this species.

Several specimens of *Panorpa* from Florida in FSCA are not assigned to species (Table 1). These include six larval *Panorpa* from Glen St. Mary, Baker County, Florida, collected on 14 January 1961 by E. W. Holder. Additionally, a single adult female from the J. & E. Turf Farm, north of LaCrosse, in the Santa Fe River floodplain, Alachua County, Florida, collected 19-26 July 1986 by G. J. Steck and B. D. Sutton, only can be assigned to the *Panorpa virginica* Banks, Virginia scorpionfly, species group (Table 1). Lastly, an undescribed species of *Panorpa* is represented by a poorly preserved male in alcohol, collected on 3 August 1948 from an unspecified locality in Putnam County, Florida (Table 1). There are not enough discernable characters preserved on this headless specimen to allow it to be described, but it is a member of the *P. virginica* species group (G. W. Byers pers. comm. 2007).

Discussion

Current records for Mecoptera in Florida, representing only 11 described and one undescribed species, indicate a low diversity for the state. Two species, *B. stigmaterus* and *P. venosa*, represent first state records. One species, *P. floridana*, is endemic, and six others, *M. tuber*, *B. punctiger*, *P. rufa*, *P. pachymera*, *P. lugubris* and *P. venosa*, are the southernmost records for these species in the continental United States. The single record for *B. texanus* in Glades County, in southern peninsular Florida (Carpenter 1931), could be erroneous (Byers 1993) or represent a nonindigenous occurrence. However, Byers (1954, 1993) makes note of a disjunct population of *B. texanus* in Kansas, more than 800 km north of its previously known range. This suggests that further collecting should be done in Glades County, Florida, and the Lake Okeechobee area in order to verify this record.

Three factors may limit mecopteran distribution and diversity in Florida, as perceived in our preliminary list. One factor may be the lack of suitable habitat, especially in the southernmost portions of the state, where warmer temperatures might be unfavorable to most species of North American mecopterans. This climatological restriction might be further enhanced by the Pleistocene biogeography (i. e., marine flooding and xerification) of Florida, especially the peninsular portions of the state (Neill 1957; Dunford et al. 2007a). Lastly, few naturalists study and collect extant mecopterans (Byers 1990; Palmer et al. 2007). This collection bias might partially explain why there is an apparent dearth of records for mecopterans in central and southern Florida, including widespread species such as *B. punctiger* and *P. lugubris*, while larger numbers and more diverse species representation come from heavily collected areas such as Leon, Okaloosa, and Alachua counties (Table 1). Alachua County, in particular the city of Gainesville, has a high density of entomologists and other naturalists, and it is not surprising that it is the type locality for Florida's only endemic mecopteran, *P. floridana*. This species has been recorded from only two counties to date, and we suspect that further intensive collecting may reveal a somewhat broader distribution for *P. floridana* in low elevation regions of northern peninsular Florida.

Our preliminary data presented herein are fragmentary and preclude generating adequate summaries of distribution, abundance, and phenologies for Floridian mecopterans. Although climatological and biogeographical limitations likely account for the seeming paucity of species and overall lower abundance of mecopterans at these southerly latitudes, further dedicated collecting may reveal a previously unrecognized diversity and allow for the generation of complete ecological, phenological and biogeographical data for this insect order in Florida. We opine that new records and undoubtedly new species of Mecoptera await discovery in the state, and strongly encourage individuals and institutional collections to publish records of mecopterans collected from Florida.

Acknowledgments

We are grateful to G. B. Edwards (Curator of Arachnida and Myriapoda, Florida State Collection of Arthropods) for initially inquiring about the number of species of scorpionflies indigenous to Florida. Our manuscript was enormously improved by critical reviews provided by George W. Byers (Snow Entomological Museum, Biodiversity Institute, The University of Kansas) and Norman D. Penny (Department of Entomology, California Academy of Sciences); both whom we acknowledge with much gratitude. We thank Gary J. Steck (Curator of Diptera and Minor Orders, FSCA) for the image of *P. lugubris*. We are indebted to Janice Kahler (Interlibrary Loan Department, Library West, George A. Smathers Libraries, University of Florida) for obtaining obscure literature for us. James R. "M.O.B." Wiley (Assistant Curator, FSCA), Wm. Charles Whitehill (Assistant Curator, FSCA), and Christopher M. Palmer (Biodiversity Conservation South, Parks and Wildlife Service, Australia) graciously assisted our literature search. Additionally, J. R. Wiley generously augmented the Mecoptera collection at FSCA. This is Entomology Contribution No. 1092, Bureau of Entomology, Nematology and Plant Pathology, Division of Plant Industry, Florida Department of Agriculture and Consumer Services, which also provided technical and library support for this research.

Literature Cited

- Abbott, I., T. Burbidge, and A. Wills. 2007. *Austromerope poultoni* (Insecta, Mecoptera) in southwest Western Australia: Occurrence, modelled geographical distribution, and phenology. Journal of the Royal Society of Western Australia 90: 97-106.
- Bockwinkel, G., and K.-P. Sauer. 1993. *Panorpa* scorpionflies foraging in spider webs Kleptoparasitism at low risk. Bulletin of the British Arachnological Society 9: 110-112.
- Boese, A. E. 1973. Descriptions of larvae and key to fourth instars of North American *Panorpa* (Mecoptera: Panorpidae). The University of Kansas Science Bulletin 50: 163-186.
- Brues, C. T., A. L. Melander, and F. M. Carpenter. 1954. Classification of insects. Bulletin of the Museum of Comparative Zoology at Harvard College 108: 1-917.
- Byers, G. W. 1954. Notes on North American Mecoptera. Annals of the Entomological Society of America 47: 484-510.
- Byers, G. W. 1963. The life history of *Panorpa nuptialis* (Mecoptera: Panorpidae). Annals of the Entomological Society of America 56: 142-149.
- Byers, G. W. 1973a. Zoogeography of the Meropeidae (Mecoptera). Journal of the Kansas Entomological Society 46: 511-516.
- Byers, G. W. 1973b. Descriptions and distributional records of American Mecoptera. III. Journal of the Kansas Entomological Society 46: 362-375.
- Byers, G. W. 1987. Order Mecoptera. p. 246-252. *In:* F. W. Stehr (editor). Immature insects. [Volume 1.] Kendall/Hunt Publishing Company; Dubuque. 754 p.
- **Byers, G. W. 1990.** Order Mecoptera. p. 177-178. *In*: M. Kosztarab and C. W. Schaefer (editors). Systematics of the North American insects and arachnids: Status and needs. Virginia Agricultural Experiment Station Information Service 90-1. Virginia Polytechnic Institute and State University; Blacksburg. 247 p.
- Byers, G. W. 1993. Autumnal Mecoptera of southeastern United States. University of Kansas Science Bulletin 55: 57-96.
- Byers, G. W. 2005 [2004]. Order Mecoptera. Scorpionflies and hangingflies. p. 662-668. *In:* C. A. Triplehorn and N. F. Johnson (editors). Borror and DeLong's introduction to the study of insects. Seventh edition. Thomson Brooks/Cole; Belmont, CA. 864 p.
- Byers, G. W., and R. Thornhill. 1983. Biology of the Mecoptera. Annual Review of Entomology 28: 203-228.
- Carpenter, F. M. 1931. Revision of the Nearctic Mecoptera. Bulletin of the Museum of Comparative Zoölogy at Harvard College 72: 205-277, pl. 1-8.
- Carpenter, F. M. 1935. New Nearctic Mecoptera, with notes on other species. Psyche 42: 105-122.
- Cheung, D. K. B., S. A. Marshall, and D. W. Webb. 2006. Mecoptera of Ontario. Canadian Journal of Arthropod Identification (1): 1-13.
- Dunford, J. C., P. W. Kovarik, L. A. Somma, and D. Serrano. 2007a. First state records for *Merope tuber* (Mecoptera: Meropeidae) in Florida and biogeographical implications. Florida Entomologist 90: 581-584.
- **Dunford, J. C., and L. A. Somma. [in press].** Scorpionflies (Mecoptera). *In*: J. L. Capinera (editor). Encyclopedia of entomology. Second edition. Springer; Dordrecht.
- Dunford, J. C., L. A. Somma, and D. Serrano. 2007b. FSCA Mecoptera resources: Mecoptera (scorpionflies, hangingflies, earwigflies, and allies) at the Florida State Collection of Arthropods [online]. Available on URL: http://www.fsca-dpi.org/MecopteraPage.htm. In: M. C. Thomas (editor). Museum of Entomology FSCA (Gainesville). Florida State Collection of Arthropods, Division of Plant Industry, Florida Department of Agriculture and Consumer Services; Gainesville. (Last accessed 20 May 2008.)
- Dunford, J. C., L. A. Somma, and D. Serrano. [in press]. Earwigflies (Mecoptera: Meropeidae). In: J. L. Capinera (editor). Encyclopedia of entomology. Second edition. Springer; Dordrecht.
- **Dunford, J. C., L. A. Somma, and D. Serrano. [in preparation].** Current disposition and holdings for earwigflies, *Merope tuber* Newman and *Austromerope poultoni* Killington (Mecoptera: Meropeidae), in the Florida State Collection of Arthropods.
- **Enderlein, G. 1910.** Über die Phylogenie und Klassifikation der Mecopteren unter Berücksichtigung der fossilen Formen. Zoologischer Anzeiger 35: 385-399.
- Esben-Petersen, P. 1915. A synonymic list of the order Mecoptera together with descriptions of new species. Entomologiske Meddelelser (København) 10: 216-242.

- Esben-Petersen, P. 1921. Mecoptera. Monographic revision. Collections Zoologiques du Baron Edm. de Selys Longchamps; Catalogue Systématique et Descriptif, Publié par les Soins de ses Fils (Bruxelles), Fascicule 5(2): 1-172, pl. I-II.
- Grassé, P.-P. 1951. Ordre des Mécoptères. p. 71-124. *In:* P.-P. Grassé (editor). Traité de zoologie. Anatomie, systématique, biologie. Tome X. Insectes superiéurs et Hémiptéroïdes. Fascicule I. Névroptéroïdes. Mécoptéroïdes Hyménoptéroïdes (Symphytes et térébrants). Masson et C^{ie} Editeurs; Paris. 975 p.
- Grimaldi, D. [A.], and M. S. Engel. 2005. Evolution of the insects. Cambridge University Press; Cambridge. 772 p.
- **Hagen, H. 1861.** Synopsis of the Neuroptera of North America. With a list of South American species. Smithsonian Miscellaneous Collections 4(1): i-xx, 1-347.
- Hine, J. S. 1898. The North American species of the genus *Bittacus*. Proceedings of the Columbus Horticultural Society 13: 105-115, pl. I-II.
- Hine, J. S. 1901a. A review of the Panorpidae north of Mexico. Bulletin of the Science Laboratories of Denison University 11: 241-264, pl. LIX-LX.
- Hine, J. S. 1901b. A review of the Panorpidae north of Mexico. University Bulletin. Ohio State University, Series V (7): 241-264, pl. LIX-LX.
- Kaltenbach, A. 1978. Mecoptera (Schnabelhafte, Schnabelfliegen). Handbuch der Zoologie 4(2); (2/28): 1-111.
- Lameere, A. 1936. Mécoptères. Précis de Zoologie, Publications de l'Institut Zoologique Torley-Rousseau (Bruxelles) 5: 5-16.
- Marshall, S. A. 2006. Insects. Their natural history and diversity. With a photographic guide to the insects of eastern North America. Firefly Books Ltd; Richmond Hill, Ontario. 718 p.
- Neill, W. T. 1957. Historical biogeography of present-day Florida. Bulletin of the Florida State Museum, Biological Sciences 2: 175-220.
- Novokschonov, V. [G.] 1998. Some problems of scorpionfly (Mecoptera) evolution [in Russian Cyrillic]. Zoologicheskii Zhurnal 77: 677-688.
- Novokschonov, V. G. 2002. Order Panorpida Latreille, 1802. The scorpionflies. p. 194-199. *In*: A. P. Rasnitsyn and D. L. J. Quicke (editors). History of insects. Kluwer Academic Publishers; Dordrecht. 517 p.
- Palmer, C. M., J. W. H. Trueman, and D. K. Yeates. 2007. Systematics of the Apteropanorpidae (Insecta: Mecoptera) based on morphological and molecular evidence. Invertebrate Systematics 21: 589-612.
- Palmer, C. M., and D. K. Yeates. 2005. Diet and feeding behavior in adults of the Apteropanorpidae (Mecoptera). Journal of Insect Behavior 18: 209-231.
- Penny, N. D. 2008. Mecoptera [online]. Available on URL: http://research.calacademy.org/research/entomology/Entomology_Resources/mecoptera/index.html. *In*: California Academy of Sciences Entomology. California Academy of Sciences; San Francisco. (Last accessed 20 May 2008.)
- Richards, A. G. 1965. The proventriculus of adult Mecoptera and Siphonaptera. Entomological News 76: 253-256.
- Richards, P. A., and A. G. Richards. 1969. Acanthae: A new type of cuticular process in the proventriculus of Mecoptera and Siphonaptera. Zoologische Jahrbücher, Abteilung für Anatomie und Ontogenie der Tiere 86: 158-176
- Rothschild, M. 1975. Recent advances in our knowledge of the Order Siphonaptera. Annual Review of Entomology 20: 241-259.
- Setty, L. R. 1940. Biology and morphology of some North American Bittacidae (Order Mecoptera). American Midland Naturalist 23: 257-353.
- Somma, L. A., and J. C. Dunford. 2007. Etymology of the earwigfly, *Merope tuber* Newman (Mecoptera: Meropeidae): Simply dull or just inscrutable? Insecta Mundi 0013: 1-5.
- **Tillyard, R. J. 1918.** The panorpoid complex. A study of the phylogeny of the holometabolous insects, with special reference to the subclasses Panorpoidea and Neuropteroidea. [Introduction]. Proceedings of the Linnean Society of New South Wales 43: 265-284.
- Tillyard, R. J. 1935. The evolution of the scorpion-flies and their derivatives (Order Mecoptera). Annals of the Entomological Society of America 28: 1-45.
- Webb, D. W., N. D. Penny, and J. C. Marlin. 1975. The Mecoptera, or scorpionflies, of Illinois. Illinois Natural History Survey Bulletin 31: 250 -316.

Whiting, M. F. 2002. Mecoptera is paraphyletic: Multiple genes and phylogeny of Mecoptera and Siphonaptera. Zoologica Scripta 31: 93-104.

Willmann, R. 1987. The phylogenetic system of the Mecoptera. Systematic Entomology 12: 519-524.

Willmann, R. 1989. Evolution und phylogenetisches System der Mecoptera (Insecta: Holometabola). Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft (544): 1-153.

Received May 21, 2008; Accepted July 8, 2008.

MECOPTERA OF FLORIDA

TABLE 1. Species of Floridian Mecoptera, listed by family, based exclusively upon specimens in the Florida State Collection of Arthropods current 20 May 2008. Florida counties from which each species was collected are listed beneath each, with number of specimens from each county provided in parentheses. Previously unpublished state records are indicated with a "*".

Meropeidae

Merope tuber Newman 1838 Leon (34), Liberty (6)

Bittacidae

Bittacus pilicornis Westwood 1846 Alachua (11), Clay (2), Jackson (1), Suwannee (3), Wakulla (1)

Bittacus punctiger Westwood 1846 Alachua (40), Collier (4), Highlands (2), Jefferson (1), Manatee (1), Miami-Dade (2), Monroe (1), Pinellas (1), Suwannee (2)

Bittacus stigmaterus Say 1823* Alachua (2), Liberty (1)

Panorpidae

```
Panorpa americana Swederus 1787
Gulf (1), Leon (1), Liberty (2), Okaloosa (26)
```

Panorpa floridana Byers 1993 Alachua (1), Clay (1)

```
Panorpa lugubris Swederus 1787
Alachua (17), Baker (30), Bradford (1), Columbia (2), Duval (2), Lake (1), Leon (63), Levy (1), Marion (2),
Okaloosa (1), Orange (6), Pinellas (1), St Johns (2), Suwannee (3), Wakulla (1), unspecified locality (1)
```

Panorpa pachymera Byers 1993 Alachua (2), Leon (4)

```
Panorpa rufa Gray 1832
Alachua (2), Gulf (1), Okaloosa (2), Wakulla (1), Walton (2)
```

- Panorpa venosa Westwood 1832* Baker (4), Leon (1)
- Panorpa sp., undescribed species¹ Putnam (1)

Panorpa sp.² Alachua (1)

Panorpa larvae Baker (7)³

²Not possible to assign to a species but in the *P. virginica* species group (determined by G. W. Byers 2002). ³Six undetermined specimens and one individual putatively assigned to *P. lugubris* (determined by G. W. Byers 1963).

¹Determined by G. W. Byers 1977. An undescribed member of the *P. virginica* species group (G. W. Byers pers. comm. 2007).