

# INSECTA MUNDI

A Journal of World Insect Systematics

---

0070

First record of the genus *Parahyliota* in the New World  
(Coleoptera: Silvanidae: Brontinae)

Michael C. Thomas  
Florida State Collection of Arthropods  
Florida Department of Agriculture and Consumer Services  
P.O. Box 1047100  
Gainesville, FL 32614-7100

Date of Issue: April 10, 2009

Michael C. Thomas  
First record of the genus *Parahyliota* in the New World  
(Coleoptera: Silvanidae: Brontinae)  
Insecta Mundi 0070: 1-4

**Published in 2009 by**

Center for Systematic Entomology, Inc.  
P. O. Box 141874  
Gainesville, FL 32614-1874 U. S. A.  
<http://www.centerforsystematicentomology.org/>

**Insecta Mundi** is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod taxon. Manuscripts considered for publication include, but are not limited to, systematic or taxonomic studies, revisions, nomenclatural changes, faunal studies, book reviews, phylogenetic analyses, biological or behavioral studies, etc. **Insecta Mundi** is widely distributed, and referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc.

As of 2007, **Insecta Mundi** is published irregularly throughout the year, not as quarterly issues. As manuscripts are completed they are published and given an individual number. Manuscripts must be peer reviewed prior to submission, after which they are again reviewed by the editorial board to insure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

**Managing editor:** Paul E. Skelley, e-mail: [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Production editor:** Michael C. Thomas, e-mail: [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Editorial board:** J. H. Frank, M. J. Paulsen

**Subject editors:** J. Eger, A. Rasmussen, F. Shockley, G. Steck, A. Van Pelt, J. Zaspel

**Printed copies deposited in libraries of:**

CSIRO, Canberra, ACT, Australia  
Museu de Zoologia, São Paulo, Brazil  
Agriculture and Agrifood Canada, Ottawa, Ontario, Canada  
The Natural History Museum, London, England  
Muzeum i Instytut Zoologii Pan, Warsaw, Poland  
National Taiwan University, Taipei, Taiwan  
California Academy of Sciences, San Francisco, CA, USA  
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA  
Field Museum of Natural History, Chicago, IL, USA  
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

**Electronic copies in PDF format:**

Printed CD mailed to all members at end of year.

Florida Center for Library Automation: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

**Author instructions** available on the Insecta Mundi page at:

<http://www.centerforsystematicentomology.org/insectamundi/>

|              |                |
|--------------|----------------|
| Printed Copy | ISSN 0749-6737 |
| On-Line      | ISSN 1942-1354 |
| CD-ROM       | ISSN 1942-1362 |

First record of the genus *Parahyliota* in the New World  
(Coleoptera: Silvanidae: Brontinae)

Michael C. Thomas

Florida State Collection of Arthropods  
Florida Department of Agriculture and Consumer Services  
P.O. Box 1047100  
Gainesville, FL 32614-7100

**Abstract.** The genus *Parahyliota* Thomas, previously known from Africa and Asia, is recorded from the New World for the first time, based on *P. balli* Thomas, a new species here described from Mexico.

**Introduction**

The genus *Parahyliota* Thomas (2004) was recently described to accommodate *Uleiota costicollis* Reitter and 12 other Asian and African species previously assigned to *Uleiota* Latreille. This paper records the presence of *Parahyliota* in the New World for the first time based on a new species from Mexico. The new species was discovered in a shipment of undetermined silvanids from the University of Alberta collection.

The new species possesses all of the diagnostic characters of the genus: absence of a scutellary striole, tarsal structure, modified male frons, and reduced parameres (Thomas 2004). This species represents only the seventh member of the tribe Brontini in the New World; the tribe is much more diverse in tropical Asia and Australia (Thomas 2004).

Collection codens used are: **CNCI**, Canadian National Collection of Insects, Arachnids and Nematodes, Ottawa, Canada; **FSCA**, Florida State Collection of Arthropods, Gainesville, FL, USA; **UASM**, E.H. Strickland Entomological Museum, University of Alberta, Edmonton, AB, Canada; **UNAM**, Universidad Nacional Autónoma de México, Mexico, Mexico D.F., Mexico; **USNM**, National Museum of Natural History, Washington, D.C., USA.

***Parahyliota balli* Thomas, new species**

Fig. 1-8

**Diagnosis.** This species is the only member of the tribe Brontini known to occur in the Neotropics. The genus is included in a key to world brontine genera in Thomas (2004).

**Description.** Holotype, male (Fig. 1-3), deposited in USNM, with following label data: "MEX. Oaxaca 13.4 km. s. Totontepec cloud forest 2530m. in litter 79-37 June 16, 1979/MEXICAN EXP. 1979 J. S. Ashe, G. E. Ball & D. Shpeley collectors".

**Body.** With the characters of *Parahyliota* Thomas as described by Thomas (2004), plus: Length, 6.2 mm; elongate, parallel-sided, dorso-ventrally compressed; dark testaceous in color (Fig. 1).

**Head.** Transverse, 1.4x wider than long; excavate over antennal insertions, with a shallow transverse impression anterior to antennal insertions and a deep transverse impression posterior to eyes. Mandibles broad, angulate, each with a vertically-directed, long curved tooth arising from the dorsal surface. Antennae elongate, filiform, longer than body; scape slightly sinuate, very elongate, as long as next three antennomeres, attaining base of pronotum; pedicel short. (Holotype missing most antennal flagellomeres. Proportions of antennomere lengths of male paratype are: 5.25:1.00:2.00:2.25:2.25:2.50:2.50:2.50:2.25:2.00:2.00; surpassing length of body by about three antennomeres.) Dorsal surface densely punctate, punctures about equal in size to an eye facet, separated by less than one puncture diameter, each puncture subtending a long pale seta; impunctate above antennal insertions; surface between punctures microreticulate laterally, smooth and shining medially; a tear shaped fovea bordered anteriorly with a blunt tubercle is present on each side of frons within antennal insertion. Eyes large, not convex, temple narrow posteriorly, about 2 eye facets in width (Fig. 3).



**Figure 1-3.** *Parahyliota balli* Thomas, new species, male holotype. 1) Dorsal habitus. 2) Head and pronotum. 3) Front of head and mandibles.

**Pronotum.** Transverse, 1.3x wider than longer, widest across apical angles; rectangular in shape, lateral margins slightly converging posteriorly; disk depressed; margins denticulate, anterior angle right, with a distinct tooth beyond anterior angle, posterior angle obtuse (Fig. 2); surface sculpture and pubescence as on head.

**Elytra.** Elongate, 2x longer than combined width, parallel-sided, converging conjointly from about apical third to suture; laterally narrowly costate, the lateral margin narrowly explanate; disc with six rows of punctate striae, striae not impressed; a single row of smaller punctures between each stria; each puncture subtending a long suberect pale seta.



**Figure 4-5.** *Parahyliota balli* Thomas, new species, female allotype. 4) Dorsal habitus. 5) Head and pronotum.



**Figure 6-8.** *Parahyliota balli* Thomas, new species. Male genitalia. **6)** Aedeagus. **7)** Tegmen. **8)** Claspers, composed of abdominal segments VIII and IX.

**Genitalia.** Aedeagus as is Fig. 6; parameres highly reduced (Fig. 7), abdominal segments VIII and IX forming claspers as in Fig. 8.

**Female.** Allotype (Fig. 4-5) with same data as holotype, deposited in USNM. Similar to male, differing as follows: Length, 6.4 mm. Head, 1.8x wider than long; pronotum, 1.3x wider than long; elytra, 2.1x longer than combined width; color castaneous; punctuation and pubescence as in male; mandibles without dorsal tooth and frons without blunt tubercles (Fig. 5); antennae about as long as body, scape attaining middle of pronotum; ratios of antennomere lengths: 5.0:1.0:2.0:2.3:2.3:2.3:2.0:2.0:1.6:2.0.

**Paratypes.** Seven with label data as follows: 2, same data as holotype; 5, "MEX. Oaxaca 16.6 km. s. Totontepec cloud forest 2540 m in litter 79-35 June 16, 1979/MEXICAN EXP. 1979 J. S. Ashe, G. E. Ball & D. Shpeley collectors". Deposited in CNCI, FSCA, UNAM, and UASM.

**Variation.** Paratypes range in color from testaceous to castaneous; the testaceous specimens appear general. Male paratype is very similar to holotype, length 5.8 mm. Female paratypes range in length from 5.7 mm to 5.8 mm.

**Etymology.** I take great pleasure in naming this species in honor of George Ball, pre-eminent carabidologist and friend, who was one of the collectors of the type series.

**Discussion.** George Ball (in litt.) wrote that his field notes indicate that the specimens were actually collected under bark and that the "in litter" referred to carabids collected at the same localities. At both localities specimens of a species of the silvanid genus *Aplatamus* Sharp were collected. Individuals of that genus seem to be infrequently collected.

The male of *P. balli* possesses mandibular horns, which otherwise occur only in the African and Madagascan members of *Parahyliota*. None of the Asian species is known to possess mandibular horns.

### Acknowledgments

I thank George Ball for the loan of specimens and Andrew Cline and Paul Skelley for manuscript review. This is Entomology Contribution No. 1136 of the Bureau of Entomology, Nematology, and Plant Pathology, Florida Department of Agriculture and Consumer Services.

### Literature Cited

**Thomas, M. C. 2004.** The Brontini of the world: A generic review of the tribe (Coleoptera: Silvanidae: Brontinae). *Insecta Mundi* 17: 1-31 (2003).

**Received February 18, 2009; Accepted March 18, 2009.**

**Subject Edited by F. Shockley.**