

INSECTA MUNDI

A Journal of World Insect Systematics

0245

First recorded introduction of the milliped order Stemmiulida
(Eugnatha: Nematophora): Potential establishment in Florida, USA,
and new records from Mexico; northward range extension into southern
Tamaulipas

Rowland M. Shelley
Research Laboratory
North Carolina State Museum of Natural Sciences
MSC #1626
Raleigh, NC 27699-1626 USA

G. B. Edwards
Florida State Collection of Arthropods
Division of Plant Industry
P. O. Box 147100
Gainesville, FL 32614-7100 USA

Arthur E. Bogan
Research Laboratory
North Carolina State Museum of Natural Sciences
MSC #1626
Raleigh, NC 27699-1626 USA

Date of Issue: August 24, 2012

Rowland M. Shelley, G. B. Edwards, and Arthur E. Bogan
First recorded introduction of the milliped order Stemmiulida (Eugnatha:
Nematophora): Potential establishment in Florida, USA, and new records from
Mexico; northward range extension into southern Tamaulipas
Insecta Mundi 0245: 1-5

Published in 2012 by

Center for Systematic Entomology, Inc.
P. O. Box 141874
Gainesville, FL 32614-1874 USA
<http://www.centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology. Manuscript preparation guidelines are available at the CSE website.

Managing editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Production editor: Michael C. Thomas, Brian Armitage, Ian Stocks

Editorial board: J. H. Frank, M. J. Paulsen

Subject editors: G.B. Edwards, J. Eger, A. Rasmussen, F. Shockley, G. Steck, Ian Stocks, A. Van Pelt, J. Zaspel

Spanish editors: Julieta Brambila, Angélico Asenjo

Printed copies (ISSN 0749-6737) deposited in libraries of:

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and Agrifood Canada, Ottawa, ON, Canada
The Natural History Museum, London, Great Britain
Muzeum i Instytut Zoologiczny PAN, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD mailed to all members at end of year.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://edocs.ub.uni-frankfurt.de/volltexte/2010/14363/>

Author instructions available on the *Insecta Mundi* page at:

<http://www.centerforsystematicentomology.org/insectamundi/>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

**First recorded introduction of the milliped order Stemmiulida
(Eugnatha: Nematophora): Potential establishment in Florida, USA,
and new records from Mexico; northward range extension into southern
Tamaulipas**

Rowland M. Shelley
Research Laboratory
North Carolina State Museum of Natural Sciences
MSC #1626
Raleigh, NC 27699-1626 USA
rowland.shelley@ncdenr.gov

G. B. Edwards
Florida State Collection of Arthropods
Division of Plant Industry
P. O. Box 147100
Gainesville, FL 32614-7100 USA
GB.Edwards@freshfromflorida.com

Arthur E. Bogan
Research Laboratory
North Carolina State Museum of Natural Sciences
MSC #1626
Raleigh, NC 27699-1626 USA
arthur.bogan@ncdenr.gov

Abstract. Based on two “uni-ocellate” females, the world’s first introductions of the milliped order Stemmiulida are recorded from Florida, United States (US). One individual was collected in 1976 in Gainesville, Alachua County (Co.), in northcentral peninsular Florida, and the other was taken in 1991 some 408 km (255 mi) to the south-southeast in Pompano Beach, Broward Co. The absence of further individuals and additional samples suggests that the introductions did not result in viable populations, and stemmiulidans are not presently established in the state; the Gainesville site was reinvestigated in 2012 without finding additional specimens. New records from Mexico include the first from Chiapas, Oaxaca, Tabasco, Yucatan, San Luis Potosí, and Tamaulipas states, with the northernmost ordinal locality now becoming Rancho del Cielo, northwest of Gómez Fariás, in the last. A northward range expansion of about 460 km (288 mi) from the previous limit, Xalapa, Veracruz, the site lies a mere 40 km (25 mi) south of the Tropic of Cancer and only some 320 km (200 mi) south of the Rio Grande and the US border at McAllen, Hidalgo Co., Texas. Indigenous Stemmiulida are not expected in the forested Rio Grande Valley of southernmost Texas, but their occurrence in the adjoining Mexican state renders such a discovery more plausible than before.

Keywords. Alachua Co., Broward Co., ocellus/ocellate, Rio Grande Valley, Tamaulipas, Tropic of Cancer, Veracruz

Introduction

The monofamilial diplopod order Stemmiulida (superorder Nematophora) is perhaps the most distinctive in the chilognath subterclass Eugnatha, as except for a couple of anophthalmous species, it is unfailingly diagnosed by the presence of one or two pairs of large ocelli caudal to the antennae (Fig. 1-4). Stemmiulida inhabit tropical America, Africa, and India/Sri Lanka, and are also known from Vietnam, Indonesia (Halmahera), and the Island of New Guinea (Hoffman 1977, 1980, 1982, 1999; Mauriès 1985, 1989; Mauriès and Golovatch 2006; Mauriès et al 2010; Shelley and Golovatch 2011). In the New World, the order ranges from Cuba and central Veracruz (Xalapa), Mexico, to southcentral Peru (Huanuco Dept.) and Brazil south of the Amazon River (Loomis 1968, Hoffman 1999, Bueno-Villegas et al. 2004, Shelley and Golovatch 2011). It is unknown north of the Tropic of Cancer, has not been encountered in

the Bahamas, and is absent from Florida, the continental United States (US/USA), and North America (Chamberlin and Hoffman 1958, Hoffman 1999, Shelley 2001, Shelley and Golovatch 2011).

Stemmiulida are also one of only two eugnathan orders for which at least potential humanly introduced populations are unknown, the other being Siphoniulida, assigned to Nematophora by Shear (2011). Copious publications, too many to cite here, document exogenous, allochthonous species in the orders Polydesmida (Merocheta) and Julida and Spirobolida (Juliformia), and known and likely ones exist in Cambalidea and Spirostreptidea (Juliformia: Spirostreptida *s. l.*) (Enghoff 1982, 1993; Mauriès 1983; Shelley 1998; Golovatch et al. 2007, 2011) and Chordeumatida (Shelley 1990, 2002; Hoffman 1999; Shear and Shelley 2004) and Callipodida (Nematophora) (Demange 1946; Korsós 1992, 1994; Korsós et al. 2002; Stoev and Enghoff 2006, Stoev et al. (2008), Shelley and Golovatch (2011). While recently perusing samples in the FSCA (codens below), RMS discovered two stemmiulidan samples collected 21 and 36 years ago in peninsular Florida with one “uni-ocellate” female each. Both are assignable to *Stemmiulus* Gervais, 1844, in today’s taxonomy (Mauriès and Golovatch 2006, Shear 2011), but without males, we can neither determine species nor verify whether female only samples are conspecific. The localities are approximately 408 km (255 mi) apart, so the specimens can only be interpreted as two accidental human importations, and the paucity of individuals, coupled with the absence of more recent Floridian samples, suggests that viable populations have not become established. However, the fact that these stemmiulidans got into Florida and were collected alive shows that importation of this order is possible, so we put them on record in case additional representatives are discovered in the state. We also present photos of the Broward County (Co.) female, depigmented after 21 years in preservative, along with a more recently collected, partly pigmented, “doubly-ocellate” stemmiulidan from Puerto Rico. Sample data are as follows:

FLORIDA: *Alachua Co.*, Gainesville, SW 23rd St., Honey Plant facilities, “*Cynodon dactylon* (L.)” [Bermuda Grass], F, 7 May 1976, W. E. Wyles. *Broward Co.*, Pompano Beach, “*Heliconia* sp.”, F, 13 March 1991, E. Tannehill.

Seven nominal genera were once recognized in Stemmiulida (Hoffman 1980), but Mauriès and Golovatch (2006) reduced this to just two – *Scoliogmus* Loomis, 1941, monotypic for *Scoliogmus teres* Loomis, 1941, in Puerto Rico, and *Stemmiulus* Gervais, 1844, for the remaining 153 nominal species. The other five generic names – *Diopsiulus* Silvestri, 1897; *Prostemmiulus* Silvestri, 1916; *Paurochaeturus* Silvestri, 1916; *Plusiochaeturus* Silvestri, 1916; and *Nesoiulus* Brolemann, 1920 – were placed in synonymies under *Stemmiulus* and their species were formally transferred into this genus. Mauriès et al. (2010) accepted this arrangement and erected *Eostemmiulus* for the first Vietnamese representative, but Shear (2011) called for further argumentation before the concept of only two genera could be fully accepted. Stemmiulida have generally received little attention from diplopodologists, and decades will likely pass before isolated females like these can be authoritatively determined, but recording even unidentifiable females from Florida is desirable to further document the state’s known and potential arthropod fauna.

With the FSCA being in Gainesville, the Alachua Co. locality was reinvestigated by GBE in June 2012. The area has changed considerably since the original collection, with additional buildings, a student coop. organic garden, and a densely overgrown protected natural area where none existed previously. The periphery of this area was examined using a rake to turn over debris on the ground and lift up compost piles, but no millipeds were found.

We also cite the following stemmiulidan samples from Mexico, which add six states to the ordinal occurrence; codens are **AMNH**, American Museum of Natural History, New York; **FSCA**, Florida State Collection of Arthropods, Gainesville; and **UCDC**, Bohart Entomological Museum, University of California at Davis. Of particular note are those from Tamaulipas, the northeasternmost state; indigenous New World occurrence now lies a mere 40 km (25 mi) south of the Tropic of Cancer. Specifically, Rancho del Cielo, northwest of Gómez Farias in southern Tamaulipas and some 460 km (288 mi) north-northwest of Xalapa, Veracruz, is now the northernmost locality in the Western Hemisphere (Fig. 5). Only 320 km (200 mi) south of the Rio Grande and the US border at McAllen, Hidalgo Co., Texas, it suggests that Stemmiulida may inhabit the Rio Grande Valley region of that state. As riparian forests on the US side in Cameron, Hidalgo, and Starr cos. have been reasonably well sampled without discovery of stemmiulidans, we believe indigenous US occurrence is unlikely, but since the order definitely inhabits the adjoining Mexican state, such a discovery is more plausible than before.



Figures 1-4. 1) Broward Co., Florida, stemmiulidan, depigmented after 20 years in preservative. 2) The same, close up of left ocellus. 3) *Stemmium heatwolei* (Vélez, 1967) from Arecibo, Puerto Rico. 4) the same, close up of right ocellus.

MEXICO: *Campeche*: 2.4 km (1.5 mi) SW Bolonchen, Cueva de Bolonchen, 31 December 1971, D. McKenzie (FSCA); 5 km (3.1 mi) SW Bolonchenticul, Grutas de Xtacumbilxunan, 19 April - 30 June 1973, 25 May 1974, B. Denson, M. Kawakatsu, R.W. Mitchell, J.R. Reddell, S. Wiley, D. McKenzie (FSCA); and 10 km (6.3 mi) W Hopelden, 27 July 1973, J.R. Reddell, J.W. Rowland (FSCA). *Chiapas*, 8 km (5 mi) NE Chiapas, 22 August 1966, J. and W. Ivie (AMNH) **New State Record**. *Oaxaca*, Temescal, 18 July 1960, J.S. Buckett, M.R. and R.C. Gardner (UCDC); and Vega del Sol, Valle Nacional, 14 July 1966, J.S. Buckett, M.R. and R.C. Gardner (UCDC) **New State Record**. *San Luis Potosí*, Talmazunchale, 24 June 1954, N.B. Causey (FSCA) **New State Record**. *Tabasco*, Teapa, Grutas del Cocona, 25 August 1972, R.W. Mitchell, W.H. Russell (FSCA) **New State Record**. *Tamaulipas*, 9.6 km (6 mi) NW Gómez Farias, Rancho del Cielo, 2 June 1967 and 9 and 25 March 1969, J.R. Reddell, F. Howell, S. Fowler, C. Tucker, R.W. Mitchell (FSCA) **Northernmost record from the New World**; 7 km (4.4 mi) S Gómez Farias, 18 February 1973, T.R. Mollhagen (FSCA); and 0.5 km (0.3 mi) NNW San Rafael de los Castrol, August 1969, W.H. Russell (FSCA) **New State Record**. *Yucatan*, Uxmal ruins, 6 August 1973, R.W. and D.R. Mitchell (FSCA); Chichen Itza, 8 August 1973, J.R. Reddell (FSCA); Mayapan Ruins, 14 August 1973, J.R. Reddell (FSCA) **New State Record**.

Acknowledgments

We thank L. Prendini and L.S. Kimsey for access to, and loans from, the AMNH and UCDC, respectively. Presubmission reviews were provided by S.I. Golovatch and the late R.L. Hoffman (1927-2012).

Literature Cited

- Bueno-Villegas, J., P. Sierwald, and J. E. Bond. 2004.** Diplopoda. p. 569-599. In: B. J. Llorente, J. J. Morrone, O. Yáñez, y I. Vargas I (eds.). Biodiversidad, Taxonomía y Biogeografía de Artrópodos de México: Hacia una síntesis de su conocimiento. IV. UNAM; México. 660 p.
- Chamberlin, R. V., and R. L. Hoffman. 1958.** Checklist of the millipeds of North America. United States National Museum Bulletin 212: 1-236.
- Demange, J.-M. 1946.** *Callipus foetidissimus gallicus denticulatus* nov. var. des catacombes du Muséum d'Histoire Naturelle de Paris (Myriapodes Diplopodes). Bulletin du Muséum National d'Histoire Naturelle, 2 sér., 18(5): 394-396.
- Enghoff, H. 1982.** Millipedes (Diplopoda) from the Cape Verde Islands. Courier Forschungsinstitut Senckenberg 52: 137-138.
- Enghoff, H. 1993.** Cape Verdean millipedes (Diplopoda). Tropical Zoology 6: 207-216.
- Golovatch, S. I., J.-J. Geoffroy, J.-P. Mauriès, and D. VandenSpiegel. 2007.** Review of the millipede genus *Glyphiulus* Gervais, 1847, with descriptions of new species from Southeast Asia (Diplopoda, Spirostreptida, Cambalopsidae). Part 2: the *javanicus*-group. Zoosystema 29(3): 417-455.
- Golovatch, S. I., J.-J. Geoffroy, J.-P. Mauriès, and D. VandenSpiegel. 2011.** New species of the millipede genus *Glyphiulus* Gervais, 1847 from the *granulatus*-group (Diplopoda: Spirostreptida: Cambalopsidae). Arthropoda Selecta 20(2): 65-114.
- Hoffman, R. L. 1977.** A stemmiulid milliped from Central Africa. Revue Suisse de Zoologie 84(4): 831-835.
- Hoffman, R. L. 1980 (1979).** Classification of the Diplopoda. Muséum d'Histoire Naturelle; Genève, Switzerland. 237 p.
- Hoffman, R. L. 1982.** Diplopoda. p. 689-724. In: S. B. Parker (ed.). Synopsis and Classification of Living Organisms 2: 1-1119.
- Hoffman, R. L. 1999.** Checklist of the millipedes of North and Middle America. Virginia Museum of Natural History Special Publication Number 8: 1-584.
- Korsós, Z. 1992.** Millipedes from anthropogenic habitats in Hungary. Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck, Supplementum 10: 237-241
- Korsós, Z., 1994.** Checklist, preliminary distribution maps, and bibliography of millipedes in Hungary (Diplopoda). Miscellanea Zoologica Hungarica 9: 29-82.
- Korsós, Z., E. Hornung, K. Szlávecz, and J. Kontschán. 2002.** Isopoda and Diplopoda of urban habitats: new data to the fauna of Budapest. Annales Historico-Naturales Musei Nationalis Hungarici 94: 193-208.
- Loomis, H. F. 1968.** A checklist of the millipedes of Mexico and Central America. United States National Museum Bulletin 266: 1-137.
- Mauriès, J.-P. 1983.** Cambalides nouveaux et peu connus d'Asie, d'Amérique et d'Océanie, I. Cambalidae et Cambalopsidae (Myriapoda, Diplopoda). Bulletin du Muséum National d'Histoire Naturelle, 4e série, section A 5: 247-276.
- Mauriès, J.-P. 1985.** Contributions to the knowledge of the mountain fauna of Malawi (Mission R. Jocqué). 5. Diplopoda Stemmiulida. Revue de Zoologie Africaine 99: 41-146.




Figure 5. Floridian localities of Stemmiulida in relation to the indigenous range in the Western Hemisphere, adjusted northward in eastern Mexico to account for occurrence in southern Tamaulipas. The depicted boundary appears to run along the Tropic of Cancer in the Gulf of Mexico, but it crosses into Mexico just south of this meridian.

- Mauriès, J.-P. 1989.** Révision des stemmiulides: espèces nouvelles et peu connues d'Afrique (Myriapoda, Diplopoda). Bulletin du Muséum National d'Histoire Naturelle, 4e série, section A, Zoologie 11(3): 605-637.
- Mauriès, J.-P., and S. I. Golovatch. 2006.** *Stemmiulus deharvengi* sp. n., the first Stemmiulida to be reported from Indonesia (Diplopoda: Stemmiulida). Arthropoda Selecta 15(2): 91-98.
- Mauriès, J.-P., S. I. Golovatch, and J.-J. Geoffroy. 2010.** A new genus and species of the order Stemmiulida from Vietnam (Diplopoda). Arthropoda Selecta 19(2): 73-80.
- Shear, W. A. 2011.** Class Diplopoda de Blainville in Gervais, 1844. p. 159-164, *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 1-237.
- Shear, W. A., and R. M. Shelley. 2004.** Introduction of the millipede *Cleidogona nantahala* Shear, 1972, in New England (Chordeumatida: Cleidogonidae). Entomological News, 115(2): 88-94.
- Shelley, R. M. 1990.** Discovery of the millipede *Craspedosoma rawlinsii* Leach in the New World (Chordeumatida: Craspedosomatidae). Entomologists Monthly Magazine 126: 117-118.
- Shelley, R. M. 1998.** Occurrence of the millipede *Glyphiulus granulatus* (Gervais) in the Hawaiian Islands (Spirostreptida: Cambalidea: Cambalopsidae). Bishop Museum Occasional Papers, 56: 36-37.
- Shelley, R. M. 2001 (2000).** Annotated checklist of the millipeds of Florida (Arthropoda). Insecta Mundi 14(4): 241-251.
- Shelley, R. M. 2002.** The millipeds of central Canada (Arthropoda: Diplopoda), with reviews of the Canadian fauna and diplopod faunistic studies. Canadian Journal of Zoology 80: 1863-1875.
- Shelley, R. M. 2003 (2002).** A revised, annotated, family-level classification of the Diplopoda. Arthropoda Selecta 11(3): 187-207.
- Shelley, R. M., and S. I. Golovatch. 2011.** Atlas of myriapod biogeography. I. Indigenous ordinal and supra-ordinal distributions in the Diplopoda: Perspectives on taxon origins and ages, and a hypothesis on the origin and early evolution of the class. Insecta Mundi 0158: 1-134.
- Stoev, P., and H. Enghoff. 2006.** A review of the millipede genus *Dorypetalum* Verhoeff, 1900 (Diplopoda: Callipodida: Dorypetalidae). Zootaxa 1254: 29-43.
- Stoev, P., P. Sierwald, and A. Billey. 2008.** An annotated world catalogue of the millipede order Callipodida (Arthropoda: Diplopoda). Zootaxa 1706: 1-50.

Received July 1, 2012; Accepted August 4, 2012.

