

INSECTA MUNDI

A Journal of World Insect Systematics

0742

Three new synonymies in *Phyllophaga* Harris, 1827
(Coleoptera: Scarabaeidae), with lectotype
and neotype designations

Kyle E. Schnepf
Florida State Collection of Arthropods
Division of Plant Industry, Florida Department of Agriculture and Consumer Services
1911 SW 34th Street
Gainesville, FL 32608, USA

Date of issue: December 23, 2019

Kyle E. Schnepf

Three new synonymies in *Phyllophaga* Harris, 1827 (Coleoptera: Scarabaeidae), with lectotype and neotype designations

Insecta Mundi 0742: 1–6

ZooBank Registered: urn:lsid:zoobank.org:pub:37A2F301-0637-41B1-8CE7-A17A54564AF9

Published in 2019 by

Center for Systematic Entomology, Inc.

P.O. Box 141874

Gainesville, FL 32614-1874 USA

<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

Chief Editor: David Plotkin, insectamundi@gmail.com

Assistant Editor: Paul E. Skelley, insectamundi@gmail.com

Head Layout Editor: Robert G. Forsyth

Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas

Review Editors: Listed on the *Insecta Mundi* webpage

Printed copies (ISSN 0749-6737) annually deposited in libraries

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agrifood Canada, Ottawa, ON, Canada

The Natural History Museum, London, UK

Muzeum i Instytut Zoologii PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Robert G. Forsyth

Three new synonymies in *Phyllophaga* Harris, 1827 (Coleoptera: Scarabaeidae), with lectotype and neotype designations

Kyle E. Schnepf

Florida State Collection of Arthropods

Division of Plant Industry, Florida Department of Agriculture and Consumer Services

1911 SW 34th Street

Gainesville, FL 32608, USA

keschnepf@gmail.com

Abstract. In the course of working on new species of North American *Phyllophaga* Harris, 1827 (Coleoptera: Scarabaeidae: Melolonthinae) some synonyms have been found and are proposed here. **New synonymies:** *Phyllophaga knausii* (Schaeffer, 1907) is synonymized with *Phyllophaga sociata* (Horn, 1878); *Phyllophaga chippewa* Saylor, 1939 is synonymized with *Phyllophaga rugosa* (Melsheimer, 1845); and *Phyllophaga falta* Sanderson, 1950 is synonymized with *Phyllophaga bipartita* (Horn, 1887). Lectotypes are here designated for the following species: *Listrochelus knausii* Schaeffer, *Listrochelus sociatus* Horn, and *Lachnosterna bipartita* Horn. A neotype for *Ancylonycha rugosa* Melsheimer is here designated from the Horn Collection.

Key words. Melolonthinae, Melolonthini, North America, taxonomy.

Introduction

Phyllophaga Harris, 1827 (Coleoptera: Scarabaeidae: Melolonthinae) is a diverse New World genus of over 850 valid species, with more than 200 of these occurring in the United States and Canada (Evans and Smith 2009). While the U.S. fauna is relatively well-known and identification guides are available (Chapin 1935; Saylor 1939a, 1940; Reinhard 1950a, 1950b; Luginbill and Painter 1953; Lago et al. 1979; Woodruff and Beck 1989; Warner and Morón 1992; Harpootlian 2001; Ratcliffe and Paulsen 2008), several species in the western U.S. remain undescribed. While examining type material for future descriptive work, some synonymies were discovered and are proposed here.

Materials and Methods

Label data. Label information is given verbatim in quotes, a slash+space (/) indicates line breaks and space+double slash+space (//) indicates a different label. Labels are typed on white paper unless otherwise indicated by information between brackets ([]).

Specimens examined. Types of all six species were studied, and additional specimens from the following collections were examined to determine variation:

FSCA	Florida State Collection of Arthropods, Gainesville, FL, USA
INHS	Illinois Natural History Survey, Champaign, IL, USA
MCZ	Museum of Comparative Zoology, Cambridge, MA, USA
USNM	United States National Museum of Natural History, Washington, D.C., USA

Specimens were examined with a Leica S6D microscope. Photographs were taken on a Leica Z16 APO microscope using a JVC KY-F75U digital camera and stacked with Synchrosopy Automontage software, version 5.01.005. Images were compiled into plates using GIMP 2 software (version 2.10.12).

Results and Discussion

Phyllophaga rugosa (Melsheimer, 1845)

Figures 3–4, 9–10

Ancylonycha rugosa Melsheimer 1845: 140. **Neotype:** MCZ, **here designated**, male.

Phyllophaga rugosa (Melsheimer): Haldeman and LeConte 1853: 59; Glasgow 1916: 371.

Lachnosterna rugosa (Melsheimer): LeConte 1856: 252.

Phyllophaga chippewa Saylor 1939b: 455. **New synonymy.** Type: USNM, male.

Ancylonycha rugosa was described by Melsheimer (1845) from Virginia. Haldeman and LeConte (1853) included it in *Phyllophaga* in their revision of Melsheimer's catalogue. LeConte (1856) used the generic name *Lachnosterna* for this species and most other *Phyllophaga* (sensu stricto). Saylor (1939b) described *Phyllophaga chippewa* from a single male collected in Schley, Minnesota (Fig. 10), and compared it with *Phyllophaga knochii* (Schoenherr and Gyllenhal). While *P. chippewa* does belong in Group IX of Horn's (1887) revision, as stated by Saylor (1939b), there are some significant differences between this species and *P. knochii*, including pronotal punctation and clypeal emargination. The parameres of *P. chippewa* are mostly developed, looking normally sclerotized, but the base of the genitalia (Fig. 4) is clearly underdeveloped and misshapen. The genitalia appear to be a deformity of *P. rugosa*, as stated by Luginbill and Painter (1953). After examination of the type, *P. chippewa* is externally within the variation of *P. rugosa* and the collection locality is also within the range of *P. rugosa* (Pike et al. 1977). Being the type of *P. chippewa* remains the only known specimen, I concur with Luginbill and Painter (1953) in considering it a deformed specimen of *P. rugosa* and here formally synonymize the two species.

A search for Melsheimer's type of *A. rugosa* turned up no specimens and is presumed lost. A neotype of *A. rugosa* is here designated to fix the name to a single specimen to allow comparison to other types. The specimen chosen is a dissected male (Fig. 3) from North Carolina in the Horn Collection. Label data (Fig. 9) are as follows: "N. C // ♂ // HornColl/ H [handwritten] 5774 // MCZ-ENT 00711292 // [on red] NEOTYPE/ *Ancylonycha rugosa*/ Melsheimer, 1845/ Det: K. E. Schnepf 2018".

Phyllophaga sociata (Horn, 1878)

Figures 1–2, 7–8

Listrochelus sociatus Horn 1878: 146. **Lectotype:** MCZ, **here designated**, male.

Phyllophaga sociatus (Horn): Saylor 1938: 131.

Listrochelus knausii Schaeffer 1907: 67. **New synonymy.** **Lectotype:** USNM, **here designated**, male.

Phyllophaga knausi (Schaeffer): Blackwelder and Arnett 1974: 58.

Horn (1878) described *Listrochelus sociatus* from "Nevada, Idaho and Oregon" without specific locality or type designation. In 1938, Saylor moved *L. sociatus* from *Listrochelus* to *Phyllophaga* without explanation. In 1907, Schaeffer described *Listrochelus knausii*, apparently without seeing the types of *L. sociatus*, stating "By description *knausii* is very close to *sociatus*, near which it has to be placed, but the latter species is said to have the posterior tibiae stout, resembling those of *Ligyris*." Schaeffer's types of *L. knausii*, a male and female, have been dissected, possibly by Schaeffer or Milton Sanderson. Both of Schaeffer's types of *L. knausii* are from Stockton, Utah and no holotype has been designated. To fix the name to a single specimen I am designating the male (Fig. 2) with the following data (Fig. 8) as the lectotype: "[handwritten label] Stockton/ V.14. Utah // [handwritten label] Warren/ Knaus // [handwritten label] Listrochelus/ knausii/ type ♂ Schae // [in red] Type [handwritten] ♂ // [handwritten label with two red border lines] Listrochelus/ knausii/ Schae. // [on red] Cotype/ No [handwritten] 42585/ U.S.N.M. // BROOKLYN/ MUSEUM/ COLL. 1929 // [on red] LECTOTYPE/ Listrochelus knausii/ Schaeffer, 1907/ Det: K. E. Schnepf 2018".

Similarly, no type has been designated for *Listrochelus sociatus* Horn. After examining the type series, a male lectotype (Fig. 1) is here designated to fix the name to a single specimen to allow comparison to other types. Label data (Fig. 7) are as follows: "Nev. // [on blue] PARA-TYPE/ [handwritten] 3635.2 // MCZ-ENT/ 00522248 // [on red] LECTOTYPE/ *Listrochelus sociatus*/ Horn, 1878/ Det: K. E. Schnepf 2018".

Sanderson's *Phyllophaga* notes are available for study at the FSCA. In them, he clearly recognized *P. knausii* as being synonymous with *P. sociata*, but no formal synonymy has been made. After studying the types, I concur with Sanderson and here formally synonymize *P. knausii* with *P. sociata*.

***Phyllophaga bipartita* (Horn, 1887)**

Figures 5–6, 11–12

Lachnosterna bipartita Horn 1887: 242. **Lectotype:** MCZ, **here designated**, male.*Phyllophaga bipartita* (Horn): Glasgow 1916: 373.*Phyllophaga falta* Sanderson 1950: 92. **New synonymy.** Type: INHS, male.

Lachnosterna bipartita was described by Horn (1887) from “Kansas, Louisiana and Texas.” Glasgow resurrected *Phyllophaga* in 1916 and included this species in the genus. *Phyllophaga falta* was described by Sanderson (1950) from a single male collected in Fayetteville, Arkansas (Fig. 12). Externally, *P. falta* is similar to *P. fervida* (Fabricius), as stated by Sanderson (1950), with the only notable difference being the origin of a ridge on the penultimate sternite of the male. This is, however, the same ridge found in some individuals of *P. bipartita*. There is a wide variation in development of this ridge, from virtually non-existent to heavily sclerotized and projecting to the apex of the segment. Externally, there are no differences between *P. falta* and some individuals of *P. bipartita*. There is considerable variation in the development of sclerotized parts of the male genitalia of *P. bipartita*, as stated by Luginbill and Painter (1953) and Ratcliffe and Paulsen (2008), including projections on the parameres as well as the internal sac. Southwestern populations of *P. bipartita* have the left paramere (when viewed with parameres face on, as in Fig. 5) lacking a strongly hamate structure, instead they look similar to the right paramere. Specimens from the central and eastern portion of this species range have an enlarged hook structure, as well as having a forward projecting process of varying size on the right paramere. The internal sac of *P. bipartita* is also quite variable, some have a pair of heavily sclerotized patches with a series of spinules on the apex, while others have a lightly sclerotized strip with no spinules where these patches would be. There are also two patches of short spines of varying development on the bottom of the sac at the base; some specimens have patches not unlike those in *P. falta*. These patches are occasionally strongly developed and each appear as a single large, dark, spinose section. There are also specimens that fall between these two extremes. *Phyllophaga falta* (Fig. 6) appears to be one of the many variations of *P. bipartita*, just with strongly reduced parameres. Since the type of *P. falta* falls within the range of external variation of *P. bipartita*, the two species are here synonymized.

After examining the type series of *L. bipartita*, a male lectotype (Fig. 5) is here designated to fix the name to a single specimen to allow comparison to other types. Label data (Fig. 11) are as follows: “Kan. // ♂ // [handwritten label] L./ bipartita/ Horn // MCZ-ENT/ 00008066 // Jan.–Jul. 2005/ MCZ Image/ Database // [on red] TYPE No. [handwritten] 3673/ [handwritten] Lachnosterna/ [handwritten] bipartita/ G. H. Horn // [on red] MCZ TYPE/ [handwritten] 8066 // [on red] LECTOTYPE/ *Lachnosterna bipartita*/ Horn, 1887/ Det: K. E. Schnepf 2018”.

Acknowledgments

I thank Crystal Maier (MCZ), Rachel Hawkins (MCZ), Floyd Shockley (USNM), and Chris Grinter (INHS) for loaning material in their care. I also thank Bill Warner and Paul Skelley (FSCA) for comments on the manuscript. Support for this work was provided by the Florida Department of Agriculture and Consumer Services, Division of Plant Industry.


Literature Cited

- Blackwelder, R. E., and R. H. Arnett, Jr. 1974.** Checklist of the beetles of Canada, United States, Mexico, Central America and the West Indies. North American Beetle Fauna Project (red version). Volume 1, part 3. The scarab beetles, ant-loving beetles, clown beetles and related groups. The Biological Research Institute of America; Latham, NY, USA. 120 p.
- Chapin, E. A. 1935.** Review of the genus *Chlaenobia* Blanchard (Coleoptera: Scarabaeidae). Smithsonian Miscellaneous Collections 94(9): 1–20.
- Evans, A. V., and A. B. T. Smith. 2009.** An Electronic Checklist of the New World Chafer (Coleoptera: Scarabaeidae: Melolonthinae). Version 3. Electronically published; Ottawa, Canada. 353 p. Available at <http://unsm-ento.unl.edu/SSSA/NW-Melo-v3.pdf> (Last accessed November 2019.)

- Glasgow, R. D. 1916.** *Phyllophaga* Harris (*Lachnosterna* Hope): A revision of the synonymy, and one new name. Bulletin of the Illinois State Laboratory of Natural History 11(5): 365–379.
- Haldeman, S. S., and J. L. LeConte. 1853.** Catalogue of the described Coleoptera of the United States by Friedrich Ernst Melsheimer, revised. Smithsonian Institution; Washington, DC. 174 p.
- Harpootlian, P. J. 2001.** Scarab beetles (Coleoptera: Scarabaeidae) of South Carolina. Biota of South Carolina. Volume 2. Clemson University; Clemson, SC. 157 p.
- Harris, T. W. 1827.** Minutes towards a history of some American species of Melolonthae particularly injurious to vegetation. Massachusetts Agricultural Journal 10(1): 1–12.
- Horn, G. H. 1878.** Revision of the species of *Listrochelus* of the United States. Transactions of the American Entomological Society 7: 137–148.
- Horn, G. H. 1887.** Revision of the species of *Lachnosterna* of America north of Mexico. Transactions of the American Entomological Society 14: 209–296.
- Lago, P. K., R. L. Post, and C. Y. Oseto. 1979.** The phytophagous Scarabaeidae and Troginae (Coleoptera) of North Dakota. North Dakota Insects Publication 12: 1–131.
- LeConte, J. L. 1856.** Synopsis of the Melolonthidae of the United States. Journal of the Academy of Natural Sciences of Philadelphia, Second Series 3: 225–288.
- Luginbill, P., and H. R. Painter. 1953.** May beetles of the United States and Canada. United States Department of Agriculture Technical Bulletin 1060: 1–102.
- Melsheimer, F. E. 1845.** Descriptions of new species of Coleoptera of the United States. Proceedings of the Academy of Natural Sciences of Philadelphia 2: 134–160.
- Pike, K. S., R. L. Rivers, and Z. B. Mayo. 1977.** Geographical distribution of the known *Phyllophaga* and *Cyclocephala* species in the North Central States. University of Nebraska Agricultural Experiment Station Miscellaneous Publication 34: 1–13.
- Ratcliffe, B. C., and M. J. Paulsen. 2008.** The scarabaeoid beetles of Nebraska. Bulletin of the University of Nebraska State Museum 22: 1–570.
- Reinhard, H. J. 1950a.** The *Phyllophaga* of Texas (Scarabaeidae, Coleoptera). Journal of the Kansas Entomological Society 23(1): 27–40.
- Reinhard, H. J. 1950b.** The *Phyllophaga* of Texas (Scarabaeidae, Coleoptera). Journal of the Kansas Entomological Society 23(2): 41–51.
- Sanderson, M. W. 1950.** New North American *Phyllophaga* (Scarabaeidae). Journal of the Kansas Entomological Society 23(3): 90–93.
- Saylor, L. W. 1938.** A new *Phyllophaga* from Nevada (Coleoptera: Scarabaeidae). Proceedings of the Entomological Society of Washington 40(5): 129–131.
- Saylor, L. W. 1939a.** Revision of the beetles of the melolonthine subgenus *Phytalus* of the United States. Proceedings of the United States National Museum 86(3048): 157–167.
- Saylor, L. W. 1939b.** Notes and descriptions of United States Scarab beetles. Journal of the Washington Academy of Sciences 29(10): 452–461.
- Saylor, L. W. 1940.** Revision of the scarabaeid beetles of the phyllophagan subgenus *Listrochelus* of the United States, with discussion of related subgenera. Proceedings of the United States National Museum 89(3095): 59–130.
- Schaeffer, C. F. A. 1907.** New Scarabaeidae. Journal of the New York Entomological Society 15(2): 60–75.
- Warner, W. B., and M. A. Morón. 1992.** A revision of the *Phyllophaga* subgenus *Triodonyx* Saylor (Coleoptera: Scarabaeidae). Journal of the Kansas Entomological Society 65(3): 321–340.
- Woodruff, R. E., and B. M. Beck. 1989.** The scarab beetles of Florida (Coleoptera: Scarabaeidae). Part II. The May or June beetles (genus *Phyllophaga*). Arthropods of Florida and Neighboring Land Areas 13: 1–225.

Received November 26, 2019; accepted December 4, 2019.

Review editor David Plotkin.



Figures 1–6. Male genitalia of *Phyllophaga* type specimens. 1) *Phyllophaga sociata*, lectotype. 2) *Phyllophaga knausii*, lectotype. 3) *Phyllophaga rugosa*, neotype. 4) *Phyllophaga chippewa*, holotype. 5) *Phyllophaga bipartita*, lectotype. 6) *Phyllophaga falta*, holotype.



Figures 7–12. Specimen labels of *Phyllophaga* type specimens. 7) *Phyllophaga sociata*, lectotype. 8) *Phyllophaga knausii*, lectotype. 9) *Phyllophaga rugosa*, neotype. 10) *Phyllophaga chippewa*, holotype. 11) *Phyllophaga bipartita*, lectotype. 12) *Phyllophaga falta*, holotype.