

Nomenclatural changes, reinstatements, new combinations, and new synonymies among American Cerambycids (Coleoptera)

Gérard Luc Tavakilian

Entomologie Forestière

ORSTOM B.P. 165

97323 CAYENNE cedex, FRENCH GUIANA

Abstract: New synonyms: *Protorma scabrosa* Waterhouse, 1880 and *Protorma recurvatum* Williams, 1829 = *Strongylaspis costifer* Thomson, 1877; *Callopsisma ruficollis* Bates, 1870 = *Eriphus croceicollis* White, 1855; *Odontocera cinctiventris* Bates, 1870 = *Odontocera simplex* White, 1855; *Tomopterus aurantiacosignatus* Zajciw, 1969 = *Tomopterus similis* Fisher, 1930; *Cerambyx spectabilis* Voet, 1778 = *Cerambyx velutinus* Fabricius, 1775; *Chion ochraceus* Bates, 1885 = *Lamia spinifera* Fabricius, 1792; *Lissonotus shepherdi* Pascoe, 1859 = *Callidium equestre* Fabricius, 1787; *Cerambyx pulverulentus* Olivier, 1790 = *Cerambyx farinosus* Linné, 1758; *Lamia crypta* Say, 1832 = *Saperda annulata* Fabricius, 1801; *Ptericoptus forsteri* Tippmann, 1960 = *Bisaltis buquetii* Thomson, 1868; *Ataxia flaviceps* Breuning, 1942 = *Stenocorus obscurus* Fabricius, 1801; *Cryptocranium gounellei* Breuning, 1980 = *Cryptocranium cazieri* Lane, 1958; *Epectasis grossepunctata* Breuning, 1942 = *Saperda juncea* Newman, 1840; *Paraclitemnestra gigantea* Breuning, 1974 = *Jamesia lineata* Fisher, 1926; *Hippopsis tremata* Galileo and Martins, 1988 = *Hippopsis meinerti* Aurivillius, 1900; *Lamia punctata* Fabricius, 1792 = *Cerambyx (Lamia) Daviesii* Swederus, 1787; *Leiopus (Oedopeza) pogonocheroides* Audinet-Serville, 1835 = *Lamia ocellator* Fabricius, 1801; *Colobothea velutina* Bates, 1865 = *Lamia macularis* Olivier, 1792; *Xylergates dorotheae* Gilmour, 1962 = *Xylergates elaineae* Gilmour, 1962; *Colobothea velutina* Bates, 1865 = *Lamia macularis* Olivier, 1792; *Pretilia telephoroides* Bates, 1866 = *Saperda tuberculata* Fabricius, 1801; *Guyanestola macrophthalma* Breuning, 1961 = *Estoloides (Estoloides) angustifrons* Breuning, 1943; *Drycothaea marmorata* Martins and Galileo, 1990 = *Guyanestola brasiliensis* Breuning, 1974.

The following species are reinstated: *Strongylaspis costifer* Thomson, 1877; *Oxymerus lineatus* Dupont, 1836; *Hebestola operaria* Erichson, 1848; *Hesycha lateralis* Thomson, 1868.

The genus *Hephialtis* Thomson, 1864 is reinstated.

New combinations: *Hephialtis ruber* (Thunberg, 1822); *Protorma costifera* (Thomson, 1877); *Knulliana cincta spinifera* (Fabricius, 1801) comb. nov.; *Ataxia operaria* (Erichson, 1848) comb. nov.; *Paraclitemnestra lineata* (Fisher, 1926) comb. nov.; *Oedopeza ocellator* (Fabricius, 1801) comb. nov.; *Toronaeus incisus* (Bates, 1864) comb. nov.; *Pretilia tuberculata* (Fabricius, 1801) comb. nov. *Drycothaea turrialbae* (Breuning, 1943) comb. nov. *Drycothaea truncatipennis* new name for *Estola stictica* Breuning, 1942 (nec Bates, 1881)

Some host-plants are given.

Introduction

During my various visits through several institutions sheltering historical collections, I noticed some unknown synonymies which are interesting to publish. The acronyms used are:

MNH for the Muséum d'Histoire Naturelle de Paris,
BMNH for the British Natural History Museum in
London,

AMNH for the American Museum of Natural History in
New York,

USNM for the United States National Museum in Wash-
ington D.C. (Smithsonian Institution),

ICCM for the Carnegie Museum in Pittsburgh, Pennsylv-
ania,

MNHU for the Museum für Naturkunde der Humboldt
Universität in Berlin,

ZMK for the Zoologisk Museum in Copenhagen,

MNRJ for the Museu Nacional da Quinta da Boa Vista
in Rio de Janeiro,

ZSMA for the Zoologische Staatssammlung des Bay-
erischen Staates in Munich,

NRS for the Naturhistoriska Riksmuseet in Stockholm.

Concerning host plant records, the scientific name is followed by the herbarium collection refer-

ence number sampled by a botanist; several parts are distributed throughout main botanical institutions of the world.

1. Reinstated species:

Subfamily Prioninae

Tribe Macrotomini

***Protorma costifera* (Thomson, 1877), reinstated and new combination.**

Strongylaspis costifer Thomson, 1877: 275

Protorma scabrosa Waterhouse, 1880: 289, **new syno-
nymy**

Stenodontes (Protorma) scabrosus; Lameere, 1903: 215

Protorma recurvatum Williams, 1929: 143, pl. V, fig. 1,
new synonymy

Protorma recurvata Blackwelder, 1946: 552 [Catalog]

Lameere, without having seen Thomson's material, simply synonymized *Strongylaspis costifer* with *Strongylaspis corticarius* (Erichson, 1848). All the following authors (Villiers, 1980: 143 and

others) accepted Lameere's judgement. Fortunately, Thomson's type of *Strongylaspis costifer* is visible in the MNHN, and in fact, belongs to the genus *Protorma*. Later on, I saw Williams' type in ICCM of *Protorma recurvatum* and came to the conclusion that Williams species is just a synonym of Thomson's species. Williams wrote "It was carefully compared by me with the type of *Protorma scabrosa* Waterhouse, from which it is quite distinct." In fact, the description of *Protorma recurvatum* was based on a single male specimen, and the type of *Protorma scabrosa* in the BMNH is a female. The three described taxa belong to the same species, and the valid name is *Protorma costifera* (Thomson, 1877).

Subfamily Cerambycinae

Tribe Trachyderini

Oxymerus lineatus Dupont, 1838 reinstated

Oxymerus lineatus Dupont, 1838: 41, pl. 211, fig. 1 [Brazil]

Oxymerus approximatus Dupont, 1838: 44, pl. 212, fig. 2 [Cayenne]

Oxymerus elongatus Dupont, 1838, 1838: 45, pl. 213, fig. 1 [Cayenne]

Oxymerus distinguendus Dupont, 1838: 48, pl. 214, fig. 2 [Demerary]

Oxymerus aculeatus subsp. *lineatus* Hüdepohl, 1979: 20, pl. I, 1(m)

Oxymerus aculeatus var. *lineatus* Rémillat, 1988: 130

From November 1982 to January 1983 the species was particularly abundant around Kourou where I saw more than a thousand. I definitely do not agree with Hüdepohl synonymies and status changing (Hüdepohl, 1979: 20). The correct synonymy, after checking the types found in the MNHN, is listed above.

Subfamily Lamiinae

Tribe Pteropliini

Fabricius described (1775: 180) a *Stenocorus annulatus* from "America meridionali" belonging to the Banks collection (today in the BMNH). Fabricius later on (1801: 310) mentioned the synonymy with *Cerambyx annulatus* Olivier, 1792 and *Cerambyx hirtipes* Degeer, 1775 (belonging to the actual genus *Colobothea* Lepeletier and Audinet-Serville, 1825). In 1792: 314 he described a *Saperda lineata* from America meridionali. The two syntypes in the ZMK belong to Degeer's species, *Cerambyx hirtipes*.

In 1801: 326, he described *Saperda annulata* and gave the short following description in latin: thorace rotundato, subspinoso, antennarum articulis basi albis, which agrees with the type in Fabricius' collection in the ZMK (conspecific with Say's *Lamia crypta*); both the locality (America meridionali) and the synonymy (*Saperda lineata* 1792: 314) were wrongly associated with the above description. The confusion was obviously caused by an error in the printing of page 326 in Fabricius' Systema Eleutheratorum. It is clear that between the *Saperda nigricornis* (n° 47) and the *Saperda tristis* (n° 51) the synonymies were wrongly skipped. Bates in the supplement of Biologia Centrali Americana (1885: 347) wrote: *Ataxia Crypta* ... Said by Erichson to be the *Saperda lineata* of Fabricius (Ent. Syst. ii p. 314), and so registered in the Munich catalogue, but it does not agree with the description, which refers also, as Fabricius states, to a South-American (Brazilian ?) insect in Lunds collection. Erichson might have seen the type and Bates read the wrong description (Fabricius, 1792: 314). The name *Saperda annulata* being used several times by Fabricius, we decide to conserve Say's species name in order to avoid more confusion. The complete synonymy of *Ataxia crypta* is the following:

Ataxia crypta (Say, 1832)

Saperda annulata Fabricius, 1801 (*nec* Fabricius, 1792): 326, **new synonymy**

Lamia crypta Say, 1832: 5

Amniscus ? cryptus Haldeman, 1847: 47

Stenosoma crypta LeConte, 1873: 302

Ataxia crypta LeConte, 1873: 344

Ataxia sordida Haldeman, 1847: 56

Stenosoma sordida LeConte, 1852: 158

Consequently *Hebestola operaria* Erichson, 1848 synonymized with *Saperda annulata* Fabricius, *Parysatis nigritarsis* Thomson, 1868b, and *Parysatis flavescens* Bates, 1880 becomes a valid taxon and we can write the following synonymy:

Ataxia operaria (Erichson, 1848), reinstated

Hebestola operaria Erichson, 1848: 574

Parysatis nigritarsis Thomson, 1868b: 120

Esthlogena operaria Lacordaire, 1872: 600

Ataxia lineata Gemminger and Harold, 1873: 3100 [Catalog]

Parysatis flavescens Bates, 1880: 112

Tribe Onciderini

Lampedusa lateralis (Thomson, 1868) reinstated

Hesycha lateralis Thomson, 1868a: 63 [Cayenne]
Lampedusa obliquator Dillon and Dillon (*nec* Fabricius, 1801), 1945: 114, pl. V, fig. 10

***Lampedusa obliquator* (Fabricius, 1801)**

Lamia obliquator Fabricius, 1801: 303
Hypsioma obliquator Thomson, 1868a: 50
Plerodia obliquator Aurivillius, 1923: 342 [Catalog]

The type of *Hesycha lateralis* Thomson, 1868 is different from the type of *Lamia obliquator* Fabricius, 1801. Dillon and Dillon wrongly interpreted Fabricius' species and consequently *Lampedusa lateralis* (Thomson, 1868) is reinstated. Both species were encountered in French Guiana and belong to the genus *Lampedusa* as described by Dillon and Dillon, 1945: 112. Following the nomenclatural code, *Lamia obliquator* remains the type species of the genus *Lampedusa* Dillon and Dillon, 1945.

2. Reinstated genus:

Subfamily Prioninae

Tribe Callipogonini

The genus *Hephialtès* Thomson, 1864 was created for *Hephialtès tricostatus* Thomson, 1864: 286 based on a female specimen from Brazil. A year later he described the male as *badium*. Both are synonyms of *Trachyderes ruber* Thunberg, 1822. Lameere (1904: 28) synonymized the genus with *Anacanthus* Audinet-Serville, 1832 which he considered as a subgenus of *Stictosomus* Audinet-Serville, 1832. Whilst reading the description of the genus *Stictosomus* (page 153) we note incompatible features with Thunberg's species «Corselet sans crénelures» and «mandibules allongées, étroites, aiguës, denticulées intérieurement». Considering the description of *Anacanthus* (page 165 of the same Audinet-Serville's paper) a name which means in Greek "without spines" we find again incompatible features (corselet mutique ... arrondi...). Neither the genus nor the subgenus retained by Lameere is correct.

Though the type species of the 3 concerned genera (*Stictosomus semicostatus* Audinet-Serville, 1832, *Anacanthus costatus* Audinet-Serville, 1832 and *Hephialtès tricostatus* Thomson, 1864) belong obviously to the same natural subdivision of the tribe Callipogonini, I think it is unreasonable to synonymize them under the same genus.

***Hephialtès* Thomson, 1864 reinstated**

Type species: *Hephialtès tricostatus* Thomson, 1864 (= *Trachyderes ruber* Thunberg, 1822) [Brazil] by original designation.

Hephialtès Thomson, 1864: 285

***Hephialtès ruber* (Thunberg, 1822), comb. nov.**

Trachyderes ruber Thunberg, 1822: 305
Orthosoma badium Dejean (nomen nudum), 1837: 342
Hephialtès tricostatus Thomson, 1864: 286
Hephialtès badium Thomson, 1865: 577
Hephialtès sulcatus Lacordaire, 1869: 147
Hephialtès badius Gemminger and Harold, 1872: 2774 [Catalog]
Hephialtès tricostatus Gahan, 1895: 84
Stictosomus (Anacanthus) tricostatus Lameere, 1904: 28
Stictosomus ruber Lameere, 1912: 164
Stictosomus (Anacanthus) ruber Lameere, 1913: 38 [Catalog]
Anacanthus ruber Villiers, 1980: 152, fig. 20

3. New synonyms and consequently new combinations:

Subfamily Cerambycinae

Tribe Rhinotragini

***Odontocera simplex* White, 1855**

Odontocera simplex White, 1855: 189
Odontocera cinctiventris Bates, 1870a: 321, new synonymy

White's type, in the BMNH, is represented by a male specimen and Bates' type, in the MNHN, a female, although the latter was doubtful concerning the gender of his *O. cinctiventris*. I personally obtained many individuals of both sexes by rearing them on their host plant. I was already alerted by the elytral similarity of the two supposed species before having them bred. They emerged from three different species of trees all belonging to the family of Papilionaceae: *Taralea oppositifolia* Aublet [Denis Loubry 1611]; *Dipteryx punctata* (Black) Amshoff [Denis Loubry 1797]; *Monopteryx inpaë* Rodrigues [Denis Loubry 1739].

Tribe Callichromatini

Callichroma velutinum (Fabricius, 1775)
Cerambyx velutinus Fabricius, 1775: 167
Cerambyx spectabilis Voet, 1778: 12, pl. X, fig. 40, new synonymy
Cerambyx velutinus Olivier, 1795: 24, (67) pl. VI, fig. 41
Callichroma velutina Serville, 1833: 557
Callichroma velutinum White, 1853: 164

Callichroma velutinus Lacordaire, 1869: 16
Callichroma porphyrogenitum Bates, 1870: 333
Callichroma (Callichroma) velutinum Schmidt, 1924: 321

Considering that the illustration is sufficient, and that the locality is "Indes Occidentales", this leaves no doubt about the identity of this taxon. The epithet "*spectabilis*" still remains in all catalogs for obscure reasons.

Tribe Lissonotini

Lissonotus equestris (Fabricius, 1787)

Cerambyx biserratus Voet, 1778: 29, pl. XXV, fig. 137 [America]
Callidium equestre Fabricius, 1787: 153 [Cajennae]
Cerambyx unidentatus Olivier, 1795: 20, (n° 67) pl. XIX, fig. 145
Cerambyx cinctus Panzer in Voet, 1798: 2, pl. XXV, fig. 136
Cerambyx cingulatus Panzer in Voet, 1798: 2, pl. XXV, fig. 137
Lissonotus cinctus Dalman in Schönherr, 1817: 364 [Cayenna]
Lissonotus unidentatus Dalman in Schönherr, 1817: 364
Lissonotus equestris Dejean, 1821: 109
Lissonotus shepherdii Pascoe, 1859: 16 [Pará], **new synonymy**
Lissonotus shepherdii Blackwelder, 1946: 592 [Brazil] Catalog

I found in the MNHN at the Library of the French Entomological Society an example of Voet's book where the first pages are missing but *Cerambyx biserratus* was already described in 1778. Since the name *equestris* has been used for over a century I think it is better to conserve the well known name. On the other hand, *Lissonotus shepherdii* is exactly similar to Fabricius' type. To summarize my observation, the most common form with the red band complete seems to be a variety of *L. equestris*. I obtained this abundant species from several legume trees and vines. Both forms emerged from the same logs (interrupted or complete red band). Though the type locality is Cayenne the interrupted form is rarer in French Guiana.

Observed host plants: *Hymenaea courbaril* Linné (Caesalpiniaceae) [Sabatier and Prévost 3490]; *Bauhinia guianensis* Aublet (Caesalpiniaceae) [Scott Mori and al. 23641, 23678]; *Macrolobium* sp. (Caesalpiniaceae) [Scott Mori and al. 23467, 23409].

Tribe Trachyderini

Subtribe Bothriospilina

The identity of *Lamia spinifera* Fabricius, 1792 is not yet resolved. Martins and Moure (1973: 79), giving an illustration of the species, came to the conclusion that it might certainly be an Oriental or Ethiopian species, maybe belonging to the genus *Plocaederus* Thomson.

Knulliana cincta spinifera (Fabricius, 1792) new combination

Lamia spinifera Fabricius, 1792: 275; Fabricius, 1801: 292; Aurivillius, 1923: 606 [Catalog]; Blackwelder, 1946: 627 [Catalog] *incertae sedis*; Zimsen, 1964: 169 [Types]; Martins and Moure, 1973: 79 figs 1-2 (male).
Chion ochraceus Bates, 1885: 243, **new synonymy**
Chion cinctus var. *ochraceus* Schaeffer, 1908: 331
Chion cinctum v. *ochraceum* Blackwelder, 1946: 562 [Catalog]
Cerasphorus cinctum ochraceum Cazier and Lacey, 1952: 10 fig. 1
Knulliana cincta ochracea Linsley, 1962: 111 fig. 35 (distribution map)

Subtribe Ancylocerina

Callancyla croceicollis (White, 1855)

Eriphus croceicollis White, 1855: 292; Bates, 1870b: 429.
Calloppisma ruficollis Bates, 1870b: 419, **new synonymy**
Ancylocera ruficollis Gemminger and Harold, 1872: 2959 [Catalog]
Callancyla ruficollis Aurivillius, 1912: 446 [Catalog]

Bates wrote (1870: 429), concerning *Eriphus croceicollis* White, 1855, "Although taken by me, I do not find the species among my own reserved collection of Amazonian Longicorns". He would never have been able to guess White's misplacement whose type (BMNH) is a female. Bates, though uncertain about the sex of his *Calloppisma ruficollis* supposed the type (MNHN) to be a male.

Subfamily Lamiinae

Tribe Lamiini

Taeniotes farinosus (Linné, 1758)

Cerambyx farinosus Linné, 1758: 390
Cerambyx farinosus DeGeer, 1775: 108, pl. XIV, fig. 1 [Suriname]
Cerambyx pulverulentus Olivier, 1790: 302, **new synonymy**

- Cerambyx surinamensis maculosus* Panzer, 1794: 15, pl. V, fig. 8 [Suriname]
Cerambix pulverulentus Olivier, 1795: 50, (n° 67) pl. VII, fig. 46 b
Stenochorus pulverulentus Schönherr, 1817: 407 [Catalogue, Suriname]
Monochamus pulverulentus Dejean, 1821: 106
Taeniotes pulverulentus Audinet-Serville, 1835: 91 [Cayenne]
Monochamus (Taeniotes) farinosus Castelnau, 1840: 479 [Brazil]
Monochamus (Taeniotes) pulverulentus Castelnau, 1840: 479 [Brazil]
Taeniotes farinosus Bates, 1865b: 110 [Brazil: Amazon, Tapajos]
Taeniotes guttularis Schwarzer, 1929: 364, fig. 16-17
Taeniotes pulverulentus m. guttularis Breuning, 1943b: 248
Taeniotes pulvurentus Bosq, 1943: 109
Taeniotes farinosus Dillon and Dillon, 1941 pl. I, fig. 6
Taeniotes farinosa Blackwelder, 1946: 594 [Catalog]
Taeniotes pulverulenta Blackwelder, 1946: 594 [Catalog]

Olivier's drawing representing *Cerambyx pulverulentus* closely resembles the species commonly collected in Cayenne. Linné's *farinosus*, which is neither in the London Linnean Society nor in Gustave-Adolphe's collection in Uppsala, has in fact no type. Linné mentioned after his laconic description the corresponding illustration in Sybille Merian's book published in 1719 (table 24, f. *infima*) and another reference published by Degeer before the tenth edition of Linné's *Systema Naturae*. The iconotype belongs to the prelinnean literature; I ignore what is to be done in this case. I consulted this very rare book with colored plates and I am certain that *C. pulverulentus* is a synonym of *C. farinosus* and their localities (Suriname and Cayenne) confirm the synonymy. The species is very common on the introduced breadfruit tree *Artocarpus altilis* (S. Parkinson) Fosberg (Moraceae). I do not understand why these two taxa were considered as different by all authors, except for Bates who had already reported the synonymy (1865b: 110).

Type Apomecynini

Bisaltes buqueti Thomson, 1868

- Bisaltes buquetii* Thomson, 1868b: 111 [Cayenne]
Bisaltes buqueti Gemminger and Harold, 1873: 3102 [Catalog]

- Bisaltes sp.* Bates, 1885: 346, pl. IX, fig. 21a
Bisaltes batesi Aurivillius, 1923: 609 [Catalog]
Ptericoptus forsteri Tippmann, 1960: 152, pl. IX, fig. 21a, new synonymy
Bisaltes (Bisaltes) buqueti Breuning, 1960: 177 [Catalog]
Bisaltes (s.str.) forsteri Breuning, 1971: 284

Tippmann's type is in the USNM and the type of Thomson in the MNHN. The species was reared on a Convolvulaceae vine belonging to the genus *Maripa* [Scott Mori *et al.* 23592].

Tethystola brasiliensis Breuning, 1940

- Tethystola brasiliensis* Breuning, 1940: 42
Tethystola flavoapicalis Breuning, 1942: 141, new synonymy

Both types are at the MNHN and were collected in the Brazilian state of Pernambuco. The type of *T. flavoapicalis* is a big specimen.

Tribe Pteropliini

Ataxia obscura (Fabricius, 1801)

- Stenocorus obscurus* Fabricius, 1801: 307 [America meridional]
Esthlogena sulcata Bates, 1866: 290
Parysatis collaris Thomson, 1868: 119
Parysatis sulcata Aurivillius, 1922: 293 (Catalog)
Parysatis obscura Aurivillius, 1922: 292 (Catalog)
Ataxia flaviceps Breuning, 1942: 137, new synonymy
Ataxia obscura Breuning, 1961: 53, fig. 7

Cryptocranium cazieri Lane, 1958

- Cryptocranium cazieri* Lane, 1958: 6, fig. 2
Ornatodesisa pulchra Breuning, 1961: 19
Cryptocranium gounellei Breuning, 1980: 68, new synonymy

Lane's type (a female from Peru) belongs to the AMNH and Breuning's type (a male with flattened legs from the Brazilian Goiás) belongs to the MNHN. The recent synonymy published by Monné and Giesbert (1992: 252) for a specimen already described by Breuning from the Brazilian state of Pará, and the presence of this species in French Guiana, lead us to consider that this very rare species is widely distributed in northern South America.

Epectasis juncea (Newman, 1840)

- Saperda juncea* Newman, 1840: 13
Epectasis attenuata Bates, 1866: 294
Epectasis grossepunctata Breuning, 1942: 175, new synonymy

Epectasis juncea m. attenuata Breuning, 1961c: 25

Breuning's type belongs to the USNM, the type of *E. attenuata* is in the MNHN.

Tribe Onciderini

Paraclytemnestra lineata (Fisher, 1926), **new combination**.

Jamesia lineata Fisher, 1926: 14

Paraclytemnestra gigantea Breuning, 1974a: 240, **new synonymy**

This incredible species from the Antilles merits its own genus. Breuning's type belongs to the MNHU, though Fisher's type belongs to the USNM.

Tribe Agapanthiini

Hippopsis meinerti Aurivillius, 1900

Hippopsis meinerti Aurivillius, 1900: 416; Aurivillius, 1923: 357 [Catalog]

Hippopsis meinerti Blackwelder, 1946: 606 [Catalog]; Galileo and Martins, 1988a: 185 figs 17-20; Galileo and Martins, 1988b: 205

Hippopsis freyi Breuning, 1955: 661

Hippopsis (Hippopsis) lemniscata m. meinerti Breuning, 1961b: 197 [Catalog]; Breuning, 1962: 10

Hippopsis (Hippopsis) freyi Breuning, 1961b: 197 [Catalog]; 1962: 18

Hippopsis (Hippopsis) lemniscata tobagoensis Breuning, 1962: 10

Hippopsis tremata Galileo and Martins, 1988a: 184, figs. 13-16, **new synonymy**

Aurivillius' type, in the NRS is a male, and the abdominal depression is absolutely identical to the drawing given in the paper of Galileo and Martins to characterize their new species *Hippopsis tremata*. I compared Aurivillius' type with my own material of both sexes from French Guiana and identified by Martins as *H. tremata*.

Tribe Acanthoderini

Acanthoderes daviesi (Swederus, 1787)

Cerambyx (Lamia) daviesii Swederus, 1787: 195, pl. VIII, fig. 6

Cerambyx daviesii Gmelin, 1790: 1837

Lamia punctata Fabricius, 1792: 272 [Cajennae], **new synonymy**

Lamia daviesii Schönherr, 1817: 380 [America, Cayenna]

Acanthoderes daviesii Dejean, 1835: 336

Acanthoderes daviesii Audinet-Serville, 1835: 29 [Cayenne]

Acanthoderes swederi White, 1855: 360, pl. IX, fig. 6 [Para]

Acanthoderes daviesi Thomson, 1864: 17

Acanthoderes daviesi var. *swederi* Gemminger and Harold, 1873: 3146 [Catalog]

Acanthoderes daviesi Bodkin, 1919: 268

Acanthoderes (Acanthoderes) daviesi Aurivillius, 1923: 384 [Catalog]

Acanthoderes (Acanthoderes) punctata Aurivillius, 1923: 385 [Catalog]

Acanthoderes (Acanthoderes) daviesi Buck, 1959: 604

The common and widely distributed epiphyte *Clusia grandiflora* Splitgerber (Clusiaceae) [Denis Loubry 1827] breeds *A. daviesi*.

Tribe Acanthocinini

Oedopeza ocellator (Fabricius, 1801), **new combination**

Lamia ocellator Fabricius, 1801: 287 [America meridionali]

Leiopus (Edopeza) pogonocheroides Audinet-Serville, 1835: 88, **new synonymy**

Leiopus pogonocheroides Castelnau, 1840: 464 [Cayenne]

Oedopeza pogonocheroides White, 1855: 394

Ozineus ocellator Aurivillius, 1923: 398 [Catalog]

I do not understand why this very common species was not recognized by Aurivillius. The type of Fabricius is a male with a characteristic swollen protarsal segment. I obtained this species from 16 different tree species, mainly belonging to the legume group: *Couma guianensis* Aublet (Apocynaceae) [Christian Feuillet 2311]; *Simarouba amara* Aublet (Simaroubaceae) [Christian Feuillet 2375]; *Theobroma cacao* Linné (Sterculiaceae) [Jean-Jacques de Granville det.]; *Swartzia lomatoxylla* (Benth) Pittier (Caesalpiniaceae) [Denis Loubry, 1585]; *Swartzia panacoco* (Aublet) Cowan (Caesalpiniaceae) [Denis Loubry, 1804]; *Machaerium* sp. (Papilionaceae) [Denis Loubry, 1815]; *Taralea oppositifolia* Aublet (Papilionaceae) [Denis Loubry, 1611]; *Abarema curvicarpa* (Irwin) Barneby and Grimes (Mimosaceae) [Grimes and Sabatier 3312]; *Abarema jupunba* (Willdenow) Barneby and Grimes (Mimosaceae) [Christian Feuillet 2307, Denis Loubry 1875 and Scott Mori 23688]; *Balizia pedicellaris* (de Candolle) Barneby and Grimes (Mimosaceae) [Christian Feuillet 1341]; *Enterolobium schomburgkii* Benth (Mimosaceae) [Daniel Sabatier 2367]; *Enterolobium* sp. (Mimosaceae) [Denis Loubry 1763]; *Hydrochorea corymbosa* (L.C.

Richard) Barneby and Grimes (Mimosaceae) [Denis Loubry 1300]; *Inga melinonis* Sagot (Mimosaceae) [Denis Loubry 1780]; *Inga* sp. (Mimosaceae) [Scott Mori 23456]; *Parkia nitida* Miquel (Mimosaceae) [Daniel Sabatier 1511 and Denis Loubry 1564].

***Toronaeus incisus* (Bates, 1864), new combination.**

Nyssodryes incisus Bates, 1864: 155

Nyssodryes incisus Gilmour, 1965: 599 [Brazil] Catalogue

Stenolis incisus Monné, 1985: 535.

The rearing of this rare species demonstrates clearly its greater affinity with the genus *Toronaeus* (where majority were bound to Burseraceae trees), confirmed by its taxonomic features. It was obtained on a still unidentified Burseraceae [Denis Loubry 1318]

***Xylergates elaineae* Gilmour, 1962**

Astynomus hilaris: Prudhomme, 1906: 38

Xylergates sp. (*hilaris* Dejean?) Duffy, 1960: 258, figs. 158-159 (Larva)

Xylergates elaineae Gilmour, 1962: 277, pl. III, fig. 3

Xylergates dorotheae Gilmour, 1962: 279, pl. III, fig. 4, **new synonymy**

Gilmour's two types belong to the American Museum of Natural History in New York. The species commonly attacks fallen trees belonging to the family of Lecythidaceae. I found this species from seven different host plants in French Guiana: *Lecythis poiteaui* O.C. Berg [Scott Mori det. 1995]; *Lecythis holcogyne* (Sandwith) Mori [D. Sabatier and M.F. Prévost 2108]; *Eschweilera sagotiana* Miers [Daniel Sabatier 1762]; *Eschweilera micrantha* (O.C. Berg) Miers [Scott Mori 23479]; *Eschweilera coriacea* (A.P. de Candolle) Martius ex O.C. Berg [Denis Loubry 1846, 1341]; *Eschweilera congestiflora* (R. Benoist) Eyma [Scott Mori 23478]; *Couratari guianensis* Aublet [Daniel Sabatier 1771 and Denis Loubry 1337].

Tribe Colobotheini

***Colobothea macularis* (Olivier, 1792)**

Lamia macularis Olivier, 1792: 474 [Suriname]

Cerambyx macularis Olivier, 1795: (67) 98 pl. XX, fig. 154 [Suriname]

Colobothea velutina Bates, 1865a: 216, **new synonymy**

Colobothea macularis: Aurivillius, 1923: 454 [Catalog]

I had the opportunity to consult and to photograph the original paintings of Olivier which served to illustrate his famous "Entomologie..." still conserved in his family, six generations later. These paintings reveal more details than the printed and post-colored plates found in the remaining examples of this very rare book and thus permit the above synonymy. I obtained several adults from *Hydrochorea corymbosa* (L.C. Richard) Barneby and Grimes (Mimosaceae) [Denis Loubry 1300]

Tribe Pretiliini

***Pretilia tuberculata* (Fabricius, 1801), new combination.**

Saperda tuberculata Fabricius, 1801: 331 [America meridionali]

Pretilia telephoroides Bates, 1866: 302; Lacordaire, 1872: 910; Gilmour, 1962: 125; Martins and Galileo, 1990b: 707 figs 1-11, **new synonymy**

Eupogonius tuberculatus Aurivillius, 1923: 312 [Catalog]; Blackwelder, 1946: 600 [Catalog]

Tribe Calliini

***Drycothaea angustifrons* (Breuning, 1943)**

Estoloides (Estoloides) angustifrons Breuning, 1943a: 56 [Guyane: Maroni]; Breuning, 1974b: 59.

Guyanestola macrophthalma Breuning, 1961a: 248, **new synonymy**

Guyanestola macrophthalma Breuning, 1963: 507 [Catalog]

Drycothaea angustifrons Tavakilian, 1991: 450; Galileo and Martins, 1991: 250

Drycothaea macrophthalma Galileo and Martins, 1991: 251

In 1990, the blocked situation concerning Frey's collection made impossible the examination of Breuning's type during my journey to Munich. The situation is now normal since the collection was transferred from Frey's house in Tutzing to the Staatliches Museum in Munich (ZSMA). I am grateful to Patrick Bleuzen who sent me the picture of *Guyanestola macrophthalma* and checked the claws, confirming that they are appendiculated as Galileo and Martins and I already guessed. This illustrates how uncomfortable can be the position of an entomologist revising a group of which the types are still in private collections.

I obtained this species from *Cecropia palmata* Willdenow (Cecropiaceae) [Denis Loubry 1570].

***Drycothaea brasiliensis* (Breuning, 1974)**

Guyanestola brasiliensis Breuning, 1974b: 49 [Brazil: Amazonas]

Drycothaea marmorata Martins and Galileo, 1990a: 611, fig. 4, **new synonymy**

Drycothaea brasiliensis Galileo and Martins, 1991: 251

Breuning's type is in the ZSMA. The host plant is still *Cecropia palmata* Willdenow (Cecropiaceae) [Denis Loubry 1570].

***Drycothaea truncatipennis* new name for *Estola stictica* Breuning, 1942 (nec Bates, 1881)**

Estola stictica Breuning, 1942: 163

The type belongs to the USNM (ex Tippmann's collection n° 41605) and is a male collected in the Brazilian state of Santa Catarina. The structure of the claws transfers this species from the tribe Desmiphorini to Calliini in the genus *Drycothaea*. Unfortunately another *Drycothaea stictica* Bates, 1881 already exists from Guatemala. I propose the specific name *D. truncatipennis* which fits better with the elytral aspect of that misplaced species.

***Drycothaea turrialbae* (Breuning, 1943), new combination**

Estoloides (Estoloides) turrialbae Breuning, 1943: 56

Estoloides (Estoloides) turrialbae Breuning, 1963: 507

Estoloides turrialbae Chemsak, Linsley and Noguera, 1992: 119 (Catalog)

The type belongs to the USNM (ex Tippmann's collection n° 41596) and is a female labelled as *Parestola turrialbae*! As seen above, this species is transferred from Desmiphorini to the tribe Caliini for the same reason.

References

- Audinet-Serville, J. G.** 1832. Nouvelle classification de la famille des longicornes. Annales de la Société Entomologique de France. Paris 1 (1): 118-201
- Audinet-Serville, J. G.** 1835. Nouvelle classification de la famille des Longicornes 3ème Tribu, Lamiaires, Lamiaires. Annales de la Société Entomologique de France. Paris 4 (1): 5-100.
- Aurivillius, C.** 1900. Verzeichniss der von Dr F. Meinert im Jahre 1891 in Venezuela gesammelten Cerambyciden. Öfversigt af Kongl. Svenska Vetenskaps-Akademiens Föreläsningar 57 (4): 409-421, 1 fig.
- Aurivillius, C.** 1912. Cerambycidae: Cerambycinae. Coleopterorum Catalogus 22 (39): 1-574. W. Junk et S. Schenkling, Berlin.
- Aurivillius, C.** 1923. Lamiinae II. Coleopterorum Catalogus 23 (74): 323-704. W. Junk et S. Schenkling, Berlin.
- Bates, H. W.** 1864. XIX. Contributions to an Insect Fauna of the Amazon Valley. Coleoptera: Longicornes. The Annals and Magazine of Natural History. London 13 (3): 144-164.
- Bates, H. W.** 1865a. XXV. Contributions to an Insect Fauna of the Amazons Valley. Coleoptera: Longicornes. The Annals and Magazine of Natural History. London 15 (3): 213-225.
- Bates, H. W.** 1865b. X. Contributions to an Insect Fauna of the Amazons Valley. Coleoptera: Longicornes. The Annals and Magazine of Natural History. London 16 (3): 101-113.
- Bates, H. W.** 1866. XXXIII. Contributions to an Insect Fauna of the Amazons Valley. Coleoptera: Longicornes. The Annals and Magazine of Natural History. London 17 (3): 288-303.
- Bates, H. W.** 1870a. XVI. Contributions to an Insect Fauna of the Amazon Valley (Coleoptera, Cerambycidae). The Transactions of the Entomological Society of London, 1870 (part III): 243-335.
- Bates, H. W.** 1870b. XIX. Contributions to an Insect Fauna of the Amazon Valley (Coleoptera, Cerambycidae). The Transactions of the Entomological Society of London, 1870 (part IV) December: 391-444.
- Bates, H. W.** 1880. Biologia Centrali-Americana, Insecta, Coleoptera, Cerambycidae, Lamiidae 5: 17-152, pls. III-XI.
- Bates, H. W.** 1885. Biologia Centrali-Americana, Insecta, Coleoptera, Cerambycidae Supplement to Longicornia. 5: 249-436, pls. XVII-XXIV.
- Blackwelder, R. E.** 1946. Checklist of the coleopterous insects of Mexico, Central America, the West Indies and South America. Part 4. Bulletin. United States National Museum 185 (4): 551-763.
- Bodkin, G. E.** 1919. Notes on the Coleoptera of British Guiana. The Entomologist's Monthly Magazine. London 55: 264-272.
- Bosq, J. M.** 1943. Agregados al Catálogo de los Longicornes de la República Argentina. Revista Argentina de Zoogeografía. Buenos Aires 3 (3): 103-112.
- Breuning, S.** 1940. Novae species Cerambycidarum VIII. Folia Zoologica et Hydrobiologica. Riga 10: 37-85.
- Breuning, S.** 1942. Novae species Cerambycidarum XI. Folia Zoologica et Hydrobiologica. Riga 11: 113-175.
- Breuning, S.** 1943 a. Novae species Cerambycidarum XII. Folia Zoologica et Hydrobiologica. Riga 12: 12-66.
- Breuning, S.** 1943 b. Études sur les Lamiaires (Coleop., Cerambycidae). Douzième tribu : Agniini Thomson. Novitates Entomologicae, Paris 3 (89-106): 137-280, figs 1-157.

- Breuning, S.** 1955. Neue Cerambyciden von der Insel Trinidad. *Entomologische Arbeiten aus dem Museum G. Frey, Tutzing bei München* 6 (2): 659-661.
- Breuning, S.** 1960. Catalogue des Lamiaires du Monde (Col. Céramb.). Verlag des Museums G. Frey, Tutzing bei München (3) juli 1960: 109-182.
- Breuning, S.** 1961a. Neue Lamiiden aus dem Museum G. Frey (Col. Cerabm.). *Entomologische Arbeiten aus dem Museum G. Frey, Tutzing bei München* 12(1): 245-250.
- Breuning, S.** 1961b. Catalogue des Lamiaires du Monde (Col. Céramb.). Verlag des Museums G. Frey, Tutzing bei München (4) Jänner 1961: 183-284.
- Breuning, S.** 1961c. Révision des Pteropliini (Col. Cerambycidae). *Pesquisas, Porto Alegre (Zoologia)* 5 (9): 5-61, 7 figs.
- Breuning, S.** 1962. Révision des Agapanthiini Muls. Américains (Col., Cerambycidae). *Pesquisas, Porto Alegre (Zoologia)* 6 (13): 5-48, 9 figs.
- Breuning, S.** 1963. Catalogue des Lamiaires du Monde (Col. Céramb.). Verlag des Museums G. Frey, Tutzing bei München (7) April 1963: 463-555.
- Breuning, S.** 1971. Révision des espèces américaines de la tribu des Apomecynini Lac. (Coleoptera, Cerambycidae). *Entomologische Abhandlungen Staatliches Museum für Tierkunde in Dresden* 37 (3): 209-335, 21 figs.
- Breuning, S.** 1974a. Neue Lamiinae (Coleoptera, Cerambycidae). *Mitteilungen aus dem Zoologischen Museum in Berlin* 50 (2): 237-241.
- Breuning, S.** 1974b. Révision des Rhodopinini Américains (Coleoptera, Cerambycidae). *Studia Entomologica, Petrópolis* 17 (1-4): 1-210, 26 figs.
- Breuning, S.** 1980. Nouveaux Cerambycidae Lamiinae néotropicaux. *Bulletin de la Société Entomologique de France, Paris* 85 (3-4): 67-71.
- Buck, P.** 1959. Cerambycidae in der Sammlung des Instituto Anchieta de Pesquisas. *Pesquisas, Porto Alegre* 3: 577-609.
- Cazier, M., and Lacey, L.** 1952. The Cerambycidae of the Bahama Islands, British West Indies (Coleoptera). *American Museum Novitates, New York* 1588: 1-55.
- Chevrolat L. A. A.** 1862. Coléoptères de l'île de Cuba. Notes, synonymies et descriptions d'espèces nouvelles. Familles des cérambycides et des parandrides. *Annales de la Société Entomologique de France, Paris* 4 (2): 245-280.
- Dalman, J. W.** In Schönherr 1817. *Synonymia insectorum, oder: Versuch einer Synonymie aller bisher bekannte Insekten; nach Fabricii Systema Eleutheratorum geordnet. Eleutherata oder Käfer* 1 (3): xi + 1-506. Stockholm.
- Degeer, C.** 1775. *Mémoires pour servir à l'histoire des insectes* 5: 1-448, 16 pls. Stockholm.
- Dejean, P. F. M. A.** 1821. Catalogue des coléoptères de la collection de M. le baron Dejean. 136 pp. Paris.
- Dejean, P. F. M. A.** 1835. Catalogue des coléoptères de la collection de M. le Comte Dejean. livr. 4, 257-360. Paris.
- Dejean, P. F. M. A.** 1837. Catalogue des coléoptères de la collection de M. le Comte Dejean. livr. 5, 361-443. [Seconde Edition] Paris.
- Dillon, L. S. and Dillon, E. S.** 1941. The tribe Monochamini in the Western Hemisphere (Coleoptera: Cerambycidae). Scientific Publications. Reading Public Museum and Art Gallery. Reading, Pennsylvania. 1: 1-135, 5 pls.
- Dillon, L. S., and Dillon, E. S.** 1945. The tribe Onciderini (Coleoptera: Cerambycidae). Part I. Scientific Publications. Reading Public Museum and Art Gallery. Reading, Pennsylvania. 5: i-xv + 1-186.
- Dupont, H.** 1838. Monographie des trachydérides de la famille des longicornes. *Magasin de Zoologie, Paris* 8: i-xiii + 29-59, pls. 186-200 and 204-224.
- Erichson, W. F.** 1848. Die Insekten in Schomburg's Reisen in Britisch-Guiana. Leipzig 3: 570-574.
- Fabricius, J. C.** 1775. *Systema entomologiae, sistens insectorum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, observationibus.* Korte; Flensburgi et Lipsiae 30 + 832 pp.
- Fabricius, J. C.** 1787. *Mantissa insectorum sistens eorum species nuper detectas adjectis characteribus genericis, differentiis specificis, emendationibus, observationibus.* Proft, Hafniae 1: xx + 348 pp.
- Fabricius, J. C.** 1792. *Entomologia systematica, emendata et aucta. Secundum classes ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus.* Proft, Hafniae 1: xx + 538 pp.
- Fabricius, J. C.** 1801. *Systema eleutheratorum secundum ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus.* Bibliopoli Academici Novi, Kiliae 2: 1-687.
- Fisher, W. S.** 1926. Descriptions of new West Indian longicorn beetles of the subfamily Lamiinae. *Proceedings of United States National Museum, Washington* 68 (22) 2623: 1-40.
- Fisher, W. S.** 1930. Notes on the Rhinotragine beetles of the family cerambycidae, with descriptions of new species. *Proceedings of United States National Museum, Washington* 77(19): 1-20.
- Gahan, C.** 1895. On the longicorn Coleoptera of the West India Islands. *Transactions of the Entomological Society of London* [1895]: 79-140.
- Galileo, M. H. M. and Martins, U. R.** 1988a. Notas sobre Agapanthiini (Coleoptera, Cerambycidae, Lamiinae). I. Hippopsis do grupo solangeae. *Revista Brasileira de Entomologia, São Paulo* 32 (2): 179-185, 20 figs.
- Galileo, M. H. M., and Martins, U. R.** 1988b. Notas sobre Agapanthiini (Coleoptera, Cerambycidae, Lamiinae). V. Revisão do gênero Hippopsis Lepeletier and A.-Serville, 1825. *Revista Brasileira de Entomologia, São Paulo* 32 (2): 199-207, 9 figs.
- Galileo, M. H. M., and Martins, U. R.** 1991. Revisão da tribo Calliini (Coleoptera, Cerambycidae, Lami-