

ORSTOM, Laboratoire de Nématologie, B.P. 1386, Dakar, Sénégal¹
University Gadjja Mada, Department of Agronomy, Yogyakarta, Indonesia²

EFFECT OF THE RHIZOSPHERE MICROFLORA ON *PASTEURIA PENETRANS* PARASITIZING *MELOIDOGYNE GRAMINICOLA*

by

R. DUPONNOIS¹, C. NETSCHER (†)² and T. MATEILLE¹

Summary. The effect of the rhizosphere microflora associated with spores of *Pasteuria penetrans* on the attachment of the spores on juveniles of *Meloidogyne graminicola* has been studied. The microflora stimulated the attachment of the spores on the juveniles and consequently reduced the invasion of the roots of tomato plants by the nematodes. The role of a helper rhizosphere microflora for biocontrol of root-knot nematodes by *P. penetrans* is suggested.

Pasteuria penetrans is an obligate endoparasite of nematodes which has potential as a biological control agent (Oostendorp *et al.*, 1991; Zaki and Maqbool, 1992). The relationship between this actinomycete and root-knot nematodes is well documented (Davies *et al.*, 1992; Davies and Danks, 1992). However, the ways in which soil factors influence the *P. penetrans* – nematode interaction are virtually unknown, particularly that between the nematodes, *P. penetrans* and the microflora. The aim of this paper is to examine the influence of the rhizosphere microflora on the parasitic relationship between *P. penetrans* Sayre *et* Starr and *Meloidogyne graminicola* Golden *et* Birchfield.

Materials and methods

A population of *Meloidogyne graminicola* was cultured for two months on tomato (*Lycopersicon esculentum* Mill.) cv. Roma. Then the tomato roots were harvested, cut into short lengths and placed in a mist chamber for one

week to allow second stage juveniles (J2) to hatch from the eggs (Seinhorst, 1950).

Seeds of tomato cv. Roma were planted in 0.3 dm³ pots previously filled with a sandy soil (sand 84.7%; silt 4.1%; clay 11.2%; pH (H₂O) 7.6) sampled in a vegetable field from the experimental station of the École Nationale Supérieure d'Agriculture (ENSA, Thiès, Sénégal) heavily infested with juveniles of *M. javanica* parasitized by *P. penetrans* (20,000 J2 per dm³, 80% of infected juveniles). Two months after sowing, the plants were uprooted, the root systems were gently washed, cut into short lengths and blended in 100 ml sterile water using an Ultraturax blender. The suspension was sieved through a 45 µm sieve and the filtrate was divided in two parts, each of which were centrifuged (3000 g, 20 min); the supernatants were discarded. The pellet of the first sample was resuspended in alcohol (97%) for 12 h at 25 °C to obtain *P. penetrans* spores free of contaminant microflora, and centrifuged (3000 g, 20 min) three times to replace the alcohol with sterile sodium-phosphate buffer (50 ml, pH 7).

The pellet of the second sample was directly re-suspended in the same sterile buffer without sterilization by alcohol. The two suspensions of spores disinfected or not were incubated for two weeks in the dark at 25 °C.

Each week, attachment tests were performed in glass tubes (18x2 cm) with the spores of *P. penetrans* disinfected or not, adding 5 µl of the suspensions (10^4 spores) to one ml of distilled water containing 100 juveniles of *M. graminicola*. There were five replicates per condition. The suspensions were incubated for 12 hours at 25 °C. The number of juveniles encumbered with *P. penetrans* spores and the number of spores per juvenile were determined with the aid of a dissecting microscope (magnification x 400). All the data were statistically analysed according to the Mann Whitney U test ($P < 0.05$). At the same time, the populations of bacteria were determined in the two suspensions of *P. penetrans* spores disinfected or not by spread-plate counts on 0.3% tryptic soy agar (Martin, 1975).

Two week-old seedlings of tomato cv. Roma were transplanted in 1 dm³ pots filled with the same sandy soil that was used for the production of spores of *P. penetrans* but which had been previously autoclaved (140 °C, 40 min). Two weeks after transplanting, the plants were inoculated either with 100 juveniles of *M. graminicola*, or with 5.9×10^6 spores of *P. penetrans* disinfected or not with alcohol following the technique described above, or with 100 juveniles of *M. graminicola* and with 5.9×10^6 spores of *P. penetrans* disinfected or not, together. There were ten replicates for each inoculum condition placed in a randomised block design in a glasshouse (25 °C day, 20 °C night). One month later, each root system was gently washed free of soil and the galls with egg masses were counted. The roots were then cut into short lengths and placed in a mist chamber for two weeks to collect the juveniles as described above. The oven dried weights of shoot and root (1 week at 65 °C) were measured and

compared to non inoculated plants (control). Then the root systems were ground to a powder and resuspended in distilled water. The spores of *P. penetrans* were recovered as described before and counted using a Mallassez counting cell. The data were analysed with the Mann Whitney U test ($P < 0.05$).

Results

In the attachment tests 100% of the juveniles of *M. graminicola* were parasitized by *P. penetrans* whatever the time of incubation of the spores in the buffer and there were no differences between the disinfected and the non disinfected spores. Only the number of spores per juvenile differed (Fig. 1). Twenty to twenty-four spores attached to the juveniles in the first test with no difference between the disinfected and the non-disinfected spores. However, after one and two weeks of incubation, the number of spores attached per juvenile decreased, more with the spores previously disinfected with alcohol than with the non-disinfected spores. In the suspension containing the non-disinfected spores of *P. penetrans*, the population of bacteria remained constant during the two weeks of incubation in the sodium-phosphate buffer. No bacteria were detected in the suspension containing the spores previously disinfected with alcohol.

One month after the inoculation of tomato plants with juveniles of *M. graminicola*, 87.8% of the juveniles were able to invade the roots and to mature in gravid females (Table I). When the nematode inoculum was supplemented with the spores of *P. penetrans*, the development rate of the juveniles was strongly reduced, significantly more when the spores were not disinfected with alcohol than when they were. Comparing with the non-inoculated plants (control), the juveniles of *M. graminicola* alone reduced the development of the roots without any effect on the shoots. When the spores of *P. penetrans* were added to the juveniles, the root growth




Fig. 1 - Evolution of the number of spores of *Pasteuria penetrans* disinfected or not with 97% alcohol on the juveniles of *Meloidogyne graminicola* (bars represent standard errors) and evolution of the total population of bacteria in the suspension of non disinfected spores (columns with the same letter are not significantly different: $P \geq 0.05$).

was restored. Plant growth was greater when the spores, especially not disinfected, were inoculated alone. Most of the next generation of juveniles which were extracted from the roots in the mist chamber were encumbered with spores of *P. penetrans* (66 to 74% of the juveniles), but without any significant effect of the alcohol disinfection previous to the inoculation of the plants. After the roots were dried and ground to powder, the number of spores of *P. penetrans* extracted was the same whether the parent spores were disinfected or not.

Discussion

The disinfection of the spores of *P. penetrans* with alcohol eliminated the associated microflora as no bacteria were detected later in the suspensions, but it did not affect the attachment of the spores to the juveniles of *M. graminicola*.

The spores of *P. penetrans* which were incubated in the sodium-phosphate buffer became less aggressive to the juveniles of *M. graminicola* over one and two weeks. The attachment is between two organisms whose external surfaces are electronegatively charged and hydrophobic (Himmelhoch *et al.*, 1979; Afolabi *et al.*, 1995; Davies *et al.*, 1996). Thus, the decrease of aggressiveness could be due to a change of hydrophobic characteristics or to a progressive saturation of the negative sites of the spore surface by the Na^{2+} cations of the buffer at neutral pH. For example, these characteristics of the Na^{2+} cations are commonly used in soil particle dispersion (Bartoli *et al.*, 1990; Feller *et al.*, 1991). Nevertheless, the spores of *P. penetrans* which were incubated together with the microflora over two weeks, were more aggressive against the nematodes than the spores whose associated microflora was eliminated with alcohol. Even if the microflora did not develop in the buffer, bacterial metabolites would have been excreted in the suspension. They could act enzymatically on the external tissues of the spores thereby exposing the parasporal fibres and enabling their attachment to the cuticle of the nematodes (Davies and Danks, 1993). They might also modify the chemical environment of the spores and facilitate their attachment by producing or excreting directly (Rozycki, 1987) simple sugars and oligosaccharides which could involve lectin-carbohydrate interaction (Bird *et al.*, 1989; Ibrahim, 1991).

The same helper potential of the microflora may act efficiently in the soil. Despite sterilization of the soil, the microflora associated with the spores of *P. penetrans* induces the best attachment of the spores to the cuticle of the juveniles, unless compounds toxic to nematodes (Hoffmann-Hergarten and Sikora, 1995) are produced in addition. Consequently, the motility of the juveniles (Stirling *et al.*, 1990) and the invasion of the plant roots (Brown and Smart, 1985; Davies *et al.*, 1988) are reduced. The subsequent production of spores detected

TABLE I - Effect of the disinfection of the spores of *Pasteuria penetrans* on the development of *Meloidogyne graminicola* on tomato plants and consecutive plant growth.

Inoculum	No of galls per plant	Shoot biomass (mg dry weight)	Root biomass (mg dry weight)	No. of spores per plant
Control		1880 b	818 b	
<i>M. graminicola</i>	87.8 a	1700 b	372 d	–
<i>P. penetrans</i> disinfected	–	2070 ab	1117 a	–
<i>P. penetrans</i> not disinfected	–	2400 a	1112 a	–
<i>M. graminicola</i> + <i>P. penetrans</i> disinfected	43.9 b	2200 ab	540 bc	4.6x10 ⁶ a
<i>M. graminicola</i> + <i>P. penetrans</i> not disinfected	20.7 c	1900 b	424 c	3.7x10 ⁵ a

In each column, data followed by the same letter are not significantly different according to the Mann Whitney U test (P < 0.05)

in the powdered roots are not different, certainly because most of the spores were spread during nematode extraction in the water mist.

Thus, the rhizosphere microflora can enhance the biocontrol of the root-knot nematodes by *P. penetrans* by increasing the attachment of the spores to the nematodes. Further research is required to study the bacteria which are concerned and the helper mechanisms they induce, and to study and manage the abiotic factors which involve such favorable biocenotic relationships.

Acknowledgements The study was supported by a grant from the EC Project STD 3 n° TS3 * CT92-0098: Biocontrol of damaging root-knot nematode (*Meloidogyne* spp.) pests of staple food and cash crops by including suppressive soils with the bacterial parasite *Pasteuria penetrans*. The authors are grateful to Dr D.L. Trudgill (SCRI, Dundee, Scotland) for reviewing the manuscript.

Literature cited

AFOLABI P., DAVIES K. G. and O'SHEA P. S., 1995. The electrostatic nature of the spore of *Pasteuria penetrans*, the bacterial parasite of root-knot nematodes. *J. appl. Bacteriol.*, 79: 244-249.

- BARTOLI F., BURTIN G. and HERBILLON A. J., 1990. Disaggregation and clay dispersion in oxisols. Na-resin, a recommended methodology for determining particule size distribution of oxisols rich in gibbsite. Trans. 14th Congr. Int. AISS, Kyoto, Japan, vol. 4: 449-450. 12-18 August, 1990.
- BIRD A. F., BONIG I. and BACIC A., 1989. Factors affecting the adhesion of microorganisms to the surfaces of plant-parasitic nematodes. *Parasitology*, 98: 155-164.
- BROWN S. M. and SMART G. C., 1985. Root penetration by *Meloidogyne incognita* juveniles infected with *Bacillus penetrans*. *J. Nematol.*, 17: 123-126.
- DAVIES K. G., AFOLABI P. and O'SHEA P., 1996. Adhesion of *Pasteuria penetrans* to the cuticle of root-knot nematodes (*Meloidogyne* spp.) inhibited by fibronectin: a study of electrostatic and hydrophobic interactions. *Parasitology*, 112: 553-559.
- DAVIES K. G. and DANKS C., 1992. Interspecific differences in the nematode surface coat between *Meloidogyne incognita* and *M. arenaria* related to the adhesion of the bacterium *Pasteuria penetrans*. *Parasitology*, 105: 475-480.
- DAVIES K. G. and DANKS C., 1993. Carbohydrate/protein interactions between the cuticle of infective juveniles of *Meloidogyne incognita* and spores of the obligate hyperparasite *Pasteuria penetrans*. *Nematologica*, 39: 53-64.
- DAVIES K. G., KERRY B. R. and FLYNN C. A., 1988. Observations on the pathology of *Pasteuria penetrans* a parasite of the root-knot nematodes. *Ann. appl. Biol.*, 112: 491-501.
- DAVIES K. G., ROBINSON M. P. and LAIRD V., 1992. Proteins involved in the attachment of a hyperparasite *Pasteuria penetrans* to its plant-parasitic nematode host, *Meloidogyne incognita*. *J. Invert. Pathol.*, 59: 18-23.
- FELLER C., BURTIN G., GARARD B. and BALESIDENT J., 1991. Utilisation des résines sodiques et des ultrasons dans le fractionnement granulométrique de la matière organique des sols. Intérêts et limites. *Science du Sol*, 29: 77-93.

- HIMMELHOCH S., ORION D. and ZUCKERMAN B. M., 1979. Partial characterization of the cuticle surface of *Meloidogyne javanica* females. *J. Nematol.*, 11: 358-362.
- HOFFMANN-HERGARTEN S. and SIKORA R. A., 1995. Influence of seed pelleting technology on the rhizobacteria *Pseudomonas fluorescens* and studies on antagonistic activity towards *Heterodera schachtii*. *Nematologica*, 41: 309-310.
- IBRAHIM S. K., 1991. Distribution of carbohydrates on the cuticle of several developmental stages of *Meloidogyne javanica*. *Nematologica*, 37: 275-284.
- MARTIN J. K., 1975. Comparison of agar media for counts of viable soil bacteria. *Soil Biol. Biochem.*, 7: 401-402.
- OOSTENDORP M., DICKSON D. W. and MITCHELL D. J., 1991. Population development of *Pasteuria penetrans* on *Meloidogyne arenaria*. *J. Nematol.*, 23: 58-64.
- ROZYCKI H., 1987. Sugar productions by bacteria and actinomycetes isolated from soil, rhizosphere and mycorrhizosphere of Pine (*Pinus sylvestris* L.). *Acta Microbiol. Polonica*, 36: 93-99.
- SEINHORST J. W., 1950. De betekenis van de toestand von de grond voor het optreden van aanstasting door het stengelaaltje (*Ditylenchus dipsaci* (Kühn) Filipjev). *Tijdschr. Plantenz.*, 56: 292-349.
- STIRLING G. R., SHARMA R. D. and PERRY J., 1990. Attachment of *Pasteuria penetrans* to the root-knot nematode *Meloidogyne javanica* in soil and its effects on infectivity. *Nematologica*, 36: 246-252.
- ZAKI M. J. and MAQBOOL M. A., 1992. Effect of spore concentrations of *Pasteuria penetrans* on the attachment of *Meloidogyne* larvae and growth of okra plants. *Pak. J. Nematol.*, 10: 69-73.