

Istituto di Nematologia Agraria del C.N.R. - 70126 Bari, Italy
Istituto di Entomologia Agraria dell'Università - 80055 Portici - Napoli, Italy

OBSERVATIONS ON THE MALE OF *XIPHINEMA ITALIAE*

by

F. LAMBERTI and F. P. D'ERRICO

Males of *Xiphinema italiae* Meyl have been reported from France and Israel (Martelli *et al.*, 1966) and Sardinia, Italy (Prota *et al.*, 1971), but never from the Italian peninsula where this species is widespread in the southern regions (Martelli and Lamberti, 1967; Roca and Lamberti, 1978).




Fig. 1 - Anterior (a) and posterior (b) regions of the male of *Xiphinema italiae* found at Portici, Naples.

Table I - *Morphometric characters of Xiphinema italiae*.

C h a r a c t e r s	L o c a l i t y						
	Portici (Naples)		Avellino		France (Martelli <i>et. al.</i> , 1966)	Israel (<i>♂</i>)	
	♀ ♀	♂	♀ ♀	♂	♂	♂ ♂	
n	3	1	3	1	1	3	2
L mm	3(2.9-3.1)	2.9	2.9(2.7-3.1)	2.6	2.4	3.3(3.2-3.4)	2.8-3.2
a	88(86-91)	77	81(76-88)	80	86	87(79-101)	84-116
b	7.5(6.8-8.3)	8.3	7.5(7.2-7.7)	7.5	6.4	8.4(7.9-9.2)	7.2-8.7
c	38(37-40)	60	37(34-40)	52	50	48(47-50)	67-72
c'	3.5(3.3-3.7)	1.6	3.4(3.0-3.7)	1.9	2	2.3(2.1-2.5)	1.6
V	45(43-48)	—	46 45 47	—	—	—	—
Odontostyle μm	97(93-100)	96	95(92-98)	92	93	104(102-108)	99-103
Odontophore μm	60(59-62)	62	59 56-60	57	56	64(63-65)	58-66
Oral aperture to guiding ring μm	93(89-96)	95	86(83-89)	84	76	98(95-102)	91-97
Tail μm	78(78-79)	49	79 67-87	51	—	—	—
Spicules μm	—	52	—	43	48	48(46-50)	42-48
Number of ventromedian supplements in addition to the adanal pair	—	3	—	4	2	1	5

A male specimen of the nematode was recently found in each of two populations of *X. italiae* collected respectively at Portici (Naples), from the rhizosphere of mulberry (*Morus alba* L.) trees on which it seems to reproduce actively (Fig. 1), and at Avellino, in the rhizosphere of grapevine.

The morphometric characters of female specimens belonging to these two populations do not differ from those of other Italian populations (Martelli and Lamberti, 1967) (Tab. I), but having noticed some differences between males, it has been thought useful to report them (Tab. I), to contribute to the knowledge of the variability of this species.

LITERATURE CITED

- MARTELLI G. P., COHN E. and DALMASSO A., 1966 - A redescription of *Xiphinema italiae* Meyl, 1953 and its relationship to *Xiphinema arenarium* Luc et Dalmaso, 1963 and *Xiphinema conurum* Siddiqi, 1964. *Nematologica*, 12: 183-194.
- MARTELLI G. P. and LAMBERTI F., 1967 - Le specie di *Xiphinema* Cobb, 1913 trovate in Italia e commenti sulla presenza di *Xiphinema americanum* Cobb (Nematoda, Dorylaimoidea). *Phytopath. medit.*, 6: 65-85.
- PROTA U., LAMBERTI F., BLEVE T. and MARTELLI G. P., 1971 - I Longidoridae (Nematoda, Dorylaimoidea) dei vigneti sardi. *Redia*, 52: 601-618.
- ROCA F. and LAMBERTI F., 1978 - Longidoridae of Italian vineyards. I. The genus *Xiphinema*. Proceedings of the VI Conference on Virus and Virus Diseases of the Grapevine (13-21 Sept., 1978) Cordoba, Spain, pp. 251-254.

Accepted for publication on 4 December 1979.