

EFFICACY OF FOUR PLANT EXTRACTS ON NEMATODES ASSOCIATED WITH PAPAYA IN SINDH, PAKISTAN

A. Khan^{1*}, M. Sayed¹, S.S. Shaukat² and Z.A. Handoo³

¹Crop Diseases Research Institute, PARC, University of Karachi, Karachi-75270

²Department of Botany, University of Karachi, Karachi-75270, Pakistan

³USDA, ARS, Nematology Laboratory, Beltsville, MD 20705, USA

Summary. This investigation attempts to evaluate the effect of ethanolic extracts of four plant species, *Azadirachta indica* (neem), *Withania somnifera* (ashwagandha), *Tagetes erecta* (marigold) and *Eucalyptus citriodora* (eucalyptus), against nematodes associated with papaya (*Carica papaya*) and to assess their influence on papaya growth and yield. While fresh shoot weight of papaya seedlings in pots was significantly increased by the plant extracts, in general, the fresh root weight remained unaltered compared to the controls. The number of *Meloidogyne incognita* juveniles, root-knot index and the number of egg masses per root were remarkably reduced by all treatments. The population densities of *Meloidogyne incognita*, *Helicotylenchus multincinctus* and *Hoplolaimus indicus*, the main nematode species associated with papaya, were significantly reduced in the field by all four plant extracts. *Meloidogyne incognita* and *H. multincinctus* were most affected by *Withania*, *H. indicus* by *Tagetes*. The papaya yield was increased by the treatments in the following order: *Withania somnifera* > carbofuran > *Azadirachta indica* > *Tagetes erecta* > *Eucalyptus citriodora*. The possible mechanisms of nematocidal effects of the plant extracts are discussed.

Key words: *Azadirachta indica*, control, *Carica papaya*, *Withania somnifera*, *Tagetes erecta*, *Eucalyptus citriodora*.

Plant-parasitic nematodes, particularly root-knot nematodes, are widely distributed and cause significant yield losses in a wide range of crops (Sasser and Freckman, 1987). It has been estimated that worldwide overall annual crop yield losses due to plant-parasitic nematodes are approximately 78 billion dollars (Barker *et al.*, 1994).

A preliminary survey of parasitic nematodes associated with papaya (*Carica papaya* L.) in Sindh province, Pakistan, disclosed that the three common species were: *Meloidogyne incognita* (Kofoid *et White*) Chitw., *Helicotylenchus multincinctus* (Cobb) Golden and *Hoplolaimus indicus* Sher (Khan *et al.*, 2007). *Meloidogyne* spp. are the most widespread parasitic nematodes, causing root-knot disease of numerous crops and responsible for 70% of the total loss due to nematodes (Sasser and Freckman, 1987). *Meloidogyne incognita* parasitizes a number of crops in the tropics and semi-tropics (Williamson and Hussey, 1996). *Helicotylenchus* spp. are ectoparasites of roots but may also occur in cortical tissue (Siddiqui, 2000). Similarly, *Hoplolaimus indicus* is also an important and widely distributed ectoparasitic nematode, causing root damage and consequently limiting crop yield.

Nematode control is largely based on synthetic nematicides, which besides being costly present potential risk to non-target organisms. In the search for more benign acceptable alternatives to chemicals, the possibilities are being investigated of exploiting nematode-an-

tagonistic plants for the management of plant parasitic nematodes (Chitwood, 2002; Akhtar, 2004). Certain cover crops as green manure have been shown to improve soil fertility as well as suppress plant parasitic nematodes (Barker and Koenig, 1998; McSorley, 1999; Wang *et al.*, 2002, 2003). Soil amendment with leaves of many plants has been shown to reduce the rhizosphere population of *Meloidogyne* spp. (Wang *et al.*, 2004; Tsay *et al.*, 2004; Wani, 2006). Shaukat and Siddiqui (2001a) found that aqueous extracts of various weeds from Karachi, Pakistan, caused significant mortality of *M. javanica* (Treub) Chitw. juveniles *in vitro*. Shaukat *et al.* (2004) demonstrated that extracts of certain weeds are able to cause substantial mortality of plant parasitic nematodes thereby improving plant growth and yield. The objectives of this study were to test 1) the effects of ethanolic extracts of four plant species, containing known toxins, on *M. incognita* soil population density, root-knot disease intensity and growth parameters of papaya seedlings, and 2) to test the effects of these extracts on the parasitic nematodes associated with papaya and on yield of papaya.

MATERIALS AND METHODS

Pot experiment. Papaya cv. Bengali seeds were grown in 20-cm diam. plastic-pots filled with 2 kg of solarized sandy loam soil. Three weeks after germination, single seedlings were transferred to 9-cm diam. plastic pots containing 250 g of solarized sandy loam soil. Pots were randomized on a glasshouse bench at 28-30 °C and relative humidity of 55-60%. One week later, the soil in each pot

* Corresponding author e-mail: dralykhan@hotmail.com

was inoculated with 200 three-day-old second stage juveniles (J_2) of *M. incognita* obtained from a pure culture raised on tomato; and one hour later the pots were treated with the plant extracts of *Azadirachta indica* A. Juss., *Withania somnifera* Dun., *Tagetes erecta* L. and *Eucalyptus citriodora* Hook., at 25 ml of extract per pot.

The extracts were prepared as follows. One hundred g of air-dried plant material was plunged into 400 ml of 80% ethanol at 50 °C, stirred well and soaked for 24 h. This material was then reduced to a gummy mass in a Rotary evaporator (Buchi Rotavapor R-200) and dissolved in 100 ml distilled water. The extract was used for the treatments.

In addition, there were positive and negative controls for different amendments. The nematicide carbofuran (commercial product, Furadan a.i. 3 percent) was used

at 2 g (dissolved in 25 ml water) per pot.

Each treatment and control were replicated four times. Sixty days after nematode inoculation, the plant in each pot was uprooted and the soil particles around the roots gently shaken off. The nematode juveniles in the soil were extracted by a sieving and centrifugation technique (Barker, 1985). The eggs/egg masses of *M. incognita* from the roots were extracted by shaking the roots in a 0.5% sodium hypochlorite solution for 3 minutes (Hussey and Barker, 1973). Egg-masses were kept in a Petri dish containing distilled water and placed in an incubator at 28 °C in the dark. J_2 were collected on the third day. Root-knot and egg mass indices were rated according to the scale of Hadisoeganda and Sasser (1982). Fresh and dry weights of shoots and roots were recorded.

Table I. Effects of different treatments on plant growth components of papaya seedlings and reproduction of *Meloidogyne incognita* in pots inoculated or not with 200 second stage juveniles (J_2) of the nematode.

Treatment	Fresh weight (g)		Dry weight (g)		Nematode J_2 / 250 g soil	Root-knot index (0-5)	Nematode egg masses/ root	Eggs/ egg mass
	Shoot	Root	Shoot	Root				
Control positive	1.19	0.27	0.11	0.052	286.5	3	8.5	20
Control negative	1.62	0.38 n.s.	0.14 n.s.	0.067 n.s.	0	0	0	0
Carbofuran + 200 J_2	1.58	0.35 n.s.	0.10 n.s.	0.06 n.s.	0	0	0	0
<i>Tagetes erecta</i> + 200 J_2	1.46	0.33 n.s.	0.13 n.s.	0.067 n.s.	2	0.25	0.25	0.50
<i>Azadirachta indica</i> + 200 J_2	1.58	0.28 n.s.	0.12 n.s.	0.04 n.s.	4	0.25	1	0.50
<i>Withania somnifera</i> + 200 J_2	1.92	0.36 n.s.	0.13 n.s.	0.035 n.s.	0	0	0	0
<i>Eucalyptus citriodora</i> + 200 J_2	1.43 n.s.	0.27 n.s.	0.10 n.s.	0.04 n.s.	4.25	0.25	0.50	1
Carbofuran	1.49	0.38 n.s.	0.16 n.s.	0.080 n.s.	0	0	0	0
<i>Tagetes erecta</i>	1.66	0.37 n.s.	0.16 n.s.	0.075 n.s.	0	0	0	0
<i>Azadirachta indica</i>	1.62	0.45	0.13 n.s.	0.062	0	0	0	0
<i>Withania somnifera</i>	2.10	0.47	0.17 n.s.	0.09 n.s.	0	0	0	0
<i>Eucalyptus citriodora</i>	1.50	0.35 n.s.	0.12 n.s.	0.065 n.s.	0	0	0	0
L.S.D. _{0.05}	0.27	1.18	0.07	0.03	28	1.7	4	1.6

Values are averages of four replicates.

Significance is given compared to positive controls.

Control positive = control in conjunction with inoculation of *M. incognita* juveniles.

Control negative = control without inoculation of *M. incognita* juveniles.

Field trial. A farmer's field in Malir, Sindh, was chosen for the trial. Papaya trees (var. Bengali) were growing in rows and were apparently free from diseases. Summer temperature in the area varies from 30 to 35 °C while the winter average is 18 °C. The average annual rainfall is 200 mm but in 2006 it was 116 mm. Soil samples were collected in Oct. 2006 (fifteen days prior to treatment) from near the tree trunks at a depth of 5–25 cm, taking 6–8 cores per tree that were pooled to form about 1.5 kg of composite sample. Soils from six rows of papaya were sampled with three samples in each row. Population density of all the stylet-bearing nematodes associated with papaya were determined by a sieving and decantation and modified Baermann funnel technique (Southey, 1986). Five ml aliquots (fifteen replicates) of nematode suspension were used for nematode counts and value converted to number of nematodes per 200 ml of soil sample. The three stylet-bearing nematodes (see below) comprised 88% of the total nematode population and the rest were free-living (non-pathogenic). The initial populations were 45 ± 3.9 *Meloidogyne incognita*, 55 ± 15.3 *Helicotylenchus multicinctus* and 30 ± 7.2 *Hoplolaimus indicus*.

The leaves of neem (*A. indica*), ashwagandha (*W. somnifera*), eucalyptus (*E. citriodora*) and marigold (*T. erecta*) were air-dried in the shade and ground to coarse powder from which the plant extracts were prepared as described earlier (see pot experiment). Four hundred ml of each extract was applied to 2 m of row of the papaya trees at 20 cm depth by mixing the extract in the soil using a spade. The extract was uniformly broadcast in the furrow and the application rate corresponded to approximately 1,300 kg plant material in the form of extract per hectare of papaya field. Each treatment and controls were replicated three times. The plant spacing was 1.5 m between rows and 1.2 m along the row. Weeding was done manually and cow manure was used as fertilizer. No fungal disease was observed and insect populations were at a low level and not problematic. For comparison of the potential phytonematicides, the chemical nematicide carbofuran (commercial product, Furadan) was applied at a rate of 15 kg/ha. Untreated trees were kept as controls. Twelve months after the application, soil samples were collected from a depth of 5–25 cm with a soil auger to determine nematode populations. Each final sample was a composite of 6–8 cores. For yield determination, the first picking started after 11 months, and a total of four pickings were taken into account over a period of three weeks. There were four replicates for each observation.

Data were subjected to either one-way analysis of variance (ANOVA) (pot experiment) or the factorial analysis of variance (FANOVA) (field experiment). Each variable was treated separately in the case of ANOVA. The post-hoc tests performed included Fisher's least significant difference (LSD) at $P = 0.05$ and Dunnett's test for comparing control with treatment means (Zar, 1999).

RESULTS

Pot experiment. Growth components were slightly elevated in the negative control over the positive control. Shoot fresh weight of papaya was significantly increased in all treatments except *Eucalyptus citriodora* + 200 nematode J₂ as compared to the positive control (Table I).

Differences in root fresh weight from the positive control were not significant in any treatments except for *A. indica* and *W. somnifera* without the nematode. Shoot dry weight also was not significantly different in all treatments compared to the positive control. Moreover, root dry weight was not significantly different in any treatments except for *A. indica* (without the nematode). Compared to negative controls, the growth parameters of treatments (either plant extract alone or in conjunction with the nematode), in general, did not exhibit any signif-




Fig. 1. Effect of different treatments, under field conditions, on nematode soil population densities of: A, *Meloidogyne incognita*; B, *Helicotylenchus multicinctus*; C, *Hoplolaimus indicus*. IN = Initial population density; CO = Control; F = Furadan; EU = *Eucalyptus citriodora*; TA = *Tagetes erecta*; AZ = *Azadirachta indica*, and WT = *Withania somnifera*. LSDs: nematodes = 5.05; treatments = 7.15.

Table II. Effect of plant extract treatments on the yield (kg/plant) of papaya, at different harvests, under field conditions.

Treatment	Harvest				Cumulative
	1	2	3	4	
Carbofuran	1.388	1.510	1.482	1.507	5.887
<i>Azadirachta</i>	1.291	1.366	1.430	1.652	5.739
<i>Withania</i>	1.304	1.328	1.646	1.774	6.052
<i>Tagetes</i>	1.121	1.245	1.368	1.504	5.238
<i>Eucalyptus</i>	1.132	1.386	1.260	1.391	5.169
Control	1.040	1.230	1.114	1.258	4.642

LSD_{0.05} (treatments) = 0.086
LSD_{0.05} (harvests) = 0.115

icant change. Total number of J₂/250 g soil, root-knot index, number of egg masses/root and mean number of eggs/egg masses were significantly reduced in all treatments compared to the positive control.

Field experiment. The effects of the treatments were significant ($F = 143.12$, $P < 0.001$). The treatment with *W. somnifera* was the most effective, followed by carbofuran and *T. erecta*. Significant differences were found for nematode species ($F = 15.47$, $P < 0.001$). While the root-knot nematode and *Helicotylenchus multicinctus* were most affected by ashwagandha (*W. somnifera*), the population of *Hoplolaimus indicus* was affected by *T. erecta*, carbofuran and *W. somnifera* extract (Fig. 1A-C). Interaction of treatments \times nematode species was found to be significant ($F = 19.15$, $P < 0.001$).

The yield of papaya was significantly increased ($P < 0.001$) by carbofuran and the four plant extracts tested (Table II). The yield was increased over the controls in the order *W. somnifera* > carbofuran > *A. indica* > *T. erecta* > *E. citriodora*. Harvests also differed significantly ($P < 0.05$) and the yield tended to increase from first to last harvest. The interaction of treatments \times harvest was also found to be significant ($P < 0.01$).

DISCUSSION

The pot and field experiments demonstrated the nematicidal potential of the four plant extracts tested, especially ashwagandha and neem, that also resulted in increased plant growth and yield of papaya.

Several mechanisms may be involved in the suppression of nematode soil populations and enhancement of plant growth and yield. Plant extracts often contain phenolic compounds, organic acids, terpenes and terpenoids, coumarin-like compounds and other secondary metabolites (Muller and Gooch, 1982; Insunza *et al.*, 2001; Shaukat *et al.*, 2004). Many of these com-

pounds, such as phenolics, have been demonstrated to have nematicidal activity (Shaukat *et al.*, 2004). The nematicidal activity of neem can be attributed to one or more of the biologically active principles it contains, including azadirachtine, nimbine, nimbidine, thionemone, kemferol, etc., that are known to possess nematicidal properties (Rossner and Zebitz, 1987; Abbasi *et al.*, 2004). *Eucalyptus* spp. possess complex mixtures of plant secondary metabolites, including terpenoids, cyanogenic glycosides, tannins, flavonoids, long-chain ketones, phloroglucinols and leaf-oil 1-8-cineole eucalyptol (Boland *et al.*, 1991; Brophy and Southwell, 2002). *Withania somnifera* contains inhibitory withanoloids and glyco-withanoloids that are known to be highly effective against *M. javanica* (Goel *et al.*, 2005). Finally, leaves and flowers of *Tagetes erecta* contain some inhibitory essential oils (Oguvawande and Olawore, 2006; Tomava *et al.*, 2005) that possess strong nematicidal activity (Wang *et al.*, 2002; Evenhuis *et al.*, 2004; Tsay *et al.*, 2004). In addition, the application of plant extracts or other control agents can also alter the indigenous bacterial or fungal communities of the rhizosphere or endorhiza in a manner that leads to enhanced soil suppressiveness to nematodes or other soil-borne plant pathogens (Vargas-Ayala *et al.*, 2000; Shaukat and Siddiqui, 2001b).

The use of plant material for the management of nematode populations is apparently effective and environmentally friendly compared to synthetic nematicides. The pot experiment was a preliminary test and the rates of plant extracts used could be considered rather large, although the quantity of secondary metabolites present in the extracts, was probably more or less equivalent to the rates used for chemical nematicides. The amount of plant material used in the field experiment are comparable to those used as organic soil amendments by other workers, and provide effective nematode control. However, whether a smaller amount would still be effective needs further investigation.

ACKNOWLEDGEMENTS

The first author is thankful to Agriculture Linkage Programme for financial support.

LITERATURE CITED

- Abbasi P.A., Riga E., Conn K.L. and Lazarovits G., 2004. Effect of neem cake soil amendment on reduction of damping-off severity and population densities of plant-parasitic nematodes and soilborne plant pathogens. *Canadian Journal of Plant Pathology*, 27: 38-45.
- Akhtar M., 2004. Current options in integrated management of plant-parasitic nematodes. *Integrated Pest Management Review*, 2: 187-197.
- Barker K.R., 1985. Nematode extraction and bioassays. Pp. 19-35. In: An Advanced Treatise on *Meloidogyne*. Vol. II Methodology. (Barker K.R., Carter C.C. and Sasser J.N., eds). North Carolina State University Press, Raleigh, USA., pp.223.
- Barker K.R. and Koenig S.R., 1998. Developing sustainable systems of nematode management. *Annual Review of Phytopathology*, 36: 165-205.
- Barker K.R., Hussey R.S., Krusberg L.R., Bird G.W., Dunn R.A., Ferris H., Ferris V.R., MacGuidwin A.E., Riddle D.R., Roberts P.A. and Schmitt D.P., 1994. Plant and soil nematodes: Social impact and focus for the future. *Journal of Nematology*, 26: 127-137.
- Boland D.J., Brophy J.J. and House A.R.N., 1991. *Eucalyptus leaf-oils-use, chemistry, distribution and marketing*. Corporate Publications, Brisbane, Australia, pp273.
- Brophy J.J. and Southwell I.A., 2002. *Eucalyptus* chemistry. Pp. 102-160. In: *Eucalyptus* (Coppin J.J.W., ed.). Taylor & Francis, London & New York.
- Chitwood D.J., 2002. Phytochemical based strategies for nematode control. *Annual Review of Phytopathology*, 87: 250-258.
- Evenhuis A., Korthalis G.W. and Molendijk L.P.G., 2004. *Tagetes patula* as an effective catch crop for long-term control of *Pratylenchus penetrans*. *Nematology*, 6: 877-881.
- Goel S.R., Madan V.K., Sharma K.K. and Nandal S.N., 2005. Nematicidal activity of various medicinal and aromatic plants under *in vitro* conditions. National Symposium on Recent Advances and Research Priorities in Indian Nematology, 9-10 December, IARI, New Delhi, India, p.23..
- Hadisoeganda W.W. and Sasser J.N., 1982. Resistance of tomato, bean, southern pea and garden pea cultivars to root-knot nematode based on host-suitability. *Plant Diseases*, 66: 145-150.
- Hussey R.S. and Barker K.R., 1973. A comparison of methods of collecting inocula of *Meloidogyne* spp. including a new technique. *Plant Disease Reporter*, 57: 1025-1028.
- Insunza V., Aballay E. and Macaya J., 2001. *In vitro* nematicidal activity of aqueous plant extracts on Chilean populations of *Xiphinema americanum sensu lato*. *Nematropica*, 31: 47-54.
- Khan A., Sayed M. and Shaukat S.S., 2007. Nematodes associated with papaya in Sindh (Pakistan). *Sarhad Journal of Agriculture*, 23: 145-147.
- McSorley R., 1999. Host suitability of potential cover crops for root-knot nematodes. *Journal of Nematology*, 31(S): 619-623.
- Muller R. and Gooch P.S., 1982. Organic amendments in nematode control and examination of the literature. *Nematropica*, 12: 319-326.
- Oguyawande I.A. and Olawore N.O., 2006. Essential oils from the leaves and flowers of African marigold, *Tagetes erecta* L. *Journal of Essential Oil Research*, 7: 30-39.
- Rossner J. and Zebitz C.P.W., 1987. Effect of neem products on nematodes and growth of tomato (*Lycopersicon esculentum*). Pp. 611-621. In: Natural Plants: the Neem Tree (*Azadirachta indica* A. Juss.) and Other Tropical Plants (Schmutter H. and Ascher K.R.S., eds). Deutsche Gesellschaft Fur Technische Zusammenarbes, Eschborn, Germany.
- Sasser J.N. and Freckman D.W., 1987. A world perspective in nematology: the role of society. Pp. 7-1. In: Vistas on Nematology (Veech J.A. and Dickson D.W., eds). Society of Nematologists, Hyattsville, Maryland, USA.
- Shaukat S.S. and Siddiqui I.A., 2001a. Effect of some weed extracts on mortality and egg hatching of *Meloidogyne javanica* (Treb.) Chitwood. *Pakistan Journal of Biological Sciences*, 4: 1251-1252.
- Shaukat S.S. and Siddiqui I.A., 2001b. *Lantana camara* in the soil changes the fungal community structure and reduces impact of *Meloidogyne javanica* on mungbean. *Phytopathologia Mediterranea*, 40: 245-252.
- Shaukat S.S., Siddiqui I.A. and Zarina B., 2004. Effects of some common weeds from Pakistan on plant-parasitic nematodes *in vitro* and population densities and survival of *Meloidogyne incognita* in okra and brinjal. *Nematologia Mediterranea*, 32: 111-115.
- Siddiqui M.R., 2000. *Tylenchida parasites of plants and insects*. 2nd ed. CABI Publishing, Wallingford, UK, pp.848.
- Southey J.F. 1986. Laboratory methods for work with plant and soil nematodes. Ministry of Agriculture, Fisheries and Food, HMSO, London. pp 202.
- Tomava B.S., Waterhouse J.S. and Doberki J., 2005. The effect of fractionated *Tagetes* oil volatiles on aphid reproduction. *Entomologia Experimentalis et Applicata*, 115: 153-159.
- Tsay T.T., Wu S.T. and Lin Y.Y., 2004. Evaluation of Asteraceae plants for control of *Meloidogyne incognita*. *Journal of Nematology*, 36: 36-41.
- Vargas-Ayala R., Rodriguez-Kabana R., Morgan-Jones G., McInory J.A. and Kloepper J.W., 2000. Shifts in soil microflora induced by velvet bean (*Mucuna deeringiana*) in cropping systems to control root-knot nematodes. *Biological Control*, 17: 11-22.
- Wang Q., Bryan H., Klassen W., Li Y., Codallo M. and Abul-Baki A., 2002. Improved tomato production with summer cover crops and reduced irrigation rates. *Proceedings of Florida State Horticulture Society*, 115: 202-207.
- Wang Q., Klassen W., Handoo Z., Abul-Baki A., Bryan H. and Li Y., 2003. Influence of summer cover crops on soil nematodes in a tomato field. *Proceedings of Soil and Crop Science Society of Florida*, 62: 86-91.
- Wang K.H., McSorley R. and Gallaher R.N., 2004. Effect of *Crotalaria juncea* amendment on squash infected with *Meloidogyne incognita*. *Journal of Nematology*, 36: 290-296.

Wani A.H., 2006. Management of root-knot nematode, *Meloidogyne incognita* on okra and lentil by soil amendment with oil cakes and leaves of different plants. *Nematologia Mediterranea*, 34: 83-87.

Williamson V.W. and Hussey R.S., 1996. Nematode pathogenesis and resistance in plants. *Plant Cell*, 8: 1735-1745.
Zar J.H., 1999. *Biostatistical Analysis* – 4th ed. Prentice Hall, Englewood, Cliffs, New Jersey, USA., pp. 718.

Accepted for publication on 18 April 2008.