

REVIEW

ROTYLENCHULUS RENIFORMIS IN COTTON: CURRENT METHODS OF MANAGEMENT AND THE FUTURE OF SITE-SPECIFIC MANAGEMENT

Scott R. Moore* and K. S. Lawrence

Auburn University, Department of Entomology and Plant Pathology, 209 Life Science Building, Auburn University, AL 36849 USA; Corresponding author: lawrekk@auburn.edu

ABSTRACT

Moore, S. R., and K. S. Lawrence. 2012. *Rotylenchulus reniformis* in cotton: current methods of management and the future of site-specific management. *Nematropica* 42:227-236.

The reniform nematode, *Rotylenchulus reniformis*, is currently one of the most limiting factors to cotton production in the United States. With no available commercial host plant resistance, options for management of *R. reniformis* are limited to the use of rotations with non-hosts and the use of nematicides, each of which varies greatly in cost-savings and effectiveness. Multiple research groups are currently pursuing the goal of site-specific management for *R. reniformis* in cotton. Site-specific application is used for a wide variety of agricultural practices, and successful programs for other species of nematodes in cotton, such as *Meloidogyne incognita* and *Hoplolaimus columbus*, are currently in use. Within this manuscript, future possibilities for the use of site-specific management for *R. reniformis* in cotton as well as potential limitations of current techniques are discussed.

Key words: Cotton, crop rotation, *Rotylenchulus reniformis*, site-specific management.

RESUMO

Moore, S. R., and K. S. Lawrence. 2012. *Rotylenchulus reniformis* en algodón: métodos actuales de manejo y futuro del manejo sitio-específico. *Nematropica* 42:227-236.

El nematodo reniforme *Rotylenchulus reniformis* es actualmente uno de los factores más limitantes en la producción de algodón en los Estados Unidos de América. Comercialmente no se encuentra disponible resistencia en plantas hospederas, por lo tanto el manejo de *R. reniformis* se limita al uso de rotaciones con plantas no hospederas o al uso de nematicidas, los cuales varían considerablemente en costo y efectividad. Actualmente múltiples grupos de investigación están en busca de estrategias de manejo sitio-específicas para *R. reniformis* en algodón. Estas estrategias son usadas en una gran variedad de prácticas agrícolas, y actualmente son empleadas en exitosos programas para otras especies de nematodos en algodón, tales como *Meloidogyne incognita* y *Hoplolaimus columbus*. En este manuscrito se discuten las futuras posibilidades para el uso de estrategias sitio-específicas de *R. reniformis* en algodón, así como potenciales limitaciones de las técnicas actuales.

Palabras clave: algodón, rotación de cultivos, *Rotylenchulus reniformis*, manejo sitio-específico.

Cotton, *Gossypium hirsutum* Linnaeus, is one of the most economically important crops in the United States. In 2010, cotton was grown in 17 states with 11 million acres devoted to cotton production valued at more than \$7.3 billion (USDA-NASS, 2011).

The reniform nematode, *Rotylenchulus reniformis* Linford & Oliveira, is a semi-endoparasite of roots that occurs in tropical and sub-tropical regions (Robinson *et al.*, 1997) and is a major pathogen affecting U. S. cotton. Currently,

R. reniformis can be found in 11 of the 17 cotton producing states and is estimated to have caused a loss of nearly 2% annually in the past decade (Blasingame *et al.*, 2002 – 2012).

Rotylenchulus reniformis is easily introduced into cotton fields on contaminated equipment and other means of soil transport. Once there, it can be spread throughout the field by tillage and water flow (Monfort *et al.*, 2008; Moore *et al.*, 2011a); however, in no-till systems, *R. reniformis* can spread independently both horizontally

and vertically (Moore *et al.*, 2010a). Vertical distribution has been well documented at depths of up to 1.5 m (Lee *et al.*, 2002; Moore *et al.*, 2010a; Robinson *et al.*, 2005a; Westphal & Smart, 2003; Westphal *et al.*, 2004), and populations below the plow layer can greatly affect cotton yields (Newman & Stebbins, 2002; Robinson *et al.*, 2005b).

Currently, there are no commercial cotton cultivars with resistance or consistent tolerance to *R. reniformis* (Usery *et al.*, 2005; Robinson, 2007). As such, management options for *R. reniformis* fall into two major categories: pesticides and crop rotation. There are many forms of pesticides available for the management of *R. reniformis*. Each varies in effectiveness and each has its limitations. Fumigants such as 1,3-dichloropropene (Telone II) and metam sodium (Vapam) are generally highly effective for management of *R. reniformis* (Kinloch & Rich, 2001; Koenning *et al.*, 2007; Lawrence *et al.*, 1990; Rich & Kinloch, 2000). They are often limited by cost, high risk to applicators, special application equipment, soil texture, and temperature and moisture requirements.

An assortment of granular pesticides have been proven effective for the management of *R. reniformis*, including aldicarb (Temik 15G) (Lawrence *et al.*, 1990; Lawrence & McLean, 2000, Rich & Kinloch, 2000), fenamiphos (Nemacur) (Koenning *et al.*, 2007; Lawrence *et al.*, 1990), and terbufos (Counter) (Lawrence *et al.*, 1990). Of the granular pesticides, aldicarb has been the most widely used in cotton production, and its continual use has resulted in reports of enhanced degradation by soil microbes thus decreasing its overall efficacy (Lawrence *et al.*, 2005). Furthermore, the future of this pesticide is currently unknown due to the discontinuance of its production (Bayer CropScience, 2010). Similarly, fenamiphos is no longer labeled for use in the United States (EPA, 2002), and terbufos is not currently labeled for use in cotton production.

Seed applied pesticides such as abamectin and thiodicarb have recently become widely used in cotton production as a part of Avicta Complete Cotton and Aeris Seed Applied System, respectively, and have been reported to provide adequate management of *R. reniformis* (Faske & Starr, 2006; Lawrence & Lawrence, 2007). Their protection of the root is limited (Faske & Starr, 2007) as is their ability to provide adequate protection against high populations of *R. reniformis* (Moore *et al.*, 2010b).

Oxamyl (Vydate C-LV) is a foliar applied

pesticide that also provides adequate management of *R. reniformis*, often in conjunction with previously mentioned pesticides (Baird *et al.*, 2000; Lawrence & McLean, 2000), but has been reported to be less effective in dry conditions (Koenning *et al.*, 2007). Additional options for *R. reniformis* management in the form of biological organisms, such as *Bacillus firmus* (Poncho/VOTiVO) and *Paecilomyces lilacinus* strain 251 (Nemout) as seed applied formulations (Castillo *et al.*, 2011), have been reported to have efficacy against *R. reniformis*. Furthermore, there are multiple known nematophagous fungi with high levels of effectiveness in greenhouse studies (Wang *et al.*, 2004; Castillo *et al.*, 2009) that could prove useful in the future. Overall, the number of pesticides for the management of *R. reniformis* is decreasing, resulting in increased challenges for producers.

Crop rotation to non-hosts, such as corn or peanuts or highly resistant varieties of soybean, is also an effective strategy for the management of *R. reniformis*. A one year rotation with corn and resistant soybean effectively increases cotton yields (Davis *et al.*, 2003; Moore *et al.*, 2010c); however, populations of *R. reniformis* quickly rebound to pre-rotational crop levels by mid-season. A two year or longer rotation with corn or resistant soybean or a one year or longer rotation with peanuts can result in *R. reniformis* populations remaining below current economic thresholds throughout the subsequent cotton crop (Stetina *et al.*, 2007; Moore *et al.*, 2010c). Many native weed species are host of *R. reniformis* to some degree and can confound the aforementioned positive effects of crop rotation if not properly controlled (Davis & Webster, 2005; Jones *et al.*, 2006; Lawrence *et al.*, 2008; Wang *et al.*, 2003).

The methods currently used to manage *R. reniformis* in cotton can be economically beneficial if utilized intelligently and with forethought. For a problem that is consistently increasing, further management strategies are needed. Site specific, or precision, management (SSM) is a concept that is increasingly utilized since being made possible by the integration of global positioning systems (GPS) technologies into agriculture. The use of SSM based on soil variability as a strategy to enhance the management of *R. reniformis* has developed into a subject of great interest in recent years. In this review, the current methods of zone delineation for SSM and their uses will be discussed along with the potential for use of known factors affecting *R. reniformis* and its

interaction with cotton. The pitfalls of SSM in regards to its use for *R. reniformis* management also will be addressed, as will an evaluation of the feasibility of using current methods of SSM for *R. reniformis*. Finally, we will determine what information is still required to facilitate a workable guideline for implementing SSM for *R. reniformis*.

The delineation of management zones for SSM based on soil variability has been a topic of research for decades. A management zone can be defined as a subregion of a field that expresses a homogeneous combination of yield limiting factors for which a single rate of a specific crop input is appropriate (Doerge, 1999). The development of management zones requires the use of some form of geostatistical analysis. There are many different methods of geostatistical analysis, both descriptive and predictive, that can be used alone or in combination, depending on the situation. Descriptive methods of geostatistical analysis allow for the detection and quantification of the major scales of spatial variability (Goovaerts, 1998). Examples of such descriptive methods include the experimental correlogram, which plots the estimated correlation coefficients of one variable as a function of the separation distance, and the experimental semivariogram, which plots the semivariances of ordered data versus distance (Goovaerts, 1998). Predictive methods are utilized in the estimation of soil properties at unsampled places between or near collected data points. Examples of predictive methods of geostatistical analysis include ordinary kriging, which estimates the value of an unsampled location as a linear combination of neighboring observations, and factorial kriging, which estimates and maps different sources of spatial variability identified by experimental semivariograms (Wackernagel 1988, 1995; Goovaerts, 1992).

Prescription maps began development based on soil type (Carr *et al.*, 1991) or topography (Fiez *et al.*, 1994). Further research has developed prescription maps on a collection of characteristics including soil type, soil color, topography, yield, aerial photos, and producer experience (Ostergaard, 1997; Fleming *et al.*, 2004). The use of soil apparent electrical conductivity (EC) has become one of the most frequently used methods of management zone delineation based on soil variability. Apparent electrical conductivity has been found to correlate highly with soil texture (Williams & Hoey, 1987). It also relates closely with a variety of other characteristics including:

cation exchange capacity and exchangeable Ca and Mg (McBride *et al.*, 1990), water content (Kachanoski *et al.*, 1988), soil organic C (Jaynes, 1996), herbicide behavior in soil (Jaynes *et al.*, 1994), depth to claypans (Kitchen *et al.*, 1999), and crop yield (Sudduth *et al.*, 1995; Heermann *et al.*, 1999).

The geostatistical analysis of soil properties and the subsequent delineation of management zones have proven effective in a variety of situations worldwide. Casa & Castrignano (2008) demonstrated the spatial relationships between soil and crop variables of durum wheat in Italy. Rab *et al.* (2009) utilized geostatistical modeling of plant-available water capacity and related soil properties to delineate management zones for the enhancement of grain yields in Australia. Liu *et al.* (2006) explored the possibilities of combining ordinary kriging with soil map-delineation to enhance the interpolation of soil properties in a paddy rice/sugarcane rotation in Taiwan. Lopez-Lozano *et al.* (2010) successfully linked leaf area index with soil properties for precision management of abiotic stress of corn in Spain. In the U. S., management zones based on soil characteristics have been used to predict grain yields (Fraisse *et al.*, 2001) and determine the risk of iron chlorosis in maize (Kyaw *et al.*, 2008).

The use of geostatistical analysis and management zone delineation also has recently been developed for the management of the Columbia lance nematode (*Hoplolaimus columbus*), the root-knot nematode (*Meloidogyne incognita*), and the ring nematode (*Criconebella* spp.) (Khalilian *et al.*, 2001; Khalilian *et al.*, 2002; Khalilian *et al.*, 2003; Monfort *et al.*, 2007; Ortiz *et al.*, 2007; Ortiz *et al.*, 2008; Wolcott *et al.*, 2004; Wolcott *et al.*, 2005). Khalilian *et al.* (2003) reported a 5% yield increase using either variable rate aldicarb or 1,3-dichloropropene for Columbia lance management with a 34% and 78% reduction of input, respectively. Monfort *et al.* (2007) observed that the combination of the initial populations of root-knot nematodes and the sand content of the soil explained 65%, 86%, and 83% of the variation in cotton yield over a three-year period, respectively. Similarly, Ortiz *et al.* (2007) observed that a model of root-knot nematode risk of a field over a specific threshold value could be produced through logistic regression using soil electrical conductivity as a predictor variable. Furthermore, it was determined that the use of variable rate application of nematicides could be effectively employed to manage root-knot

nematodes in cotton (Ortiz *et al.*, 2008).

Although there are several successful examples of site-specific management of nematodes, there are studies that address certain pitfalls of this technique. Wyse-Pester *et al.* (2002) conducted a study to determine the scale of sampling required to obtain correlated observations of density in order to reduce sampling costs for three species of nematodes on corn. The results of the study indicated that correlations between nematode density and soil attributes were inconsistent between field and species, and thus the cost of sampling was not reduced. Similarly, Evans *et al.* (2002) found that coarse sampling grids, which are required to make SSM a commercially viable option for the management of potato cyst nematodes (*Globodera pallida* and *G. rostochiensis*), are likely to produce misleading population distribution maps resulting in yield penalties. Farias *et al.* (2002) were able to construct an accurate distribution model of *R. reniformis* within a cotton field; however, the number of sampling points used (64 points within a 48 x 32 m area) would be cost prohibitive in a commercial setting. In a study assessing sampling grid size for variable rate application of nematicides for the management of *R. reniformis*, Ellis *et al.* (2004) found that fewer rate changes occurred with increasing grid size. This relationship has one of two possible consequences. The first is increased input of nematicides where they are not needed, which would result in a cost penalty. The second consequence would be not applying nematicides where needed, which would result in a yield penalty.

Technological pitfalls are also a possibility in the development of site-specific management. Choosing the correct analysis of spatial data is vital to producing accurate prescription maps. In a study of the accuracy of interpolating elevation data, a measurement commonly used in conjunction with EC for management zone delineation, Weng (2006) determined that accuracy was subject to a number of interpolation parameters that may significantly improve or worsen the accuracy. Similarly, it has been reported that apparent soil electrical conductivity is affected by soil transient properties such as volumetric soil water content and exhibits large changes throughout the season (McCutcheon *et al.*, 2006). Factors such as these can result in unreliable data and must be considered during management zone creation.

To create management zones within a field for *R. reniformis*, the factors of influence must first be

characterized through quantitative research and then the data can subsequently developed into a useable form. As was discussed earlier, soil texture distribution can be easily measured within a field by utilizing soil apparent electrical conductivity and has been used in management strategies for other species of nematodes. Consequently, this factor has been investigated as a starting point for zone delineation for *R. reniformis*. While *R. reniformis* is known to exist and cause damage in a wide variety of soil types (Gazaway & McLean, 2003), some research has suggested that *R. reniformis* is more prevalent in fine-textured soils (Robinson *et al.*, 1987; Starr *et al.*, 1993; Monfort *et al.*, 2008). Other research on the effects of soil type on *R. reniformis* populations has suggested that the productivity of the soil, not specifically soil texture, is the driving force behind population development (Koenning *et al.*, 1996; Herring *et al.*, 2010) as well as response to nematicides (Overstreet *et al.*, 2007, 2011, 2012).

Another consideration for zone delineation is initial populations of *R. reniformis* and economic damage threshold values. More often than not, management decisions and subsequently economic threshold values are based on post-harvest nematode sampling. Although little is known about the overwinter survivorship of *R. reniformis*, it has been observed that overwinter survivorship was lowest in areas of high sand content and increased with increasing clay content (Still & Kirkpatrick, 2006). Studies of overwinter survivorship on *Meloidogyne incognita* have suggested that population density and cultural practices have the greatest impact on overwinter survivorship (Ferris, 1985). Studies have shown that *R. reniformis* populations are adversely affected by post-harvest conventional tillage compared to non-tillage and ridge tillage (Cabanillas *et al.*, 1999). Economic thresholds are established based on the relationships between the degree of control and cost and nematode densities and crop value (Ferris, 1978). Current thresholds are established on a state-by state basis, but it has recently been suggested that different economic thresholds be considered based on soil type and productivity (Moore *et al.*, 2011b).

Studies exploring the possibilities of SSM and variable rate nematicide applications for *R. reniformis* have been conducted in recent years. Variable rate application based on populations of *R. reniformis* have been conducted with the fumigant nematicides 1,3-dichloropropene and metam sodium with promising results (Lawrence

et al., 2002; Ellis *et al.*, 2005). Farias *et al.* (2002) created a risk-benefit analysis for the treatment of *R. reniformis* in a Brazilian cotton field by utilizing geostatistical methods to interpolate population distribution over short distances (4-6 m). Another tool in development is the use of remotely sensed hyperspectral data to detect stress levels in cotton. Doshi *et al.* (2010) conducted a study comparing hyperspectral reflectance of cotton plants grown in microplots to *R. reniformis* populations in the plant rhizosphere and determined that this method could accurately estimate *R. reniformis* populations affecting the cotton plant. The use of remote sensing to detect cotton plant stress due to issues with subsurface drip irrigation has also illustrated this tool's ability to detect differences in cotton response to stress in field settings (Fulton *et al.*, 2008).

The successful use of site-specific management for *R. reniformis* on cotton is dependent on the resolution of several issues. The first and most important issue is to what spatial scale (single field, soil region, state, etc.) can general recommendations be developed and be reliable? Second, what parameters, or combination of parameters, will provide the most accurate measure of economic risk and subsequent usefulness in management zone creation? Third, can the two aforementioned issues be resolved in a manner which will result in a method that is easily adaptable for producers and will provide them with cost savings?

The issue of the size of the spatial scale upon which to separate recommendations includes two major considerations. *R. reniformis* is known to have geographical variation with respect to reproduction, pathogenicity, morphometrics, temperature effects on embryogenesis, and genetics (Agudelo *et al.*, 2001; Agudelo *et al.*, 2005; Arias *et al.*, 2009; Leach *et al.*, 2009; McGawley *et al.*, 2010), some of which vary within a single state. A second consideration is the diversity of soils within regions and states. For example, Alabama has six major soil areas where cotton is produced, each with quite different characteristics and levels of in-field variability. It is also well known that certain soils, such as those found in the Mississippi River Delta region, support far greater populations of *R. reniformis* in comparison to the soils found in the Coastal Plain region of the Southeast, yet the amount of yield loss in each region is similar.

The second issue is which parameters provide the best indicators of economic risk and subsequent usefulness in management zone creation? As was

detailed earlier, soil texture distribution has been studied quite extensively in relation to predicting which location in a field is more favorable to *R. reniformis* reproduction. While this technique has been used successfully for other species of plant-parasitic nematodes, the success of *R. reniformis* to reproduce and cause damage in a wide variety of soil textural distributions renders this method much less useful. Economic threshold level of *R. reniformis* is another parameter to be considered. Potential soil productivity has been shown to affect this relationship (Moore *et al.*, 2011a) as well as the possibilities of additional stress due to the lack of water throughout the growing season (Moore *et al.*, 2011c). The use of yield maps from previous years, if they exist, is another strong possibility for guidance of zone creation. Massey *et al.* (2008) determined that utilizing yield maps to assess profitability of corn, soybean, and grain sorghum based on field features and input costs could provide producers with information to assess management options.

Can SSM for *R. reniformis* on cotton become an easily adaptable and cost-saving tool for cotton producers? The answer depends on two major issues; spatial scale and zone creation parameters. Spatial scale and zone creation parameters are currently a focal point of research throughout areas affected by *R. reniformis*. Furthermore, many of the techniques for site-specific management are used for a variety of other issues and could be easily adapted with the correct guidelines. The identification of parameters to quantify economic risk and the understanding of how these parameters will differ over geographical areas will determine if SSM can enable cotton producers to gain an economic advantage over *R. reniformis*.

LITERATURE CITED

- Agudelo, P., R. T. Robbins, and J. M. Stewart. 2001. Morphometric variation of reniform nematode geographic populations from cotton-growing regions in the United States. Arkansas Agricultural Experiment Station, Research Series 497:87-91. <http://arkansasagnews.uark.edu/1292.htm>
- Agudelo, P., R. T. Robbins, and J. M. Stewart, A. Szalanski. 2005. Intraspecific variability of *Rotylenchulus reniformis* from cotton-growing regions in the United States. *Journal of Nematology* 37:105-114.
- Arias, R. S., S. R. Stetina, J. L. Tonos, J. A. Scheffler, and B. E. Scheffler. 2009. Microsatellites reveal genetic diversity in *Rotylenchulus reniformis* populations. *Journal of Nematology* 41:146-156.

- Baird, R. E., J. R. Rich, R. G. McDaniel, and B. G. Mullinix. 2000. Effects of nematicides on *Rotylenchulus reniformis* in cotton. *Nematologia Mediterranea* 28:83-88.
- Bayer CropScience. 2010. Bayer CropScience plans to discontinue aldicarb by 2014. Bayer CropScience 2010-0423E. [http://www.bayercropscience.com/bcsweb/cropprotection.nsf/id/EN_20100816/\\$file/2010-0423e.pdf](http://www.bayercropscience.com/bcsweb/cropprotection.nsf/id/EN_20100816/$file/2010-0423e.pdf)
- Blasingame, D., and M. V. Patel. 2002. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2002. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Blasingame, D., and M. V. Patel. 2003. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2003 Pp. 252-253. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Blasingame, D., and M. V. Patel. 2004. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2004 Pp. 459-450. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Blasingame, D., and M. V. Patel. 2005. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2005 Pp. 259-262. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Blasingame, D. 2006. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2006 Pp. 155-157. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Blasingame, D., J. C. Banks, P. D. Colyer, R. M. Davis, W. S. Gazaway, N. Goldberg, R. C. Kemerait, T. L. Kirkpatrick, S. R. Koenning, J. Muller, M. A. Newman, M. Olsen, P. M. Phipps, G. L. Scuibato, R. Sprenkel, J. E. Woodward, A. Wrather, M. V. Patel. 2008. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2008 Pp. 294-297. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Blasingame, D., W. Gazaway, K. Lawrence, A. Wrather, M. Olsen, N. Goldberg, T. Kirkpatrick, S. Koenning, M. Davis, J. C. Banks, R. K. Sprenkel, J. Muller, R. Kemerait, M. Newman, P. Colyer, J. Woodward, G. Scuibato, P. Phipps. 2009. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2009. Pp. 94-96. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Blasingame, D., M. V. Patel, K. Lawrence, W. Gazaway, M. Olsen, T. Kirkpatrick, S. Monfort, M. Davis, J. Marios, R. Kemerait, P. Colyer, G. Scuibato, G. Lawrence, A. Wrather, N. Goldberg, S. Koenning, J. T. Pitts, J. Muller, M. Newman, J. Woodward, T. Wheeler, P. Phipps. 2010. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2010. Pp. 237-240. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Blasingame, D., and M. V. Patel. 2011. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2011 Pp. 306-308. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Blasingame, D., and M. V. Patel. 2012. Cotton disease loss estimate committee report. Proceedings of the Beltwide Cotton Conferences, 2012. Pp. 341-344. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Cabanillas, H. E., J. M. Bradford, and J. R. Smart. 1999. Effect of tillage system, soil type, crop stand, and crop sequence on reniform nematodes after harvest. *Nematropica* 29:137-146.
- Carr, P. M., G. R. Carlson, J. S. Jacobsen, G. A. Nielsen, and E. O. Slogles. 1991. Farming soils, not fields: A strategy for increasing fertilizer profitability. *Journal of Production Agriculture* 4:57-61.
- Casa, R., and A. Castrignano. 2008. Analysis of spatial relationships between soil and crop variables in a durum wheat field using a multivariate geostatistical approach. *European Journal of Agronomy* 28:331-342.
- Castillo, J. D., K. S. Lawrence, and E. van Santen. 2009. Efficacy of *Arthrobotrys dactyloides*, *Dactylaria brochopaga*, *Fusarium oxysporium*, and *Paecilomyces lilacinus* for biocontrol of reniform nematode (*Rotylenchulus reniformis*). Proceedings of the Beltwide Cotton Conferences, 2009 Pp. 144-150. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Castillo, J. D., K. S. Lawrence, and J. W. Kloepper. 2011. Evaluation of *Bacillus firmus* (Votivo®) and *Paecilomyces lilacinus* strain 251 (Nemout®) for the biocontrol of reniform nematode *Rotylenchulus reniformis*. Proceedings of the Beltwide Cotton Conferences, 2011 Pp. 242-246. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Davis, R. F., S. R. Koenning, R. C. Kemerait, T. D. Cummings, and W. D. Shurley. 2003. *Rotylenchulus reniformis* management in cotton with crop rotation. *Journal of Nematology* 35:58-64.
- Davis, R. F., and T. M. Webster. 2005. Relative status of selected weeds and crops for *Meloidogyne incognita* and *Rotylenchulus reniformis*. *The Journal of Cotton Science* 9:41-46.
- Doerge, T. 1999. Defining management zones for precision farming. *Crop Insight*. Vol. 8 No. 21. Pioneer Hybrids, Johnston, IA.
- Doshi, R. A., R. L. King, and G. W. Lawrence. 2010. Classification of *Rotylenchulus reniformis*

- numbers in cotton using remotely sensed hyperspectral data on self-organizing maps. *Journal of Nematology* 42:179-193.
- Ellis, G. R., G. W. Lawrence, S. Samson, W. A. Givens, and K. S. Lawrence. 2004. Variable rate nematicides applications on cotton for reniform nematode management. *Journal of Nematology* 36:316.
- Ellis, G. R., G. W. Lawrence, S. A. Samson, and W. A. Givens. 2005. Variable rate applications of Telone II on cotton for reniform nematode management. 2005. Proceedings of the Beltwide Cotton Conferences, 2005 Pp. 195-196. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Environmental Protection Agency. 2002. Iterim re-registration eligibility decision (IRED) fenamiphos. EPA 738-R-02-004. http://www.epa.gov/REDs/fenamiphos_ired.pdf
- Evans, K., R. M. Webster, P. D. Halford, A. D. Barker, and M. D. Russell. 2002. Site-specific management of nematodes: Pitfalls and practicalities. *Journal of Nematology* 34:194-199.
- Farias, P. R. S., X. Sanchez-Vila, J. C. Barbosa, S. R. Vieira, L. C. C. B. Ferraz, and J. Solis-Delfin. 2002. Using geostatistical analysis to evaluate the presence of *Rotylenchulus reniformis* in cotton crops in Brazil: Economic implications. *Journal of Nematology* 34:232-238.
- Faske, T. R., and J. L. Starr. 2006. Sensitivity of *Meloidogyne incognita* and *Rotylenchulus reniformis* to abamectin. *Journal of Nematology* 38:240-244.
- Faske, T. R., and J. L. Starr. 2007. Cotton root protection from plant-parasitic nematodes by abamectin-treated seed. *Journal of Nematology* 39:27-30.
- Ferris, H. 1978. Nematode economic thresholds: Derivation, requirements, and theoretical considerations. *Journal of Nematology* 10:341-350.
- Ferris, H. 1985. Density-dependant nematode seasonal multiplication rates and overwinter survivorship: a critical point model. *Journal of Nematology* 17:93-100.
- Fiez, T. E., B. C. Miller, and W. L. Pan. 1994. Assessment of spatially variable nitrogen fertilizer management in winter wheat. *Journal of Production Agriculture* 7:86-93.
- Fleming, K. L., D. F. Heermann, and D. G. Westfall. 2004. Evaluating soil color with farmer input and apparent soil electrical conductivity for management zone delineation. *Agronomy Journal* 96:1581-1587.
- Fraisse, C. W., K. A. Sudduth, and N. R. Kitchen. 2001. Delineation of site-specific management zones by unsupervised classification of topographic attributes and soil electrical conductivity. *Transactions of the American Society of Agricultural Engineers* 44:155-166.
- Fulton, J. P., J. N. Shaw, D. Sullivan, M. P. Dougherty, and C. Brodbeck. 2008. Use of remote sensed thermal imagery for in-season stress detection and site-specific management of cotton. Alabama Agricultural Experiment Station Research Report Series No. 32:26-27. <http://www.ag.auburn.edu/aaes/communications/researchreports/07cottonrr.pdf>
- Gazaway, W. S., and K. S. McLean, 2003. Plant pathology and nematology: A survey of plant-parasitic nematodes associated with cotton in Alabama. *J. Cot. Sci.* 7:1-7.
- Goovaerts, P. 1992. Factorial kriging analysis: a useful tool for exploring the structure of multivariate geostatistics. *Geoderma* 62:93-107.
- Goovaerts, P. 1998. Geostatistical tools for characterizing the spatial variability of microbiological and physico-chemical soil properties. *Biology and Fertility of Soils* 27:315-334.
- Heermann, D. F., J. Hoeting, H. R. Duke, D. G. Westfall, G. W. Buckleiter, P. Westra, F. B. Peairs, and K. L. Fleming. 1999. Interdisciplinary irrigated precision farming team research. Pp. 212-230 in J. V. Stafford (ed.) *Precision Agriculture '99*. Vol. 1. Soil and crop factors: Location, sensing, and sampling. Oxford, UK.
- Herring, S. L., S. R. Koenning, and J. L. Heitman. 2010. Impact of *Rotylenchulus reniformis* on cotton yield as affected by soil texture and irrigation. *Journal of Nematology* 42:319-323.
- Jaynes, D. B., J. M. Novak, T. B. Moorman, and C. M. Cambardella. 1994. Estimating herbicide partition coefficients from electromagnetic induction measurements. *Journal of Environmental Quality* 24:26-41.
- Jaynes, D. B. 1996. Improved soil mapping using electromagnetic induction surveys. Pp. 169-179. Proceedings of the 3rd International Conference on Site-Specific Management for Agricultural Systems. ASA, CSSA, and SSSA, Madison, WI.
- Jones, J. R., K. S. Lawrence, and G. W. Lawrence. 2006. Evaluation of winter cover crops in cotton cropping for management of *Rotylenchulus reniformis*. *Nematropica* 36:53-66.
- Kachanoski, R. G., E. G. Gregorich, and I. J. Van Wesenbeck. 1988. Estimating spatial variations of soil water content using noncontacting electromagnetic inductive methods. *Canadian Journal of Soil Science* 68:715-722.
- Khalilian, A., J. D. Mueller, Y. J. Han, and F. J. Wolak. 2001. Predicting cotton nematodes distribution utilizing soil electrical conductivity. Proceedings of the Beltwide Cotton Conferences, 2001 Pp. 146-148. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Khalilian, A., J. Mueller, S. Lewis, and Y. Han. 2002. Relationship of Columbia lance and root-

- knot nematodes to soil type. Proceedings of the Beltwide Cotton Conferences, 2002. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Khalilian, A., J. D. Mueller, and Y. J. Han. 2003. Performance of variable rate nematicides application systems. Proceedings of the Beltwide Cotton Conferences, 2003 Pp. 578-582. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Kinloch, R. A., and J. R. Rich. 2001. Management of root-knot and reniform nematodes in ultra-narrow row cotton with 1,3-dichloropropene. *Journal of Nematology* 33:311-313.
- Kitchen, N. R., K. A. Sudduth, and S. T. Drummond. 1999. Soil electrical conductivity as a crop productivity measure for clay pan soils. *Journal of Production Agriculture* 12:607-617.
- Koenning, S. R., S. A. Walters, and K. R. Barker. 1996. Impact of soil texture on the reproductive and damage potentials of *Rotylenchulus reniformis* and *Meloidogyne incognita* on cotton. *Journal of Nematology* 28:527-536.
- Koenning, S. R., D. E. Morrison, and K. L. Edmisten. 2007. Relative efficacy of selected nematicides for management of *Rotylenchulus reniformis* in cotton. *Nematropica* 37:227-235.
- Kyaw, T., R. B. Ferguson, V. I. Adamchuk, D. B. Marx, D. D. Tarkalson, and D. L. McCallister. 2008. Delineating site-specific management zones for pH-induced iron chlorosis. *Precision Agriculture* 9:71-84.
- Lawrence, G. W., K. S. McLean, W. E. Batson, D. Miller, and J. C. Borbon. 1990. Response of *Rotylenchulus reniformis* to nematicides applications on cotton. *Journal of Nematology* 22:707-711.
- Lawrence, G. W., and K. S. McLean. 2000. Effect of foliar applications of oxamyl with aldicarb for the management of *Rotylenchulus reniformis* on cotton. *Journal of Nematology* 32:542-549.
- Lawrence, G. W., K. S. McLean, W. A. Givens, R. K. Mehrle, H. K. Lee, and A. T. Kelley. 2002. Reniform nematode management on cotton with VRT and site specific applications. Proceedings of the Beltwide Cotton Conferences, 2002. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Lawrence, K. S., Y. Feng, G. W. Lawrence, C. H. Burmester, and S. H. Norwood. 2005. Accelerated degradation of aldicarb and its metabolites in cotton field soils. *Journal of Nematology* 37:190-197.
- Lawrence, K. S., and G. W. Lawrence. 2007. Performance of the new nematicides seed treatments on cotton. Proceedings of the Beltwide Cotton Conferences, 2007 Pp. 602-605. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Lawrence, K. S., A. J. Price, G. W. Lawrence, J. R. Jones, and J. R. Akridge. 2008. Weed hosts for *Rotylenchulus reniformis* in cotton fields rotated with corn in the southeast United States. *Nematropica* 38:13-22.
- Leach, M., P. Agudelo, and P. Gerard. 2009. Effect of temperature on the embryogenesis of geographic populations of *Rotylenchulus reniformis*. *Journal of Nematology* 41:23-27.
- Lee, H. K., G. W. Lawrence, A. T. Kelley, and K. S. McLean. 2002. Horizontal and vertical distribution of *Rotylenchulus reniformis* in a Mississippi cotton field. Proceedings of the Beltwide Cotton Conferences, 2002. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Liu, T., K. Juang, and D. Lee. 2006. Interpolating soil properties using kriging combined with categorical information of soil maps. *Soil Science Society of America Journal* 70:1200-1209.
- Lopez-Lozano, R., M. A. Casterad, and J. Herrero. 2010. Site-specific management units in a commercial maize plot delineated using very high resolution remote sensing and soil properties mapping. *Computers and Electronics in Agriculture* 73:219-229.
- Massey, R. E., D. B. Myers, N. R. Kitchen, and K. A. Sudduth. 2008. Profitability maps as an input for site-specific management decision making. *Agronomy Journal* 100:52-59.
- McBride, R. A., A. M. Gordon, and S. C. Shrive. 1990. Estimating forest soil quality from terrain measurements of apparent electrical conductivity. *Soil Science Society of America Journal* 54:290-293.
- McCutcheon, M. C., H. J. Farahani, J. D. Stednick, G. W. Buchleiter, and T. R. Green. 2006. Effect of soil water on apparent soil electrical conductivity and texture relationships in a dryland field. *Biosystems Engineering* 94:19-32.
- McGawley, E. C., M. J. Pontif, and C. Overstreet. 2010. Variation in reproduction and pathogenicity of geographic isolates of *Rotylenchulus reniformis* on cotton. *Nematropica* 40:275-288.
- Monfort, W. S., T. L. Kirkpatrick, C. S. Rothrock, and A. Mauromoustakos. 2007. Potential for site-specific management of *Meloidogyne incognita* in cotton using soil textural zones. *Journal of Nematology* 39:1-8.
- Monfort, W. S., T. L. Kirkpatrick, and A. Mauromoustakos. 2008. Spread of *Rotylenchulus reniformis* in an Arkansas cotton field over a four-year period. *Journal of Nematology* 40:162-166.
- Moore, S. R., K. S. Lawrence, F. J. Arriaga, C. H. Burmester, and E. van Santen. 2010a. Natural migration of *R. reniformis* in a no-till cotton system. *Journal of Nematology* 42:307-312.
- Moore, S. R., K. S. Lawrence, B. V. Ortiz, J. N. Shaw, and J. Fulton. 2010b. Evaluation of nematicides

- for the management of *Rotylenchulus reniformis* across management zones created using soil electrical conductivity. *Phytopathology* 100:S86.
- Moore, S. R., W. S. Gazaway, K. S. Lawrence, B. Goodman, and J. R. Akridge. 2010c. Value of rotational crops for profit increase and reniform nematode suppression with and without a nematicides in Alabama. Proceedings of the Beltwide Cotton Conferences, 2010 Pp. 260-268. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Moore, S. R., K. S. Lawrence, F. J. Arriaga, C. H. Burmester, and E. van Santen. 2011a. Influence of water movement and root growth in the downward dispersion of *Rotylenchulus reniformis*. *Nematopica* 41:75-81.
- Moore, S. R., and K. S. Lawrence. 2011b. Effects of soil type on the reproductive potential of *Rotylenchulus reniformis* on cotton. Proceedings of the Beltwide Cotton Conferences, 2011 Pp. 235-240. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Moore, S. R., K. S. Lawrence, B. Ortiz, J. Shaw, and J. Fulton. 2011c. Evaluation of the effects of soil moisture on the damage potential of *Rotylenchulus reniformis* on cotton. *Phytopathology* 101:S123
- Newman, M. A., and T. C. Stebbins. 2002. Recovery of reniform nematodes at various soil depths in cotton. Proceedings of the Beltwide Cotton Conferences, 2002. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Ortiz, B. V., D. Sullivan, C. Perry, and G. Vellidis. 2007. Geostatistical analysis of the spatial variability of cotton-parasitic nematodes and the factors favoring its occurrence. Proceedings of the Beltwide Cotton Conferences, 2007 Pp. 920-928. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Ortiz, B. V., C. Perry, D. Sullivan, R. Kemerait, A. R. Ziehl, R. Davis, G. Vellidis, and K. Rucker. 2008. Cotton yield response to variable rate nematicides according to risk zones. Proceedings of the Beltwide Cotton Conferences, 2008 Pp. 573-582. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Ostergaard, H. G. S. 1997. Agronomic consequences of variable N fertilization. Pp. 315-320 in J. V. Stafford (ed.) *Precision Agriculture '97*. Vol. 1. Spatial variability in soil and crop. Oxford, UK.
- Overstreet, C., E. Burris, D. R. Cook, E. C. McGawley, B. Padgett, and M. Wolcott. 2007. Telone II fumigation. Proceedings of the Beltwide Cotton Conferences, 2007 Pp. 587-597. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Overstreet, C., R. Barbosa, D. Burns, R. L. Frazier, E. C. McGawley, G. B. Padgett, and M.C. Wolcott. 2011. Using electrical conductivity to determine nematode management zones in alluvial soils of the mid-South. Proceedings of the Beltwide Cotton Conferences, 2011 Pp. 252-258. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Overstreet, C., E. C. McGawley, D. Burns, R. L. Frazier, and R. Barbosa. 2012. The influence of apparent electrical conductivity of the soil on nematicides in cotton. Proceedings of the Beltwide Cotton Conferences, 2012 Pp. 288-292. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Rab, M. A., P. D. Fisher, R. D. Armstrong, M. Abuzar, N. J. Robinson, and S. Chandra. 2009. Advances in precision agriculture in south-eastern Australia. IV. Spatial variability in plant-available water capacity of soil and its relationship with yield in site-specific management zones. *Crop & Pasture Science* 60:885-900.
- Rich, J. R., and R. A. Kinloch. 2000. Influence of aldicarb and 1,3-dichloropropene applications on cotton yield and *Rotylenchulus reniformis* post-harvest populations. *Nematopica* 30:47-53.
- Robinson, A. F., C. M. Heald, S. L. Flanagan, W. H. Thames, and J. Amador. 1987. Geographical distributions of *Rotylenchulus reniformis*, *Meloidogyne incognita*, and *Tylenchulus semipenetrans* in the Lower Rio Grande Valley as related to soil texture and land use. *Annals of Applied Nematology* (Journal of Nematology 19, Supplement) 1: 20-25.
- Robinson, A. F., R. N. Inserra, E. P. Caswell-Chen, N. Vovlas, and A. Troccoli. 1997. *Rotylenchulus* species: identification, distribution, host ranges, and crop plant resistance. *Nematopica* 27:127-180.
- Robinson, A. F., C. G. Cook, A. Westphal, and J. M. Bradford. 2005a. *Rotylenchulus reniformis* below plow depth suppresses cotton yield and root growth. *Journal of Nematology* 37:285-291.
- Robinson, A. F., R. Akridge, J. M. Bradford, C. G. Cook, W. S. Gazaway, T. L. Kirkpatrick, G. W. Lawrence, G. Lee, E. C. McGawley, C. Overstreet, B. Padgett, R. Rodriguez-Kabana, A. Westphal, and L. D. Young. 2005b. Vertical distribution of *Rotylenchulus reniformis* in cotton fields. *Journal of Nematology* 37:265-271.
- Robinson, A. F. 2007. Reniform in U.S. cotton: when, where, why, and some remedies. *Annual Review of Phytopathology* 45:263-288.
- Starr, J. L., C. M. Heald, A. F. Robinson, R. G. Smith, and J. P. Krausz. 1993. *Meloidogyne incognita* and *Rotylenchulus reniformis* and associated soil textures from some cotton production areas of Texas. *Journal of Nematology* 25:895-899.
- Stetina, S. R., L. D. Young, W. T. Pettigrew, and H. A. Bruns. 2007. Effect of corn-cotton rotations on reniform nematode populations and crop yield. *Nematopica* 37:237-248.
- Still, J. A., and T. L. Kirkpatrick. 2006. Ecology and

- over-wintering ability of *Rotylenchulus reniformis* on cotton in Arkansas. Research Series 552. Summaries of Arkansas Cotton Research 2006. <http://arkansasagnews.uark.edu/1993.htm>
- Sudduth, K. A., D. F. Hughes, and S. T. Drummond. 1995. Electromagnetic induction sensing as an indicator of productivity on claypan soils. Pp. 671-681. Proceedings of the 2nd International Conference on Site-Specific Management for Agricultural Systems. ASA, Madison, WI.
- United States Department of Agriculture, National Agricultural Statistical Service. 2011. www.nass.usda.gov
- Usery, S. R., K. S. Lawrence, G. W. Lawrence, and C. H. Burmester. 2005. Evaluation of cotton cultivars for resistance and tolerance to *Rotylenchulus reniformis*. *Nematropica* 35:121-133.
- Wackernagel, H. 1988. Geostatistical techniques for interpreting multivariate spatial information. Pp. 393-409 in Chung, C. F., A. Fabbri, G. R. Sinding-Larsen. (eds) Quantitative analysis of mineral and energy resources. Reidel, Dordrecht.
- Wackernagel, H. 1995. Multivariate geostatistics: and introduction with applications. Springer, Berlin Heidelberg New York.
- Wang, K. H., B. S. Sipes, and D. P. Schmitt. 2003. Intercropping cover crops with pineapple for the management of *Rotylenchulus reniformis*. *Journal of Nematology* 35:39-47.
- Wang, K., R. D. Riggs, and D. Crippen. 2004. Suppression of *Rotylenchulus reniformis* on cotton by the nematophagous fungus ARF. *Journal of Nematology* 36(2):186-191.
- Weng Q. 2006. An evaluation of spatial interpolation accuracy of elevation data. Pp. 805-824 in *Progress in spatial data handling*. Riedl, A., W. Kainz, G. A. Elmes eds. Springer Berlin Heidelberg.
- Westphal, A., and J. R. Smart. 2003. Depth distribution of *Rotylenchulus reniformis* under different tillage and crop sequence systems. *Phytopathology* 93:1182-1189.
- Westphal, A., A. F. Robinson, A. W. Scott, Jr., and J. B. Santini. 2004. Depth distribution of *Rotylenchulus reniformis* under crops of different host status and after fumigation. *Nematology* 6:97-107.
- Williams, B. G., and D. Hoey. 1987. The use of electromagnetic induction to detect spatial variability of the salt and clay content of soils. *Australian Journal of Soil Research* 25:21-27.
- Wolcott, M. C., C. Overstreet, B. Padgett, and E. Burris. 2004. Using soil electrical conductivity to denote potential nematode management zones. Proceedings of the Beltwide Cotton Conferences, 2004 Pp. 349-353. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Wolcott, M. C., C. Overstreet, E. Burris, D. Cook, D. Sullivan, G. B. Padgett, and R. Goodson. 2005. Evaluating cotton nematicides response across soil electrical conductivity zones using remote sensing. Proceedings of the Beltwide Cotton Conferences, 2005 Pp. 215-220. National Cotton Council of America. Online. <http://www.cotton.org/beltwide/proceedings.htm>
- Wyse-Pester, D. Y., L. J. Wiles, and P. Westra. 2002. The potential for mapping nematode distributions for site-specific management. *Journal of Nematology* 34:80-87.

Received:

13/I/2012

Accepted for publication:

13/VI/2012

Recibido:

Aceptado para publicación: