

ELECTRONIC ARTICLE/ARTICULO ELECTRONICO

(Available at Nematropica Online: <http://palmm.fcla.edu/nematode/>)**PHYLOGENETIC RELATIONSHIPS AMONG MEXICAN POPULATIONS OF *NACOBBUS ABERRANS* (NEMATODA, PRATYLENCHIDAE) REVEAL THE EXISTENCE OF CRYPTIC (COMPLEX) SPECIES**

A. J. Cabrera-Hidalgo, N. Marban-Mendoza, and E. Valadez-Moctezuma 1-11

FURTHER ELUCIDATION OF THE HOST RANGE OF *GLOBODERA ELLINGTONAE*

A. B. Peetz, H.V. Baker, and I. A. Zasada..... 12-17

DISTRIBUTION AND DIVERSITY OF CYST NEMATODE (NEMATODA: HETERODERIDAE) POPULATIONS IN THE REPUBLIC OF AZERBAIJAN, AND THEIR MOLECULAR CHARACTERIZATION USING ITS-RDNA ANALYSIS

A. A. Dababat, H. Muminjanov, G. Erginbas-Orakci, G. Ahmadova Fakhreddin, L. Waeyenberge, S. Yildiz, N. Duman, and M. Imren 18-30

PATHOGENICITY AND REPRODUCTION OF ISOLATES OF RENIFORM NEMATODE, *ROTYLENCHULUS RENIFORMIS*, FROM LOUISIANA ON SOYBEAN

M. T. Kularathna, C. Overstreet, E. C. McGawley, S. R. Stetina, C. Khanal, F. M. C. Godoy, and B. K. McInnes..... 31-41

INFLUENCE OF TWO *NACOBBUS ABERRANS* ISOLATES FROM ARGENTINA ON THE GROWTH OF THREE TOMATO CULTIVARS

V. A. Cabrera, N. Dottori, and M. E. Doucet 42-48

CHARACTERIZATION OF A POPULATION OF *HELICOTYLENCHUS DIHYSTERA* (COBB, 1893) SHER, 1961, PARASITIZING MAIZE ROOTS, IN SOUTHERN CÓRDOBA, ARGENTINA

E. Brücher, E. E. Vuletic, F. A. Guerra, R. L. De Rossi, M. C. Plazas, G. D. Guerra, M. G. Molina, M. M. Gabriela, and M. E. Doucet 49-58

EFFECT OF SEED GALL NEMATODE, *ANGUINA TRITICI*, ON GRAIN YIELD OF BREAD WHEAT CULTIVAR UNDER FIELD CONDITIONS

A. Tülek, A. A. Dababat, İ. Kepenekci, and İ. Öztürk..... 59-62

**PLANTING DIFFERENT CROPS IN SUCCESSION TO MANAGE
PRATYLENCHUS ZEAE IN SUGARCANE**

- S. M. Santana-Gomes, C. R. Dias-Arieira, F. Biela, M. Ragazzi, S. S. Baldisera, and
 R. P. Schwengber..... 63-70

**DISTRIBUTION OF *MELOIDOGYNE* SPECIES (TYLENCHIDA:
 MELOIDOGYNIDAE) IN TOMATO CROP IN SINALOA, MEXICO**

- J. A. Carrillo-Fasio, J. A. Martínez-Gallardo, R. Allende-Molar, S. Velarde-Félix,
 C. E. Romero-Higareda, and J. E. Retes-Manjarrez 71-82

**EVALUACIÓN DE DOS CEPAS DE NEMATODOS
 ENTOMOPATÓGENOS (STEINERNEMATIDAE,
 HETERORHABDITIDAE) PARA EL CONTROL DEL SALIVAZO
 (HEMIPTERA: CERCOPIDAE) EN CAÑA DE AZÚCAR**

- P. F. Grifaldo-Alcántara, R. Alatorre-Rosas, J. A. Villanueva-Jiménez, F. Hernández-Rosas,
 S. P. Stock, y G. Ramírez-Valverde 83-90

**SCREEN HOUSE RESPONSE OF SEVEN ELITE CASSAVA (*MANIHOT
 ESCULENTA* CRANTZ) VARIETIES TO *MELOIDOGYNE
 INCognITA* INFECTION**

- A. K. Akinsanya and S. O. Afolami 91-98

***POCHONIA CHLAMYDOSPORIA* NO CONTROLE DO NEMATOIDE DE
 GALHAS EM BANANEIRA**

- R. T. Barbosa, T. S. A. Monteiro, R. R. Coutinho, J. G. Silva, and L. G. Freitas 99-106

**INCIDENCE OF *APHELENCHOIDES BESSEYI* IN RICE IN LOUISIANA
 AND HOST STATUS OF THE MOST WIDELY PLANTED
 CULTIVARS**

- F. M. C. Godoy, C. Overstreet, E. C. McGawley, C. A. Hollier, M. T. Kularathna, C. Khanal,
 and B. McInnes 107-123

**DAMAGE POTENTIAL OF *MELOIDOGYNE INCognITA*
 POPULATIONS ON SELECTED TOMATO GENOTYPES IN
 ETHIOPIA**

- A. Seid, C. Fininsa, T. M. Mekete, W. Decraemer, and W. M. L. Wesemael 124-133