

RESEARCH/INVESTIGACIÓN

PLANT-PARASITIC NEMATODES ASSOCIATED WITH PLANTAIN (*MUSA PARADISIACA*) IN TALAMANCA, LIMÓN, COSTA RICA

W. Peraza-Padilla^{1*}, R. Artavia-Carmona¹, E. Arboleda-Julio²,
R. Rodríguez-Porras³, and S. Orozco-Cayasso⁴

¹Laboratory of Nematology; ²Agribusiness; ⁴Laboratory of Entomology; and ⁵Laboratory of Phytopathology, National University of Costa Rica, 86-3000, Heredia, Costa Rica. *Corresponding author: walter.peraza.padilla@una.cr

ABSTRACT

Peraza-Padilla, W., R. Artavia-Carmona, E. Arboleda-Julio, R. Rodríguez-Porras, and S. Orozco-Cayasso. 2020. Plant-parasitic nematodes associated with plantain (*Musa paradisiaca*) in Talamanca, Limón, Costa Rica. *Nematopica* 50:151-159.

Plantain is one of the most important food and commercial crops in the Huetar Atlantica region of Costa Rica. Plant-parasitic nematodes (PPN) are one of the major constraints of plantain production and are therefore a threat to food security. The present study was part of an investigation that was carried out in Talamanca, Limón, Costa Rica with the objective to quantify and identify the main PPN genera associated with plantain cultivation. Nematodes were extracted from a total of 76 samples; 63 from soil and 13 from roots of plantain collected from 10 smallholder farms. In terms of absolute frequencies of distribution, the most abundant PPN genus in soil was *Pratylenchus*, which was detected in over 76.9% samples, followed by *Helicotylenchus* (69.2%), *Meloidogyne* and *Radopholus* (61.5%). In the roots, *Helicotylenchus* (100%), *Meloidogyne* (90.5%), *Pratylenchus* (74.6%), and *Radopholus* (66.7%) were the most frequently detected genera. In general, frequency and population densities of PPN were high in comparison to other similar surveys. Overall, the data suggest that PPNs could pose a serious threat to plantain production in Costa Rica and that control measures should be further developed and implemented.

Key words: *Helicotylenchus*, *Meloidogyne*, plant-parasitic nematodes, *Pratylenchus*, *Radopholus*

RESUMEN

Peraza-Padilla, W., R. Artavia-Carmona, E. Arboleda-Julio, R. Rodríguez-Porras, and E. Orozco-Cayasso. 2020. Nematodos vegetales asociados con plátano (*Musa paradisiaca*) en Talamanca, Limón, Costa Rica. *Nematopica* 50:151-159.

El plátano es uno de los cultivos alimentarios y comerciales más importantes de la región Huetar Atlántica de Costa Rica y del mundo. Los nematodos parásitos de plantas (NPP) son una de las principales limitaciones de la producción de plátano y, por lo tanto, son una amenaza para la seguridad alimentaria. El presente estudio fue parte de una investigación que se llevó a cabo en Talamanca, Limón y cuyo objetivo fue cuantificar e identificar los principales NPP asociados al cultivo de plátano. Los nematodos se extrajeron de un total de 76 muestras, 63 de suelo y 13 de raíz de plantas de plátano recolectadas en diez fincas de pequeños agricultores. En términos de frecuencias absolutas de distribución, el género más abundante en el suelo fue *Pratylenchus*, que se detectó en más del 76,9% de las fincas, seguido de

Helicotylenchus (69,2%), *Meloidogyne* y *Radopholus similis* (61,5%). En el caso de la raíz, el género más frecuentemente encontrado fue *Helicotylenchus* (100%), *Meloidogyne* (90.5%), *Pratylenchus* (74.6%), finalmente, *R. similis* (66.7%). En general, la frecuencia y la densidad de población de PPN fue alta. Esto probablemente se debió al manejo de las condiciones agronómicas y climáticas observadas durante los últimos años. En general, los datos sugieren que las PPN podrían representar una grave amenaza para las plantaciones de plátano y que deberían desarrollarse e implementarse medidas de control.

Palabras clave: *Helicotylenchus*, *Meloidogyne*, nematodos parásitos de plantas, *Pratylenchus*, *Radopholus*

INTRODUCTION

Plantain, *Musa paradisiaca*, is in the Musaceae family. The fruit is less sweet than that of banana, the other important species in the genus *Musa*. It is estimated that *Musa* is cultivated in more than 125 tropical and subtropical countries (Jones, 2000). Plantains are a major source of carbohydrates for millions of people in Africa, the Caribbean, Latin America, Asia, and the Pacific (FAO, 2003).

Plantain and banana have been ranked as the fourth most important food crop in the world after rice (*Oryza sativa*), maize (*Zea mays*), and wheat (*Triticum* spp.) (Frison and Sharrock, 1999). Plantain and banana are used for food, beverages, fermentable sugars, medicines, flavorings, and cooking (Nelson *et al.*, 2006; Phillip *et al.*, 2009). According to Oladejo and Sanusi (2008), plantain contributes to nutrition and food security in rural and urban areas. The main producing countries of plantain are Democratic Republic of Congo, Ghana, Cameroon, Uganda, Colombia, Philippines, Nigeria, and Perú (FAO, 2018). In sub-Saharan Africa, for example, it is estimated that over 30 million people subsist on banana and plantain as the principal source of dietary carbohydrate (Karamura *et al.*, 1999). *Musa* plantations are a major economic base in Central American, Caribbean, and some South American countries. According to (FAO, 2018), the production area and yield of plantain globally was estimated at 5,643,475 ha and 39,482,164 tons, respectively.

In Costa Rica, *Musa* spp. is one the most important crops (Sánchez and Zúñiga, 2001), representing almost 18.8% of the agricultural gross national products just below pineapple, which represents 19.1%. Exported *Musa* products in 2017 were valued at \$1,042,171 (USD) produced on 42,921 ha (Mora and Quirós, 2018).

Historically, plantain cultivation has had an

important role in the social development and economy of Costa Rica. Plantain production in Costa Rica is concentrated in the Talamanca region and often is creatively integrated into various cropping systems on small-scale farms. The region has the potential for development of industrial processing of plantain into snack products and plantain flour. This commercial potential has increased the interest of small producers to grow plantain; however, there are problems in commercialization and prices. In addition, there are several limiting production factors, including plant-parasitic nematodes (PPN) and Black Sigatoka disease (*Mycosphaerella fijiensis*), which reduces bunch weight and increases the time needed for fruit to develop to maturity (Araya, 2002).

Plant-parasitic nematodes are the primary pest of *Musa* crops worldwide (Gowen and Quénehervé, 1990; Kshaija *et al.*, 1994; Pattison, 2011). In Costa Rican plantations, mixtures of the following PPN usually occur: *R. similis* (Cobb, 1893, Thorne, 1949, Sher, 1968), *Helicotylenchus multicinctus* (Cobb, 1893, Golden, 1956), *H. dihystra* (Cobb, 1893; Sher, 1961), *Meloidogyne incognita* (Kofoid and White, 1919, Chitwood, 1949), *M. javanica* (Chitwood, 1949, Treub, 1985), and *Pratylenchus* spp. (Linford and Oliveira, 1940). Plantain production in Costa Rica is characterized as being highly dependent on agrochemicals, which increase costs and productive vulnerability. The regular application of nematicides is the most widely used strategy to control PPN in Costa Rica; however, the environmental implications and restrictions in international markets dictate a rational use of nematicides at minimum dosages.

Considering the economic importance of plantain as well as the destructive nature of PPN to plantain, the objective of this study was to provide quantitative information about population densities and frequencies of the major PPN associated with

plantain in smallholder farmer fields in Talamanca, Limón. This information will be used to identify the most important PPN associated with plantain in Costa Rica.

MATERIALS AND METHODS

Study area and description

The survey on PPN associated with plantain was conducted from January 2016 to December 2017 among smallholder plantain farmers of Bratsi, Telire, and Sixaola Districts, Talamanca Canton, Costa Rica. The farms were located at an altitude of 10 to 70 m above sea level (Fig. 1). Ten farms from the production area were surveyed. According to Holdridge (1982), this region is located in a humid tropical forest with annual rainfall ranging from 1 m to 2 m. In this region, mostly native varieties are grown in rural and indigenous communities in small fields. In some locations near the Sixaola District, farmers produce plantain cv. Curraré enano.

Soil and root samples

A total of 76 samples were collected. Each sample consisted of 20 soil cores (30 cm width × 20 cm deep). Holes were dug in the rhizosphere of each of the sampled plants. Samples were spaced at roughly equal intervals in a zig-zag pattern. Soil and root samples were placed in sealed plastic bags and protected from both direct sunlight and high temperatures. The samples were stored at 4°C before being analyzed to minimize changes in nematode population densities. A handheld global positional system device (GPS) (Garmin, GPS Map 60CSx, Garmin, Olathe, KS) was used to identify the locations.

Nematode extraction

Nematodes were extracted from 100 g of soil and 10 g of roots. Soil samples were processed by the flotation-centrifugation method (Jenkins, 1964). Root samples were washed in running tap water and chopped into small pieces (~ 1 cm),




Figure 1. Map of Costa Rica showing the site studied, Talamanca, Limón province, 2016-2017.

macerated in a kitchen blender for 20 sec at low speed, and the resultant suspension passed through two sieves (400 μm and 100 μm) to collect nematodes. Nematodes were killed and fixed in hot 4% formaldehyde (70°C) and subsequently infiltrated with glycerin, using Seinhorst's modified slow method (Seinhorst, 1959, 1962) and mounted on slides for observation and preservation. All specimens of this study were deposited in the Laboratory of Nematology Collection at Universidad Nacional, Heredia, Costa Rica.

Nematode identification and enumeration

Nematodes were identified and counted to determine the population densities of all PPN in both the soil and root samples. PPN were identified to genus and/or species level (if possible) using an Olympus BX50 microscope (Tokyo, Japan). The identification of PPN was carried out with the help of taxonomic keys developed by Thorne (1961) and Mai *et al.* (1996).

Data analysis

Frequency distribution analysis for each nematode genus and species by farmer field was determined, and the absolute frequency of each nematode genus was calculated (Barker, 1985). *Radopholus similis* and *Pratylenchus* sp. males and females, *Helicotylenchus* sp. females, males, and juveniles, and *Meloidogyne* sp. juveniles were included in the data analysis.

RESULTS

Six genera and one species of PPN were found to be associated with the surveyed plantations, including: *Meloidogyne* sp., *Helicotylenchus* sp., *Pratylenchus* sp., *Tylenchus* sp., *Aphelenchus* sp., *Aphelenchoides* sp. and *R. similis*.

Stereoscopic examination of the root system revealed the presence of necrotic lesions. These lesions in some roots were yellow, typical of the beginning of PPN injury. However, other reddish-brown to black lesions were also observed, showing advanced and necrotic states of the lesion produced by the migratory endoparasitic nematodes *Pratylenchus* spp. and *R. similis*. The magnitude of necrosis observed in the current study may have been the result of either high population

densities of *R. similis* (9,520 nematodes/10 g root) or *Pratylenchus* sp. (1,762 nematodes/10 g root).

Frequency of PPN in plantain soil

From the 10 farmer fields surveyed in Talamanca canton, the major PPN associated with plantain in the soil were in the following order of frequencies: *Helicotylenchus* spp., *Meloidogyne* spp., *Pratylenchus* spp. and *R. similis* (Table 1). Population densities of these nematodes varied among farms. Out of the 10 farms, the most abundant nematode based on total number of nematodes per each genus was *Helicotylenchus* spp. with an average population density of 78 nematodes/100 g soil, followed by *Meloidogyne* spp., and *Pratylenchus* spp. which had average population densities of 33 and 25 nematodes/100 g soil, respectively. In the case of *R. similis* in Costa Rica, this nematode is considered to be one of the most important PPN associated with plantain plantations. However, it was the least abundant with a population density of 12 nematodes/100 g soil. The average population density of all nematode species for the 4 genera from the 10 farms was 147 nematodes/100 g soil (Table 1). *Pratylenchus* sp. and *Helicotylenchus* sp. were widespread and occurred in soil in 77% and 69% of the farms surveyed, respectively. *Radopholus similis* and *Meloidogyne* sp. occurred in soil in 61.5% of the farms surveyed (Table 1).

Frequency of PPN in plantain roots

From the 10 farmer fields surveyed, the major PPN associated with plantain roots were: *R. similis*, *Pratylenchus* spp., *Helicotylenchus* spp., and *Meloidogyne* spp. (Table 1). The most abundant nematode was *R. similis* with a highest average population density of 9,520 nematodes/100 g root, followed by *Pratylenchus* spp. and *Helicotylenchus* spp., which had average population densities of 176 and 561 nematodes/100 g root, respectively. *Meloidogyne* spp. was the least abundant with a population density of 222 nematodes/100 g root; however, this nematode was present in 10 of 13 samples (Table 1). The average population density of all nematode species for the 4 genera from the 10 farms was 1,207 nematodes/100 g root (Table 1). *Helicotylenchus* spp. and *Meloidogyne* spp. were widespread and occurred in roots in 100% and 91% of the farms

Table 1. Analysis of plant-parasitic nematode (PPN) from soil and root samples of plantain grown in Talamanca, Limón, Costa Rica in 2016-2017.

Nematode	No. samples collected		No. positive samples		Avg no. of nematodes		Frequency of distribution (%)	
	Soil	Root	Soil	Root	100 g soil	100 g root	Soil	Root
<i>Radopholus similis</i>	63	13	42	8	12	9,520	62	67
<i>Pratylenchus</i> spp.	63	13	47	8	25	1,762	77	75
<i>Helicotylenchus</i> spp.	63	13	63	9	78	561	69	100
<i>Meloidogyne</i> spp.	63	13	57	10	33	222	62	91

surveyed, respectively, while *Pratylenchus* spp. and *R. similis* occurred in root in 75% and 67% of the farms surveyed, respectively (Table 1).

DISCUSSION

The PPN detected in the rhizosphere of plantain in the Talamanca region, *Meloidogyne* sp., *Helicotylenchus* sp., *Pratylenchus*, and *R. similis*, have been reported in Costa Rica (López, 1980; Araya *et al.*, 2002; Araya, 2002, 2003, 2005; Chavez and Araya, 2010), Uganda (Kashaija *et al.*, 1994), Ecuador (Chavez and Araya, 2010), India (Srinivasan *et al.*, 2011), Greece (Tzortzakakis, 2008), Australia (Pattison, 2002, 2011), Hawaii (Wang and Hooks, 2009), Belize (Bridge, 1996), and Rwanda (Gaidashova *et al.*, 2004).

Previous studies by Quénéhervé (1989) and by Volker and Gamboa (1991) identified *R. similis*, *P. coffeae*, and *P. brachyurus* as the most damaging PPN to plantain. Additionally, species like *H. multicinctus* and *H. dishytera* were also damaging to plantain (Lamptey and Karikari, 1977; McSorley and Parrado, 1986; Kamira *et al.*, 2013). Roman (1986) also reported *Helicotylenchus* sp. and *Rotylenchus reniformis* on plantain. Root-associate and fungal-feeding nematodes, including *Tylenchus* sp., *Aphelenchus* sp., and *Aphelenchoides* sp., considered to be of minor importance in plantain production, were not considered further in this study. Although found in association with plantain cultivation, they were found at low frequencies and densities; therefore, they probably do not have economic importance in

the system. The results of all the genera reported in this study, except *R. reniformis*, are consistent with those by Gowen *et al.* (2005) and Roman (1986), which both reported that nematode parasitism in *Musa* is characterized by several nematode species causing simultaneous infections.

Based on the nematode population of *Helicotylenchus* sp. with 78 nematodes/100 g soil and *R. similis* with 9,520 nematodes/100 g root, these were the most abundant PPN in plantain fields in the Talamanca canton region. *Helicotylenchus* and *R. similis* are diverse pests of *Musa* (Gowen and Quénéhervé, 1990) causing considerable damage (Kashaija *et al.*, 1994; Fogain and Gowen 1997; Elsen *et al.*, 2000; Barekye *et al.*, 2005). These findings are similar to those obtained by Marin *et al.* (1998) and Castillo-Russi *et al.* (2010) where these two genera were more abundant and widespread than *Pratylenchus*. The low frequency and abundance of *Pratylenchus* sp. could be attributed to its low competitiveness and suppression by high densities of *R. similis* since both nematodes have similar feeding strategies. However, the frequency and abundance of each of these nematodes can change depending on the plantain variety and the agroecological conditions during the year (Araya, 2003).

Nematodes like *R. similis* and *H. multicinctus* are of enormous concern among the many plant-parasitic nematodes associated with *Musa* (Wang and Hooks, 2009), and they usually occur together in plantain roots (Gowen, 1993). *Radopholus similis* can significantly weaken root systems, reduce yields, topple plants before harvest, reduce

fertilizer uptake and utilization, and make plants more prone to wind knockdowns. Nelson *et al.* (2006) indicated that *R. similis* can cause lesions to root and rhizomes, plantain decline, yield losses, and toppling; the nematode has a relatively wide non-*Musa* host range. Brooks (2004) reported that *Helicotylenchus* sp. is a worldwide problem in *Musa*-growing regions causing yield losses of 40% and 30-60% in Africa and India, respectively. It is possible that the abundance of weeds associated with plantain plantations is a reservoir of PPN around plantain in the studied area. This condition could increase the presence of these PPN in the plantations.

López (1980) determined the presence of *R. similis* in Costa Rica, along with some species of *Helicotylenchus*, *Meloidogyne*, and *Pratylenchus*. However, lower population densities of *H. multicinctus* were observed associated with the presence of high population densities of *R. similis* that may have impeded its development. In this investigation, *Helicotylenchus* sp. was found in 100% of the root samples. We believe this is an important finding and merits future research related to the damage caused by this nematode to plantain. This study focused on determining the nematodes associated with plantain cultivation only. According to Sikora *et al.* (1989) and Gowen and Quénehervé (1990), there are nematodes with different feeding habits that generally exist as mixed populations in *Musa* fields. We support this view because, in a tropical country such as Costa Rica, it is easy to find a mixture of species in a single crop such as plantain.

Previous research (López, 1980; Araya *et al.*, 2002; Araya, 2002, 2003, 2005; Chavez and Araya, 2010) supports the relative importance of *R. similis* as a *Musa* pest in Costa Rica. For example, the high population densities and frequencies of *R. similis* in many plantain fields in Sixaola, Talamanca can be attributed to the long-term monoculture practices that lead to ideal conditions for nematode survival and much variation in nematode populations. However, in Alto Talamanca (indigenous zone), the management practices are different because agrochemicals are not used, and plantain is grown in association with timber and fruit trees. Nevertheless, we also found high population densities of PPN. We believe the introduction of contaminated plantain plants could have also contributed to the high diversity of nematodes found in the fields. It is also important to

investigate if weeds growing in *Musa* plantations are hosts of the identified nematodes, and whether they are of concern for plantain cultivation. In this sense, López (1980) found that weeds such as *Croton trinitatis*, *Stachytarpheta* sp., and *Cyathula achyranthoides* are hosts of *M. javanica* and that *Laportea aestuans* and *Piper auritum* are of *M. incognita*. These weeds can be found frequently in plantain plantations and may be alternate hosts of PPN.

In conclusion, *R. similis*, *Pratylenchus* spp., *Helicotylenchus* spp., and *Meloidogyne* are the major PPN associated with plantain in Talamanca canton. Considering the destructive nature of these nematodes in various *Musa*-producing countries, research and management practices have to be in place to minimize crop damage as well as improve income and well-being from the small plantain producers of Talamanca. Future studies should evaluate the extent of plantain damage caused by PPN in Talamanca canton, followed by development of integrated pest management strategies against these nematodes. An alternative would be establishing nematode-free plantain planting material in fields not previously cultivated with plantain.

ACKNOWLEDGMENTS

Research was supported by grant of Universidad Nacional (UNA) through the Institutional Fund for Academic Development (FIDA).

LITERATURE CITED

- Araya, M. 2002. Metodología utilizada en el laboratorio de nematología de CORBANA S.A. para la extracción de nematodos de las raíces de banana (*Musa* AAA) y plátano (*Musa* AAB). CORBANA 28:97-110.
- Araya, M. 2003. Situación actual del manejo de nematodos en banano (*Musa* AAA) y plátano (*Musa* AAB) en el trópico americano. Pp. 79-102 *en* Rivas, G., y Rosales, F. (eds.), Manejo Convencional y alternativo de la Sigatoka Negra, Nematodos y Otras Plagas Asociadas al Cultivo de Musáceas en los Trópicos. Francia: INIBAP.
- Araya, M. 2005. Study of the effect of *Musa* genotype, soil type, and weed management on nematodes and nematode damage

- distribution. Doctoral Thesis No 655, Katholieke Universiteit Leuven, Leuven, Belgium.
- Araya, M., D., De Waele, and R. Vargas. 2002. Occurrence and population densities of nematode parasites of banana (*Musa AAA*) roots in Costa Rica. *Nematropica* 32:21-33.
- Barekye, A., I. N, Kashaija, W. K, Tushemereirwe, and E. Adipala. 2005. Comparison of damage levels caused by *Radopholus similis* and *Helicotylenchus multicinctus* on bananas in Uganda. *Annals of Applied Biology* 137:273-278.
- Barker, K. R. 1985. Sampling nematode communities. Pp. 3-17 in K. R. Barker, C. C. Carter, and J. N. Sasser (eds.) *An Advanced Treatise on Meloidogyne*. Volume II: Methodology. Raleigh, NC, USA: North Carolina State University Graphics.
- Bridge, J. 1996. Nematode management in sustainable and subsistence agriculture. *Annual Review of Phytopathology* 34:201-225.
- Brooks, F. 2004. Banana Nematodes. American Samoa Community College and Natural Resources Cooperative Research and Extension 9:1-3.
- Castillo-Russi, J. D., M, Araya-Vargas and L. F. Patiño-Hoyos. 2010. Respuesta a la aplicación de nematicida en banano en la zona de Urubá, Colombia. *Agronomía Mesoamericana* 21:307-317.
- Chávez, C., and M. Araya. 2010. Spatial-temporal distribution of plant-parasitic nematodes in banana (*Musa AAA*) plantations in Ecuador. *Journal of Applied Biosciences* 33:2057-2069.
- Elsen, A., P. R, Speijer, R, Swennen and D. Waele. 2000. Nematode species densities, root damage and yield of bananas (*Musa spp.*) in Uganda. *African Plant Protection* 6:31-36.
- FAO (Food and Agriculture Organization of the United Nations). 2003. FAOSTAT Statistics Database. Rome: Food and Agriculture organization. Available at <<http://faostat.fao.org>>.
- FAO (Food and Agriculture Organization of the United Nations). 2018. FAOSTAT. Top 10 Country Production of Plantains. Available at: http://www.fao.org/faostat/en/#rankings/countries_by_commodity.
- Fogain, R., and S. R. Gowen. 1997. Damage to roots of *Musa* cultivars by *Radopholus similis* with and without protection of nematicides. *Nematropica* 27:27-32.
- Frison, E., and S. Sharrock. 1999. The economic, social and nutritional importance of banana in the world. Pp. 21-35 in C. Picq, E. Foure, and E. A. Frison (eds.). *Proceedings of International symposium on bananas and food security*, Cameroon, Douala, 10-14 November 1998. IPGRI.
- Gaidashova, S. V., S, Okech, E, Van den Berg., M, Marais., C. M, Gatarayiha, and P. E. Ragama. 2004. Plant-parasitic nematodes in banana-based farming systems in Rwanda: Species profile, distribution and abundance. *African Plant Protection* 10:27-33.
- Gowen, S. R. 1993. Yield losses caused by nematodes on bananas and some management techniques appropriate to farmers' fields in Africa. Pp. 199-208 in S. Gold and B. Gemmill (eds.) *Biological and Integrated Control of Highland Banana and Plantain Pests and Diseases*. *Proceedings of a Research Coordination Meeting*, Cotonou, Benin (C). IITA, Ibadan, Nigeria, 12-14 November 1991.
- Gowen, S. R. and P. Quénéhervé. 1990. Nematode parasites of bananas, plantains and abaca. Pp. 431-460 in M. Luc, R. A. Sikora and J. Bridge (eds.) *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*. Wallingford, UK: CAB International.
- Gowen, S. R., P. Quénéhervé, and R. Fogain. 2005. Nematode Parasites of Bananas and Plantains. Pp. 611-643 in M. Luc, R. A. Sikora, and J. Bridge (eds.) *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*, 2nd edition. Wallingford, U.K.: CAB International.
- Holdridge, L. 1982. *Ecología basada en zona de vida*. Trad. del inglés por Jiménez, H. *Segunda reimpresión*. Instituto Interamericano de Cooperación para la Agricultura. San José, Costa Rica.
- Jenkins, W. R. 1964. A rapid centrifugal-flotation technique for separating nematodes from soil. *Plant Disease Reporter* 48:692.
- Jones, D.R. 2000. The genera *Musa* and *Ensete*. Pp. 1-36 in D. R. Jones (ed.). *Diseases of banana, abaca and enset*. Wallingford, U.K.: CAB International.
- Kamira M., S. Hauser, P. Van Asten, D. Coyne, and H. L. Talwana. 2013. Nematodos

- fitoparásitos asociados a los cultivos de banano y plátano en el este y oeste de la República Democrática del Congo. *Nematropica*, 43:216-225.
- Karamura, E. B., E. Frison, D. A. Karamura, and S. Sharrock. 1999. Banana production systems in eastern and southern Africa. Pp. 401-412 in C. Picq, E. Fouré, and E. Frison (eds.). *Bananas and Food Security. Proceedings of the International Symposium, Douala, Cameroon, INIBAP, Montpellier, 10-14 November 1998.*
- Kashaija, I. N., P. R. Speijer, C. S. Gold and S. R. Gowen. 1994. Occurrence, distribution and abundance of plant parasitic nematodes of bananas in Uganda. *African Crop Science Journal* 2:99-104.
- Lamprey, P., and S. K. Karikari. 1977. *Helicotylenchus dihystra* associated with banana and plantain in Ghana. *Acta Horticulturae* 53:291-292.
DOI: 10.17660/ActaHortic.1977.53.39
- López, R. 1980. Determinación de los nematodos fitoparásitos asociados al plátano (*Musa acuminata* * *M. balbisiana*, AAB) en Río Frío. *Agronomía Costarricense* 4:143-147.
- McSorley, R., and J. L. Parrado. 1986. *Helicotylenchus multicinctus* on bananas: An international problem. *Nematropica* 16:73-91.
- Mai, W. F., P. G. Mullin, H. H. Lyon, and K. Loeffler. 1996. Plant parasitic nematodes. A pictorial key to genera. Fifth edition. Comstock Publishing Associates, Cornell University Press.
- Marín, D. H., T. B. Sutton, and K. R. Barker. 1998. Dissemination of bananas in Latin America and the Caribbean and its relationship to the occurrence of *Radopholus similis*. *Plant Disease* 82:964-974.
- Mora, R. S., y B. Y. Quirós. (2018). Boletín Estadístico Agropecuario №28 Serie Cronológica 2014 – 2017. SEPSA (Secretaría Ejecutiva de Planificación Sectorial Agropecuaria).
- Nelson, S. C., R. C. Ploetz, and A. K. Kepler. 2006. *Musa* species (banana and plantains), ver. 2.2. Pp. 1-33 in C. R. Elevitch (ed.). *Species profiles for Pacific Island Agroforestry: Permanent Agriculture Resources (PAR), Hôlualoa, Hawaii.*
<www.traditionaltree.org>
- Oladejo J. A., and W. A. Sanusi. 2008. Marketing analysis of plantain in Owo and Ose local government areas of Ondo State, Nigeria. *International Journal of Agriculture and Economic Rural Development* 1:93-101.
- Pattison, A. B. 2011. The importance of the antagonistic potential in the management of populations of plant-parasitic nematodes in banana (*Musa* AAA) as influenced by agronomic factors. Ph.D. Thesis, Rheinischen Friedrich-Wilhelms-Universität, Bonn, Germany.
- Pattison, A. B., J. M. Stanton, J. A. Cobon and V. J. Doogan. 2002. Population dynamics and economic threshold of the nematodes *Radopholus similis* and *Pratylenchus goodeyi* on banana in Australia. *International Journal of Pest Management* 48:107-111.
- Phillip, B., A. M. Shittu. I. O. Aiyelaagbe, and T. Addokun. 2009. Economic potentials of plantain and fluted pumpkin intercropping as a poverty reduction strategy in South-Western Nigeria. *World Journal of Agriculture Sciences* 5:525-534.
- Quénéhervé, P. 1989. Population of nematodes in soils of banana cv Poyo, in the Ivory Coast. 3. Seasonal dynamics of population in mineral soil. *Revue Nematology* 12:149-160.
- Quénéhervé, P. 1990. Spatial arrangement of nematodes around the banana plant in the Ivory Coast: Related comments on the interaction among concomitant phytophagous nematodes. *Acta Ecologica* 11:875-886.
- Sánchez, O., and M. Zúñiga. 2001. Informe estadístico de Exportaciones Bananeras 1970-2000. Dirección de Estadística y Política Bananera. Costa Rica. CORBANA, S.A., Costa Rica.
- Seinhorst, J. W. 1959. A rapid method for the transfer of nematodes from fixative to anhydrous glycerin. *Nematologica* 4:67-69.
- Seinhorst, J. W. 1962. On the killing, fixation, and transferring to glycerin of nematodes. *Nematologica* 8:29-32.
- Sikora R. A., N. D. Bafokuzara, A. S. S. Mbwana, G.W. Oloo, B. Uroni, and K. V. Seshu Reddy. 1989. Interrelationship between banana weevil, root lesion nematode and agronomic practices and their importance for banana decline. *FAO Plant Protection Bulletin* 37:151-157.
- Srinivasan, R., S. Kulothungan, P. Sundararaju, and C. Govindasamy. 2011. Biodiversity of plant parasitic nematodes associated with

- banana in Thanjavur district of Tamil Nadu. *International Journal of Plant, Animal and Environmental Sciences* 1:63-69.
- Thorne, G. 1961. *Principles of nematology*. New York: McGraw-Hill Book Company, Inc.
- Tzortzakakis, E. A. 2008. Plant parasitic nematodes associated with banana crop in Crete, Greece. *Acta Agriculturae Slovenica* 91:97-101.
- Volkers, E. E. L., and A. Gamboa. 1991. Nematodes on plantain. *Nematologica* 60: 98-99.
- Wang, K.-H., and C. R. R. Hooks. 2009. Plant-parasitic nematodes and their associated natural enemies within banana (*Musa* spp.) plantings in Hawaii. *Nematropica* 39:57-73.

Received:

28/IV/2020

Accepted for publication:

6/VIII/2020

Recibido:

Aceptado para publicación: