

## SCIENTIFIC NOTE: *LAGERSTROEMIA SPECIOSA* (L.) PERS. (LYTHRACEAE), A NEW HOST FOR THE DEFOLIATOR *OIKETICUS KIRBYI* GULDING, [1827] (LEPIDOPTERA: PSYCHIDAE)

**Wagner S. Tavares<sup>1</sup>, José E. Serrão<sup>2</sup>, Rodolfo A. Barbosa<sup>3</sup>, and José C. Zanuncio<sup>4</sup>**

<sup>1</sup>. Departamento de Fitotecnia, Universidade Federal de Viçosa, 36570-000, Viçosa, Minas Gerais, Brazil; wagner.tavares@ufv.br; <sup>2</sup>. Departamento de Biologia Geral, Universidade Federal de Viçosa, 36571-000, Viçosa, Minas Gerais, Brazil; jeserrao@ufv.br; <sup>3</sup>. Departamento de Engenharia Florestal, Universidade Federal de Viçosa, 36571-000, Viçosa, Minas Gerais, Brazil; rodolfo.barbosa@ufv.br; <sup>4</sup>. Departamento de Biologia Animal, Universidade Federal de Viçosa, 36571-000, Viçosa, Minas Gerais, Brazil; zanuncio@ufv.br

**Abstract** - The native vegetation of the Zona da Mata region in Minas Gerais, Brazil has largely been replaced by coffee plantations. Coffee, *Coffea* spp. (L.) (Rubiaceae), is the preferred host of the bagworm *Oiketicus kirbyi* Guilding, [1827] (Lepidoptera: Psychidae) in Brazil. This insect was found to cause damage to plants of *Lagerstroemia speciosa* (L.) Pers. (Lythraceae) in the Municipality of Viçosa, Minas Gerais, Brazil. Three individuals of various sizes were found in seven trees planted along a parking lot. The branches containing the insects were collected, brought to the laboratory, and transferred to plastic pots (1.0 L) provided with a cotton wool pad moistened with distilled water. The caterpillars, of nocturnal habit, were observed partly emerging from the larval cases to feed on leaves. The biggest larval case was opened, which contained a caterpillar that was identified and deposited in the Universidade Federal de Viçosa (UFV) collections. This is the first report of *O. kirbyi* defoliating *L. speciosa*, a plant grown in parking lot and gardens to provide shade and as an ornamental plant for flower production. *Oiketicus kirbyi* should be monitored on *L. speciosa* to determine its potential economic impact on this plant.

**Key words:** bagworm, giant crape-myrtle, ornamental plant, tropical region, Zona da Mata region

### INTRODUCTION

The Zona da Mata region is located in the southeast of Minas Gerais, Brazil, covering an area of 36,058 km<sup>2</sup>, accounting for 6.2 % of this state (Resende et al., 2009). Historically, its native vegetation was tropical forest, with dense tree cover, and represented part of the Atlantic Forest region (Portugal et al., 2010). These forests were reduced to small patches and shrubs on steep slopes, and were replaced over time by coffee plantations [*Coffea* spp. (L.) (Rubiaceae)], pastures (Poaceae) and other crops (Souza et al., 2009). The terrain of the Zona da Mata region is uneven, with many hills and narrow valleys and some mountains formed of ancient crystalline rocks. The altitude ranges from 100 m in the Rio Pomba and Paraíba do Sul valleys to 1,889 m in the most mountainous region (Nunes et al., 2009). The tropical climate has hot summers with temperatures reaching 25 °C and lower temperatures at altitudes above 1,000 m above sea level (Santana et al., 2010). The rainfall varies from 1,200 to 1,400 mm (Reis et al., 2007).

Giant Crape-myrtle, *Lagerstroemia speciosa* (L.) Pers. (Lythraceae) is native to tropical southern Asia (Ellis et al., 2007), being cultivated in southern Asia, India and Philippines and in tropical and subtropical regions as an ornamental plant (Hashim & Hughes, 2010). The attractiveness of *L. speciosa*, with numerous panicles and each flower with six white to purple petals and foliage suitable for shade in parking lots, has increased its cultivation in cities in the Zona da Mata region. In addition, pruned branches of this plant can be composted and used as fertilizer (Sannigrahi, 2009). This plant is also valued for its medicinal qualities. Extract of *L. speciosa* leaves is used as a natural health supplement and may support blood sugar balance and aid weight loss. The leaves of the *L. speciosa* and other parts are widely used in the Philippines, Taiwan, and Japan as a tea preparation. This tea is largely consumed because of its various medicinal properties, especially regarding the kidney (e.g. for dissolving kidney stones, for kidney cleansing,

and kidney health in general) (Unno et al., 2000). Moreover, improved varieties of *L. speciosa* have flowers with higher durability, resistance to pests and to environmental factors. However, this plant can be damaged by sucking and chewing pests, such as aphid *Sarucallis kahawaluokalani* Kirkaldy, [1907] (Hemiptera: Aphididae), beetles *Altica litigata* Fall, [1910] (Coleoptera: Chrysomelidae) and diseases (Pettis & Braman, 2007; Herbert et al., 2009).

Caterpillars of *Oiketicus kirbyi* Guilding, [1827] (Lepidoptera: Psychidae) build a self-enclosing bag or case with silk and pieces of plants for protection (Rhainds et al., 2009). In tropical America this insect has been found to cause significant damage to coffee leaves, but also feeds on plants of the Aquifoliaceae, Apocynaceae, Arecaceae, Betulaceae, Bignoniaceae, Caprifoliaceae, Casuarinaceae, Clusiaceae, Combretaceae, Ebenaceae, Erythroxylaceae, Euphorbiaceae, Lamiaceae, Laureaceae, Fabaceae, Malpighiaceae, Malvaceae, Myrtaceae, Moraceae, Musaceae, Palmaceae, Pinaceae, Polygonaceae, Rhizophoraceae, Rosaceae, Rubiaceae, Rutaceae, Sapindaceae, Ulmaceae, Vochysiaceae, and Zygophyllaceae families (Stephens, 1962; Rhainds et al., 2002; Rhainds & La Rosa, 2010). In Brazil, it feeds on *Eriobotrya japonica* (Thunb.) Lindl. (Rosaceae), *Eucalyptus* spp. (Myrtaceae), *Tectona grandis* L. (Lamiaceae), and *Terminalia catappa* L. (Combretaceae) and is considered a polyphagous pest (Gara et al., 1990; Rhainds et al., 2008; Rhainds & Sadof, 2009).

The aim of this paper is to report damage by *O. kirbyi* on *L. speciosa* in the Zona da Mata region in Viçosa, Minas Gerais, Brazil.

### MATERIALS AND METHODS

Three larval cases of different sizes, made from silk and plant material (Figs. 1A and 1B), were found on March, 3, 2010 on a *L. speciosa* tree (Fig. 2) in a parking lot near the Departamento

Table 1. Host plants of *Oiketicus kirbyi*.

<b>Host plants</b>	<b>Family</b>	<b>Country</b>	<b>References</b>
<i>Artocarpus heterophyllus</i>	Moraceae	Cuba	Núñez, 2006
<i>Bactris gasipaes</i>	Arecaceae	Costa Rica	Rhainds et al., 1996
<i>Bauhinia</i> sp.	Fabaceae	U.S.A.	Jones & Parks, 1928
<i>Bauhinia variegata</i>	Fabaceae	Brazil	Garcia, 1999
<i>Bernardia myricaefolia</i>	Euphorbiaceae	U.S.A.	Davis, 1964
<i>Byrsinima sericea</i>	Malpighiaceae	Brazil	Flinte et al., 2006
<i>Calophyllum antillanum</i>	Clusiaceae	Cuba	Núñez, 2006
<i>Casuarina equisetifolia</i>	Casuarinaceae	Cuba	Núñez, 2006
<i>Celtis laevigata</i>	Ulmaceae	Cameroon	Davis, 1964
<i>Citrus</i> spp.	Rutaceae	Cuba	Núñez, 2006
<i>Citrus sinensis</i> var. Natal	Rutaceae	Brazil	Gravena & Almeida, 1982
<i>Citrus sinensis</i> var. Valênciâ	Rutaceae	Brazil	Gravena & Almeida, 1982
<i>Coccocloba uvifera</i>	Polygonaceae	Cuba	Núñez, 2006
<i>Cocos nucifera</i>	Arecaceae	Costa Rica	Rhainds et al., 1995a
<i>Coffea</i> spp.	Rubiaceae	Brazil	Gravena & Almeida, 1982
<i>Coffea arabica</i>	Rubiaceae	Cuba	Núñez, 2006
<i>Cupania americana</i>	Sapindaceae	Cuba	Núñez, 2006
<i>Diospyros texana</i>	Ebenaceae	U.S.A.	Davis, 1964
<i>Elaeis guineensis</i>	Arecaceae	Costa Rica	Rhainds et al., 1995b
<i>Eryobotrya japonica</i>	Rosaceae	Cameroon	Davis, 1964
<i>Erythroxylum tortuosum</i>	Erythroxylaceae	Brazil	Diniz & Morais
<i>Eucalyptus</i> spp.	Myrtaceae	Brazil	Arce et al., 1987
<i>Eucalyptus urophylla</i>	Myrtaceae	Brazil	Pereira et al., 2001
<i>Eucalyptus urograndis</i>	Myrtaceae	Brazil	Peres Filho et al., 2009
<i>Ilex paraguariensis</i>	Aquifoliaceae	Brazil	Iede & Machado, 1989
<i>Ixora coccinea</i>	Rubiaceae	Cuba	Núñez, 2006
<i>Kielmeyera coriacea</i>	Clusiaceae	Brazil	Diniz et al., 1999
<i>Leucaena leucocephala</i>	Fabaceae	U.S.A.	Davis, 1964
<i>Leucaena pulviflora</i>	Fabaceae	Cameroon	Davis, 1964
<i>Lonicera japonica</i>	Caprifoliaceae	Cameroon	Davis, 1964
<i>Mimosa pudica</i>	Fabaceae	U.S.A.	Jones & Parks, 1928
<i>Musa</i> spp.	Musaceae	Costa Rica	Stephens, 1962
<i>Musa AAB</i> var. Horn	Musaceae	Venezuela	Ramírez et al., 1999
<i>Musa paradisiaca</i>	Musaceae	Cuba	Núñez, 2006
<i>Ostrya</i> sp.	Betulaceae	U.S.A.	Jones & Parks, 1928
<i>Persea</i> sp.	Lauraceae	Cuba	Núñez, 2006
<i>Persea</i> sp.	Lauraceae	Peru	Rhainds & La Rosa, 2010
<i>Persea americana</i>	Lauraceae	Cameroon	Davis, 1964
<i>Philodendron</i> sp.	Araceae	U.S.A.	Jones & Parks, 1928
<i>Porlieria angustifolia</i>	Zygophyllaceae	U.S.A.	Davis, 1964
<i>Prosopis glandulosa</i>	Fabaceae	Cameroon	Davis, 1964
<i>Psidium guajava</i>	Myrtaceae	Cuba	Núñez, 2006
<i>Pyracantha coccinea</i>	Rosaceae	Cameroon	Davis, 1964
<i>Qualea multiflora</i>	Vochysiaceae	Brazil	Diniz & Morais, 1997
<i>Robinia</i> sp.	Fabaceae	U.S.A.	Jones & Parks, 1928
<i>Spathodea campanulata</i>	Bignoniaceae	Brazil	Garcia, 1999
<i>Tabebuia rosea</i>	Bignoniaceae	Colombia	Madrigal, 1986
<i>Tectona grandis</i>	Lamiaceae	Brazil	Ferreira et al., 2008
<i>Terminalia</i> sp.	Combretaceae	Cuba	Núñez, 2006
<i>Terminalia catappa</i>	Combretaceae	Brazil	Garcia, 1999
<i>Theobroma cacao</i>	Malvaceae	Trinidad	Kirkpatrick, 1953
<i>Thuja orientalis</i>	Pinaceae	Cameroon	Davis, 1964
<i>Rhizophora</i> spp.	Rhizophoraceae	Ecuador	Gara et al., 1990
<i>Rosa</i> sp.	Rosaceae	Cameroon	Davis, 1964
<i>Ulmus</i> sp.	Ulmaceae	U.S.A.	Jones & Parks, 1928
<i>Vinca</i> sp.	Apocynaceae	U.S.A.	Jones & Parks, 1928



Fig. 1. *Oiketicus kirbyi*: case of late instar larva (A) and smaller case of young larva (B). Zona da Mata region in Viçosa, Minas Gerais, Brazil.



Fig. 2. *Lagerstroemia speciosa*, host plant.

de Entomologia of Universidade Federal de Viçosa (UFV) in Viçosa, Minas Gerais, Brazil ( $20^{\circ} 45' S$ ,  $42^{\circ} 51' W$ , 651 m). Seven trees up to 4 years in age and 6 m tall were searched. This area has little sharp relief, fragments of secondary forest and diversified fauna (Pereira et al., 2009a, b, 2010).

Branches of *L. speciosa* with cases were removed, brought to the Laboratório de Controle Biológico de Insetos (LCBI) of the Departamento de Biologia Animal (DBA) of UFV and transferred to plastic pots (1.0 L) with cut ends placed in cotton wool pads moistened with distilled water. The material was maintained at a temperature of  $25 \pm 1^{\circ} C$ , 12 h photoperiod and relative air humidity of  $70 \pm 10\%$ . The caterpillars, of nocturnal habit, partly emerged from the cases to feed on leaves of *L. speciosa*. Caterpillars of the two smaller cases died and were discarded. The biggest case was opened and was found to contain a caterpillar that was preserved and identified by comparison with material deposited in the UFV collections and with the published account by Davis (1964). Taxonomist Dr. Paulo Sérgio Fiúza Ferreira of UFV identified the material collected from *L. speciosa* (caterpillars and their larval cases). Characteristics of length, width, shape, silk, and pieces of plant material used in the tree cases were compared, and it was concluded that they belonged to the same species. Morphological aspects of the head capsule, lateral stigmata, spiracles, thoracic legs, prolegs and crochets, besides the length and color of caterpillar, were used for species identification.

## RESULTS AND DISCUSSION

Fragments of plant material in the cases as opposed to silk only or other materials are characteristics of the macro-moth family Psychidae (Davis, 1964). The three larval cases collected were of this type and thus recognized as Psychidae. The older case was 85 mm long and had a diameter of 27.5 mm. This case was heavily constructed of short sections of twigs and a few tiny leaf fragments arranged in circular pattern around the bag. A thin layer of grayish-brown silk covered the entire structure, concealing all but the outline of the plant fragments.

The bigger caterpillar had a grayish color and was 47 mm long, with black spots of irregular size and shape located on the dorsum of the three thoracic segments and head, suggesting it was a female (Davis, 1964). These spots are possibly for camouflage, since they only appear on the parts exposed during feeding, locomotion, and during construction of cases (Davis, 1964). The thoracic legs were well-developed in contrast to the abdominal and anal prolegs. Characteristics of the head capsule, lateral stigmata, spiracles, thoracic legs, prolegs and crochets along with length and color were compared with determined material in UFV and the descriptive account by Davis (1964) in order to confirm the identity of the species.

Based on the aforementioned characters, the caterpillar found on *L. speciosa* was identified as the bagworm *O. kirbyi*. The association of *O. kirbyi* with this plant adds to the more than 50 host plants previously known for this polyphagous species (Table 1) and confirms its Neotropical distribution. Cultivars of *Lagerstroemia* spp. were found to be resistant to damage caused by the aphid *S. kahawaluokalani* (Herbert et al., 2009). Our observations on damage caused by this insect suggest it

has the potential to become a pest of *L. speciosa*. This is a bold conclusion based on finding only three larvae. However, further studies with other cultivars are needed.

The occurrence of *O. kirbyi* in Viçosa can be explained by the large area of coffee plantations in this municipality, since *Coffea* spp. is the preferred host for this insect (Ellis et al., 2007). Moreover, the egg stage of *O. kirbyi* is of shorter duration in tropical regions, indicating its adaptation to this region and resulting in greater fecundity in the Neotropical region (Mishra, 1978). The species is less common in temperate regions (Morden & Waldbauer, 1980) where it hibernates in the egg stage (Neal et al., 1987). *Oiketicus kirbyi* caterpillars draw most nutrients from leaves of plants with high pH levels (Rhainds et al., 2009). Although the chemical composition of plants may vary depending on environmental factors, which may prevent or reduce insect damage in some seasons (Sidhu et al., 2004), a comparative study of host pH levels and relative growth of *O. kirbyi* on *L. speciosa* and other potential ornamental hosts could be useful for the evaluation of the potential for pest damage.

This is the first report of *O. kirbyi* defoliating *L. speciosa*. We suggest *Lagerstroemia speciosa* should be monitored for *O. kirbyi* presence on a permanent basis. Its potential for defoliation should be evaluated and control measures developed, if necessary, to minimize the impact to these ornamental plants.

#### ACKNOWLEDGEMENTS

We acknowledge Don Davis for the review and suggestions concerning Table 1, Thomas Sobczyk for the review and suggestions concerning the manuscript, Santin Gravena for the review of the manuscript and providing references, Shuhei Niitsu and Mônica Ruediger de Britto Velho for editing the manuscript. The authors thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG) for financial support.

#### REFERENCES CITED

- Arce, J. J. C., O. Peres Filho, and E. Berti Filho**  
1987. Biología do bicho do cesto *Oiketicus kirbyi* (Lands.-Guilding, 1827) (Lepidoptera, Psychidae) em folhas de *Eucalyptus* spp. *Anais da Escola Superior de Agricultura Luiz de Queiroz* 44: 341-358.
- Basri, M. W., K. Norman, and A. B. Hamdan**  
1995. Natural enemies of the bagworm, *Metisa plana* Walker (Lepidoptera: Psychidae) and their impact on host population regulation. *Crop Protection* 14: 637-644.
- Cox, D. L. and D. A. Potter**  
1988. Within-crown distributions of male and female bagworm (Lepidoptera: Psychidae) pupae on juniper as affected by host defoliation. *Canadian Entomologist* 120: 559-567.
- Davis, D. R.**  
1964. Bagworm moths of the Western Hemisphere. *Bulletin of the United States National Museum* 244: 1-233.
- Diniz, I. R. and H. C. Morais**  
1997. Lepidopteran caterpillar fauna of cerrado host plant. *Biodiversity and Conservation* 6: 817-836.
- Diniz, I. R., H. C. Morais, A. M. F. Botelho, F. Venturoli, and B. C. Cabral**  
1999. Lepidopteran caterpillar fauna on lactiferous host plants in the central Brazilian cerrado. *Revista Brasileira de Biologia* 59: 627-635.
- Ellis, R. H., T. Mai-Hong, T. D. Hong, T. T. Tan, N. D. Xuan-Chuong, L. Q. Hung, B. Ngoc-Tam, and V. T. Le-Tam**  
2007. Comparative analysis by protocol and key of seed storage behaviour of sixty Vietnamese tree species. *Seed Science and Technology* 35: 460-476.
- Entwistle, P. F.**  
1963. Observations on the biology of four species of Psychidae (Lepidoptera) on *Theobroma cacao* L. in western Nigeria. *Proceedings of the Royal Entomological Society of London. Series A, General Entomology* 38: 145-152.
- Ferreira, R. A., W. F. G. Tosta, V. G. Giacometti, G. O. Souza, and J. M. S. Silva**  
2008. Entomofauna observada na cultura da teca (*Tectona grandis* L.f), no campo. *Revista Científica Eletrônica de Engenharia Florestal* 12: 1-23.
- Flinte, V., C. O. Araújo, M. V. Macedo, and R. F. Monteiro**  
2006. Insetos fitófagos associados ao murici da praia, *Byrsonima sericea* (Malpighiaceae), na Restinga de Jurubatiba (RJ). *Revista Brasileira de Entomologia* 50: 512-523.
- Gara, R. I., A. Sarango, and P. G. Cannon**  
1990. Defoliation of an Ecuadorian mangrove forest by the bagworm *Oiketicus kirbyi* Guilding (Lepidoptera: Psychidae). *Journal of Tropical Forest Science* 3: 181-186.
- Garcia, A. H.**  
1999. Levantamento, identificação e avaliação dos danos de insetos em árvores ornamentais na área urbana de Goiânia (GO). *Pesquisa Agropecuária Tropical* 29: 77-81.
- Gravena, S. and J. C. V. Almeida**  
1982. Inimigos naturais de *Oiketicus kirbyi* Lands Guilding, 1827 e *Oiketicus geyeri* Berg, 1877 no agroecossistema citrícola. *Científica* 10: 99-104.
- Hashim, N. R. and F. M. R. Hughes**  
2010. The responses of secondary forest tree seedlings to soil enrichment in Peninsular Malaysia: an experimental approach. *Tropical Ecology* 51: 173-182.
- Herbert, J. J., R. F. Mizell, and H. J. Mcauslane**  
2009. Host preference of the crapemyrtle aphid (Hemiptera: Aphididae) and host suitability of crapemyrtle cultivars. *Environmental Entomology* 38: 1155-1160.
- Iede, E. T. and D. C. Machado**  
1989. Pragas da erva-mate (*Ilex paraguariensis* St Hill.) e seu controle. *Boletim de Pesquisa Florestal* 18/19: 51-60.
- Jones, F. M. and H. B. Parks**  
1928. The bagworms of Texas. *Texas Agricultural Experiment Station Bulletin* 382: 1-36.
- Kirkpatrick, T. W.**  
1953. Notes on minor insect pests of cacao in Trinidad. Lepidoptera (butterflies and moths). *A Report on Cacao Research* 2: 67-72.
- Madrigal, A.**  
1986. Reconocimiento de insectos dañinos en plantaciones forestales de la Costa Atlántica Colombiana. *Miscelánea Sociedad Colombiana de Entomología* 12: 8-12.
- Mishra, S. C.**  
1978. Observations on the biology of bag worm: *Clania cramerii* Westwood (Psychidae: Lepidoptera). *Indian Journal of Forestry* 104: 135-141.
- Morden, R. D. and G. P. Waldbauer**  
1980. Diapause and its termination in the psychid moth, *Thyridopteryx ephemeraeformis*. *Entomologia Experimentalis et Applicata* 28: 322-333.
- Neal, J. W., M. J. Raupp, and L. W. Douglass**  
1987. Temperature-dependent model for predicting larval emergence of the bagworm, *Thyridopteryx ephemeraeformis* (Haworth) (Lepidoptera: Psychidae). *Environmental Entomology* 16: 1141-1144.
- Núñez, R.**  
2005. El gusano de cartucho mayor. *Boletín de la Sociedad cubana de Zoología* 1: 1-2.
- Nunes, L. A. P. L., L. E. Dias, I. Jucksch, N. F. Barros, M. C. M. Kasuya, and M. E. F. Correia**  
2009. Impact of monocultivation coffee on biological indicators of quality soil in the Zona da Mata (MG), Brazil. *Ciência Rural* 39: 2467-

2474.

- Pereira, J. M. M., T. V. Zanuncio, J. C. Zanuncio, and A. Pallini**  
2001. Lepidoptera pests collected in *Eucalyptus urophylla* (Myrtaceae) plantations during five years in Três Marias, State of Minas Gerais, Brazil. *Revista de Biologia Tropical* 49: 1073-1082.
- Pereira, A. I. A., J. C. Zanuncio, H. R. Gil-Santana, F. S. Ramalho, G. L. D. Leite, and J. E. Serrão**  
2009a. *Harpactor angulosus* (Reduviidae: Harpactorinae), a predator of Neotropical saturniids, *Hylesia* spp. in Brazil. *Entomological News* 120: 206-212.
- Pereira, A. I. A., C. R. S. Curvêlo, D. R. Smith, J. M. M. Pereira, T. V. Zanuncio, and J. C. Zanuncio**  
2010. Fatores de mortalidade natural do Symphyta Neotropical *Haplostegus nigricrus* (Hymenoptera: Pergidae). *Bioscience Journal* 26: 115-120.
- Pereira, A. I. A., V. V. Zanuncio, A. S. Lorenzon, H. Bolognani, B. V. Fernandes, O. H. H. Mielke, J. E. Serrão, and J. C. Zanuncio**  
2009b. Biological and morphological characteristics of *Hylesia paulex* (Lepidoptera: Saturniidae) fed with *Eucalyptus urophylla* (Myrtaceae). *Interciencia* 34: 645-649.
- Peres Filho, O., R. S. Gusmão, A. B. S. Januário, and A. Dorval**  
2009. Primeiro registro de *Oiketicus kirbyi* em clone de *Eucalyptus urophylla* x *Eucaliptus grandis* (*urograndis*) no estado de Mato Grosso, Brasil. *Revista de Agricultura* 84: 175-178.
- Pettis, G. V. and S. K. Braman**  
2007. Effect of temperature and host plant on survival and development of *Altica litigata* Fall. *Journal of Entomological Science* 42: 66-73.
- Portugal, A. F., O. D. V. Costa, and L. M. Costa**  
2010. Physical and chemical properties of a soil under different production systems and forest in the Zona da Mata region of Minas Gerais State (Brazil). *Revista Brasileira de Ciência do Solo* 34: 575-585.
- Ramírez, R., O. Domínguez, O. Liscano, M. Vilchez, R. Urdaneta**  
1999. Importancia de *Antichloris viridis* Druce como lepidóptero defoliador del plátano (*Musa* AAB cv. Hartón) en la zona sur del Lago de Maracaibo, Venezuela. *Revista de la Facultad de Agronomía* 16: 88-94.
- Reis, H., J. R. S. Scolforo, A. D. Oliveira, A. T. Oliveira, and J. M. Mello**  
2007. Floristic composition analysis, diversity and similarity of Atlantic forest fragments in Minas Gerais State-Brazil. *Cerne* 13: 280-290.
- Resende, J. L. P., C. T. J. Padua, A. D. Oliveira, J. R. S. Scolforo, and L. M. Coelho**  
2009. Characterization of a forest incentive program in Minas Gerais State. *Cerne* 15: 295-302.
- Rhainds, M. and C. T. Ho**  
2002. Size-dependent reproductive output of female bagworms (Lepidoptera: Psychidae): implications for intergenerational variations of population density. *Applied Entomology and Zoology* 37: 357-364.
- Rhainds, M. and C. S. Sadof**  
2009. Fitness consequence of pupation site for bagworms (Lepidoptera: Psychidae) on arborvitae. *Applied Entomology and Zoology* 44: 323-329.
- Rhainds, M. and J. C. C. La Rosa**  
2010. *Oiketicus kirbyi* (Lepidoptera, Psychidae), a key pest in Peruvian orchards of avocado. *International Journal of Pest Management* 56: 103-107.

**Rhainds, M., G. Gries, and G. Castrillo**1995a. Pupation site affects the mating success of small but not large female bagworms, *Oiketicus kirbyi* (Lepidoptera: Psychidae). *Oikos* 74: 213-217.**Rhainds, M., G. Gries, and C. Chinchilla**1995b. Pupation site and emergence time influence the mating success of bagworm females, *Oiketicus kirbyi*. *Entomologia Experimentalis et Applicata* 77: 183-187.**Rhainds, M., G. Gries, and C. Chinchilla**1996. Development of a sampling method for first instar *Oiketicus kirbyi* (Lepidoptera: Psychidae) in oil palm plantations. *Journal of Economic Entomology* 89: 396-401.**Rhainds, M., G. Gries, and M. M. Min**1999. Size- and density-dependent reproductive success of bagworms, *Metisa plana*. *Entomologia Experimentalis et Applicata* 91: 375-383.**Rhainds, M., S. R. Leather, and C. Sadof**2008. Polyphagy, flightlessness, and reproductive output of females: a case study with bagworms (Lepidoptera: Psychidae). *Ecological Entomology* 33: 663-672.**Rhainds, M., D. R. Davis, and P. W. Price**2009. Bionomics of Bagworms (Lepidoptera: Psychidae). *Annual Review of Entomology* 54: 209-226.**Rhainds, M., G. Gries, C. T. Ho, and P. S. Chew**2002. Dispersal by bagworm larvae, *Metisa plana*: effects of population density, larval sex, and host plant attributes. *Ecological Entomology* 27: 204-212.**Sannigrahi, A. K.**2009. Biodegradation of leaf litter of tree species in presence of cow dung and earthworms. *Indian Journal of Biotechnology* 8: 335-338.**Santana, S. W. J., R. Barros, J. B. Torres, and M. G. C. Gondim**2010. Thermal requirements of the coconut pest *Athelocoa subrufella* (Hulst) (Lepidoptera: Phycitidae). *Neotropical Entomology* 39: 181-186.**Sidhu, O. P., V. Kumar, and H. M. Behl**2004. Variability in triterpenoids (nimbin and salanin) composition of neem among different provenances of India. *Industrial Crops and Products* 19: 69-75.**Souza, R. M., N. Anjos, and J. C. Sorgato**2009. Occurrence of *Naupactus cervinus* (Boheman) in coffee plantation in the region of Zona da Mata mineira. *Ciência e Agrotecnologia* 33: 1967-1971.**Stephens, C. S.**1962. *Oiketicus kirbyi* (Lepidoptera: Psychidae): a pest of bananas in Costa Rica. *Journal of Economic Entomology* 55: 381-386.**Thompson, V. P., A. B. Nicotra, and M. J. Steinbauer**2001. Influence of previous frost damage on tree growth and insect herbivory of *Eucalyptus globulus* *globulus*. *Austral Ecology* 26: 489-499.**Unno, T., I. Sakane, and T. Kakuda**2000. Inhibition of xanthine oxidase by an aqueous extract of banaba leaves (*Lagerstroemia speciosa*). *Journal of the Japanese Society for Food Science and Technology* 47: 740-943.